

This is the peer reviewed version of the following article: Shirai, A., Watanabe, T. and Matsuki, H. (2017), Inactivation of foodborne pathogenic and spoilage micro-organisms using ultraviolet-A light in combination with ferulic acid. *Lett Appl Microbiol*, 64: 96–102. doi:10.1111/lam.12701, which has been published in final form at <https://doi.org/10.1111/lam.12701>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

Inactivation of foodborne pathogenic and spoilage microorganisms using ultraviolet-A light in combination with ferulic Acid

Journal:	<i>Applied Microbiology</i>
Manuscript ID	LAM-2016-2310.R1
Journal Name:	Letters in Applied Microbiology
Manuscript Type:	LAM - Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Shirai, Akihiro; Tokushima University, Bioresource Chemistry and Technology Watanabe, Takashi; Tokushima University, Biological Science and Technology Matsuki, Hitoshi; Tokushima University, Bioresource Chemistry and Technology
Key Words:	Disinfection, Biocontrol, Food preservation, Non-thermal processes, Post harvest

1 **Inactivation of foodborne pathogenic and spoilage microorganisms**
2 **using ultraviolet-A light in combination with ferulic Acid**

3 A. Shirai¹, T. Watanabe² and H. Matsuki¹

4 1 Department of Bioresource Chemistry and Technology, Graduate School of Bioscience and
5 Bioindustry, Tokushima University, Japan

6 2 Department of Biological Science and Technology, Faculty of Engineering, Tokushima
7 University, Japan

8 Running headline: Synergistic antimicrobial activity

9 Corresponding author: Akihiro Shirai, 2-1, Minamijosanjima-cho, Tokushima 770-8513,
10 Japan.

11 E-mail: a.shirai@tokushima-u.ac.jp

12

13

14

15

16

17

18

19

20 SIGNIFICANCE AND IMPACT OF THE STUDY (100 words)

21 Microbial contamination is one of the most serious problems for foods, fruit and sugar thick
22 juices. UV light is suitable for non-thermal decontamination of food products by inactivation
23 of contaminating microorganisms. However, UV-A exposure is insufficient for disinfection.
24 This study demonstrates that the combination of UV-A LED light (350-385 nm), which is not
25 hazardous to human eyes and skin, and ferulic acid (FA), a known phytochemical and food
26 additive, provides synergistic antimicrobial activity against foodborne pathogenic and spoilage
27 microorganisms. Therefore, FA addition to UV-A light treatment may be useful for
28 improvement of UV-A disinfection technology to prevent food deterioration.

29 ABSTRACT (200 WORDS)

30 The low energy of UV-A (315–400 nm) is insufficient for disinfection. To improve UV-A
31 disinfection technology, the effect of ferulic acid (FA) addition on inactivation by UV-A
32 light-emitting diode (LED) light (350–385 nm) was evaluated in the eliminating of
33 suspensions of various food spoilers and pathogens (seven bacteria and four fungi).
34 Photoantimicrobial assays were performed at FA concentrations below the MIC. The MIC of
35 the isomerized FA, consisting of 93% *cis*-form and 7% *trans*-form, was very similar to that of
36 the commercially available FA (*trans*-form). Irradiation with UV-A (1.0 J cm^{-2}) in the
37 presence of 100 mg l^{-1} FA resulted in enhanced reducing of all of the tested bacterial strains. A
38 combination of UV-A (10 J cm^{-2}) and 1000 mg l^{-1} FA resulted in enhanced reducing of
39 *Saccharomyces cerevisiae* and one of the tested filamentous fungi. These results demonstrated
40 that the combination of a short-term application of UV-A and FA at a low concentration
41 yielded synergistic enhancement of antimicrobial activity, especially against bacteria.

42 **Keywords:** ferulic acid, photoantimicrobial activity, UV-A, food spoilers, foodborne
43 pathogens

44

45 **Introduction**

46 UV-bactericidal technology, which produces no residual chemicals and has little influence on
47 the environment, is a convenient method for **disinfection** of gases, liquids, and solid surfaces.

48 In recent years, **a versatile UV source** has been provided by the development of light-emitting
49 diodes (LEDs), which yield constant illumination at a specific wavelength and do not contain
50 mercury. LEDs are also advantageous because of their durability and low heat generation
51 (Shin *et al.* 2016). A LED illuminating UV-C light (266 nm or 275 nm), which is defined as
52 radiation with wavelengths 200–280 nm and has traditionally been used as an effective
53 germicidal disinfectant, exhibited high bactericidal activity (Kim *et al.* 2016; Shin *et al.* 2016).

54 UV-C LED irradiation (266 nm; irradiance $4 \mu\text{W cm}^{-2}$) reduced the viability of *Escherichia*
55 *coli* O157:H7 by 6 log-units at 0.5 mJ cm^{-2} . UV-A (315–400 nm) has been also reported to
56 exhibit photobactericidal activity (Hamamoto *et al.* 2007). Notably, exposure to UV-A is not
57 hazardous to human eyes and skin; in contrast, UV-B and UV-C light (i.e. at wavelengths
58 $<315 \text{ nm}$) can induce sunburn, cell mutations, and cell death (Gruijl 2002). Moreover, in the

59 food industry, UV-C exposure is known to inactivate polyphenolic acids (chlorogenic acid and
60 phloridzin) that are abundant in apple juice, thus decreasing the antioxidative activity of this
61 product (Islam *et al.* 2016). These findings suggest that UV-A light may be suitable for
62 non-thermal decontamination of food products by inactivation of contaminating
63 **microorganisms**. However, UV-A light, being of lower energy than UV-C, exhibits lower
64 photobactericidal activity. To completely inactivate *E. coli* using UV-A LED irradiation alone

65 (365 nm), a 315-J cm⁻² fluence (at an irradiance of 70 mW cm⁻² for 75 min) was required
66 (Hamamoto *et al.* 2007).

67 To increase the inactivation efficiency of UV-A, the synergistic bactericidal activity of the
68 combination of UV-A light and organic substrates, a quaternary ammonium salt (Shirai *et al.*
69 2014), and natural compounds [coumaric acid and ferulic acid (FA)] and their
70 derivatives(Shirai *et al.* 2015a, 2015b), has been investigated in our laboratory. In those
71 reports, the addition of FA enhanced the photobactericidal activity of the UV-A such that the
72 combination of approximately 20 mg l⁻¹ FA with UV-A (irradiance 4.09 mW cm⁻², 30 min)
73 resulted in a >5-log decrease in the survival of *E. coli*.

74 FA, one of the most abundant phenolic acids in plants such as rice, wheat, barley, citrus
75 fruits, and tomatoes, occurs as esters conjugated covalently with mono-, di-, and
76 polysaccharides and with lignin of cell walls (Graf 1992; Paiva *et al.* 2013). It is a potential
77 therapeutic agent with demonstrated antioxidant, antimicrobial, hepatoprotective, and UV
78 protective activities (D Paiva *et al.* 2013). In Japan, FA is approved for use as an antioxidant
79 food additive (JFCRF 1996).

80 Microbial contamination is one of the most serious problems for foods, fruit and sugar
81 thick juices (Braun *et al.* 1999; Fleet 2007; Juste *et al.* 2008; Tribst *et al.* 2009). In those
82 reports, *Salmonella* are typical pathogenic bacteria associated with fruit juices.
83 *Alicyclobacillus* and *Sporolactobacillus* have been isolated from spoiled fruit juices and can
84 survive heat treatments by forming heat-tolerant spores. Yeasts affect negatively juice's flavor,
85 turbidity and odor. The microorganisms that we tested included two heat-tolerant filamentous
86 fungi (*Byssoschlamys fulva* and *Eupenicillium lapidosum*) and a third filamentous fungus,
87 *Cladosporium cladosporioides*, found in canned or bottled fruit. *E. coli* is commonly used as

88 an indicator organism. As an UV-A disinfection technology for applications in the food
89 industry, the goal of this study was to investigate the synergistic effect of FA on the
90 photoantimicrobial activity of UV-A (wavelength range 350–385 nm produced by a LED
91 source) using those multiple food spoilers and pathogens and to reveal enhancement of
92 photoantimicrobial activity by FA addition.

93 Results and Discussion

94 Antimicrobial activity of FA isomers

95 FA (*trans*-form) and isomerized FA, consisting of 93% *cis*-form and 7% *trans*-form, were
96 evaluated for antimicrobial activity on the basis of MIC assays against various microorganisms
97 (Table 1). The MICs for both FAs were 125 to 1000 mg l⁻¹ or >1000 mg l⁻¹ for the tested
98 microorganisms. The MICs of isomerized FA and *trans*-FA were very similar and no
99 significant differences were observed between the isomers.

100 >Table 1<

101 Other work has shown that *cis*-form phenyl compounds substituted with propenoic acid
102 moieties have notable biological properties. For instance, the antimicrobial activity and the
103 anti-invasive activity (against adenocarcinoma cells) of cinnamic acid depend on its
104 isomerization (Chen *et al.* 2011; Yen *et al.* 2011). The *cis*-isomer of cinnamic acid has a strong
105 bactericidal effect against a multidrug-resistant *Mycobacterium tuberculosis* at a concentration
106 two orders of magnitude lower than that observed for the *trans*-isomer (Chen *et al.* 2011).
107 Cinnamic acid acts by causing a disruption in the cell wall of the bacterium, resulting in a
108 wrinkled and rough colony phenotype in micrographs. Similarly, other phenolic acid
109 compounds (caffeic acid and gallic acid) have been shown to bind to (or be incorporated into)

110 the cytoplasmic membrane of *E. coli* and *S. aureus* (Nakamura *et al.* 2015). FA is isomerized
111 from the *trans*-form to the *cis*-form through phenoxy radical formation during UV exposure
112 (Graf 1992). Photoisomerization of FA to the *cis*-isomer could directly injure the cell
113 membrane (as *cis*-cinnamic acid does). By analogy to cinnamic acid, we expected that the
114 antimicrobial activity of FA would exhibit isomeric specificity, such that the inhibitory activity
115 of the *cis*-form of FA would be much higher than that of the *trans*-form. However, the MICs
116 of isomerized FA and *trans*-FA were very similar in our experiments, suggesting that the
117 antimicrobial activity of FA does not depend on isomerization. Therefore, the photo-induced
118 conversion of FA would have no effect on its photoantimicrobial activity when combined with
119 UV-A irradiation.

120 **Photoantimicrobial activity**

121 **Photoantimicrobial assays were performed at FA concentrations below its MIC.** Bactericidal
122 activity against *E. coli* was investigated by treating this organism with a combination of FA
123 and UV-A irradiation. A total UV-A fluence at 30 J cm^{-2} (58.3-min irradiation) in the absence
124 of FA resulted in a 2.82-log unit reduction in the viable cell count [Fig. 1(A)]. The addition of
125 FA to the suspension before UV-A irradiation enhanced bactericidal activity in a manner that
126 was dependent on FA dose and UV-A fluence. The combination of FA (at 50, 100, and 150
127 mg l^{-1}) with UV-A light [at 1.5 J cm^{-2} (2.92-min irradiation), 1.0 J cm^{-2} (1.95-min irradiation),
128 and 0.75 J cm^{-2} (1.46-min irradiation), respectively] yielded viable cell counts of $<10 \text{ CFU}$
129 ml^{-1} [Fig. 1(B)]. Compared with the bactericidal activity of UV-A at 3.0 J cm^{-2} fluence in the
130 absence of FA, **the combination of FA with UV-A yielded significant ($P < 0.01$) decreases in**
131 **the viable cell count.** Exposure to 150 mg l^{-1} FA for an equivalent time interval (5.83 min, the
132 incubation time used above to provide a fluence of 3.0 J cm^{-2} UV-A) in the absence of UV-A

133 (i.e., in the dark) had a much smaller antibacterial effect (0.37-log reduction in *E. coli* cell
134 density).

135 >Figure 1<

136 The efficacy of the combination of FA (100 mg l⁻¹) and UV-A (1.0 J cm⁻²) was tested with
137 six other bacterial strains (Fig. 2). Under these conditions, cell counts of *Salm. enterica*, *Staph.*
138 *aureus*, and *Sporolact. inulinus* decreased below the detection limit of 10 CFU ml⁻¹. Cell
139 counts of *B. cereus*, *K. rhizophila* and *A. acidoterrestris* decreased by 4.04-logs, 3.75-logs and
140 2.20-logs, respectively. In contrast, no significant decrease in viability was observed for any of
141 these strains (except for *Sporolact. inulinus*) when exposed to FA in the absence of UV-A (for
142 1.95 min, the incubation time used above to provide a fluence of 1.0 J cm⁻² UV-A), or to the
143 UV-A fluence in the absence of FA. For *Sporolact. inulinus*, a decrease in viable cells
144 (3.15-logs) was observed upon treatment with FA in the dark. Notably, the combined
145 treatments with UV-A and FA provided significant ($P < 0.01$) increases in bactericidal activity
146 compared with those with FA addition alone or UV-A irradiation alone.

147 >Figure 2<

148 In other work, Nakamura et al. (2015) showed that irradiation with short wavelength light
149 (380–420 nm; irradiance 0.26 W cm⁻²) in combination with caffeic acid (a polyphenol similar
150 to FA) killed each of four bacterial strains; notably, however, these effects required a fluence
151 of 78 J cm⁻² and a caffeic acid concentration of 1000 mg l⁻¹. In contrast, inactivation of
152 bacteria by the methods described in the present study required much lower light fluence and
153 reagent concentration. Potent inactivation (to microbial densities below 10 CFU ml⁻¹, the
154 lower limit of detection) of *E. coli* was achieved at total fluences of 1.0 J cm⁻² in combination
155 with a FA concentration of 100 mg l⁻¹. Similar enhancement of UV-A bactericidal efficacy

156 was also demonstrated for the other six tested bacterial strains. The bactericidal synergy of
157 UV-A and FA against some Gram-positive bacteria, including some *Bacillus*, *Kocuria* and
158 *Alicyclobacillus*, was lower than against Gram-negative bacteria. As shown in a previous
159 report by Nakamura et al. (2015), the affinity of FA, which is an analog of caffeic acid, to
160 Gram-positive bacteria should be lower than to Gram-negative bacteria. These results
161 demonstrated that the combination of UV-A and FA yielded synergistic enhancement of
162 bactericidal activity while using a short-term application of UV-A light (1.0 J cm^{-2}) in
163 combination with 100 mg l^{-1} FA. We infer that the high synergism of UV-A light with FA
164 reflects the production of phenoxy radicals that in turn leads to the production of hydrogen
165 peroxide as supported by the fact that photobactericidal activity is quenched by the addition of
166 catalase (Shirai *et al.* 2015b). In this work, microbial inactivation by the FA + UV-A regime
167 was assayed against vegetative cells. Future work on the synergistic efficacy will be needed to
168 investigate activity against spores of organisms like *B. subtilis*, *B. cereus*, *A. acidoterrestris*
169 and *Sporolact. inulinus*.

170 We investigated antifungal activity by testing the effect of the combination of FA (1000
171 mg l^{-1}) and UV-A (10 J cm^{-2} fluence) on *S. cerevisiae* and three filamentous fungi.

172 UV-A irradiation alone, at fluences of up to 10 J cm^{-2} , yielded little decrease in viable
173 yeast cell counts (Fig. 3). Exposure to 1000 mg l^{-1} FA for an equivalent time interval (19.4 min,
174 the incubation time used above to provide a fluence of 10 J cm^{-1} UV-A) in the absence of
175 UV-A (i.e., in the dark) resulted in a very small amount of anti-yeast activity (0.05-log
176 reduction in yeast cell density). The anti-yeast activity of UV-A was enhanced in the presence
177 of 1000 mg l^{-1} FA; at a fluence of 10 J cm^{-2} in the presence of 1000 mg l^{-1} FA, yeast viability
178 fell below the detection limit of 10 CFU ml^{-1} .

179

>Figure 3<

180 The photoantifungal activity against *B. fulva* was significantly increased in the presence of
181 FA, with **survival** ratios without and with FA of **0.01-logs** and 1.49-logs, respectively ($P <$
182 0.01). In contrast, FA addition did not significantly enhance photoinactivation of *C.*
183 *cladosporioides* by UV-A irradiation ($P > 0.05$), though the treatment yielded a nominal
184 decrease in viability. Similarly, FA addition did not significantly enhance photoinactivation of
185 *E. lapidosum*. For the three filamentous fungi, FA addition alone and UV-A irradiation alone
186 had a low **fungicidal** activity. Additional investigations with various combinations of
187 irradiance and FA concentration may reveal conditions suitable for the **inactivation** of those
188 fungi; such efficacy would be of great value, given that these organisms are often resistant to
189 heating and UV-C exposure (254 nm) (Hamanaka *et al.* 2010).

190 Several laboratories have shown that UV-A LED light exhibits much lower
191 photobactericidal activity (Hamanoto *et al.* 2007; Shirai *et al.* 2014) than UV-C LED light
192 (Kim *et al.* 2016). Inactivation with UV-A light alone requires high irradiation and long
193 exposure times. In the present work, to increase the inactivation efficiency of UV-A, we
194 investigated the FA + UV-A combination with respect to its photoantimicrobial activity
195 against various foodborne pathogenic and spoilage microorganisms related to foods, fruit and
196 sugar thick juices.

197 In conclusion, a combination of FA with UV-A irradiation resulted in a significant
198 enhancement in their individual efficacy toward the tested organisms. The high synergistic
199 activity against bacteria was observed when UV-A at low fluence and short time exposure was
200 combined with FA at a low concentration. **Reducing** of yeast and at least one filamentous
201 fungus was observed upon treatment with longer UV-A exposure and higher FA concentration.

202 Therefore, FA combined with UV-A light treatment may be useful for the improvement of
203 UV-A disinfection technology. Also, in future work, surface decontamination of fruits will be
204 investigated using the combination of UV-A light and FA for applications in postharvest
205 disinfection.

206 **Materials and Methods**

207 **Microbial strains**

208 **Microorganisms** used in the determination of antimicrobial activity were purchased from NITE
209 Biological Resource Center (NBRC), American Type Culture Collection (ATCC) and Institute
210 of Food Microbiology (IFM). **Those strains are listed in Table 1.** Endospore-forming bacteria
211 (*Bacillus*, *A. acidoterrestris* and *Sporolact. inulinus*) were tested in their vegetative state.

212 **UV-A source and irradiation.**

213 A device equipped with a UV-A LED (NCSU033B; Nichia Corp., Anan, Japan), as previously
214 described, was used in all photoexperiments (Shirai *et al.* 2015b). The LED has a radiation
215 angle of about 120° as the full width at half maximum. The peak wavelength was 365 nm and
216 the wavelength range was 350–385 nm, which was measured with a cumulative UV meter
217 (MCPD-3700A; Otsuka Electronics Co. Ltd., Hirakata, Japan) (Fig. S1).

218 **Chemistry**

219 FA (PubChem CID: 445858) was purchased from Tokyo Chemical Industry Co. Ltd. (Tokyo,
220 Japan). Except as noted, FA used for testing consisted of the *trans*-form obtained as
221 commercially available FA.

222 **MIC assay**

223 Details regarding preparation, separation and purity of isomerized FA are summarized in the
224 Supporting Information (Text S1). The antimicrobial activity of FA and the isomerized FA,
225 consisting of 93% *cis*-form and 7% *trans*-form, against the organisms listed above was
226 evaluated by determining MICs using the broth dilution method (Shirai *et al.* 2005). Preculture
227 of organisms was performed with the conditions described in Table S1. Cell density was
228 determined from OD₆₆₀ using a UV-1700 spectrophotometer (Shimadzu Ltd., Kyoto, Japan)
229 for bacteria and yeast, and using a hemocytometer (Burker-Turk; depth 0.1 mm, 1/400 qmm;
230 Erma Inc., Tokyo, Japan) for filamentous fungi.

231 FA solutions for testing against bacteria (except *A. acidoterrestris* and *Sporolact. inulinus*)
232 were generated by diluting the FA stock solution [100 g l⁻¹ in 80% dimethylsulfoxide
233 (DMSO)] with nutrient broth (Becton, Dickinson and Company, Franklin Lakes, NJ, USA) to
234 generate the highest testing concentration of 1000 mg l⁻¹; this solution was then subjected to a
235 two-fold serial dilution using nutrient broth. To generate FA solutions for testing against *A.*
236 *acidoterrestris* and *Sporolact. inulinus*, all dilutions (starting from the stock solution) were
237 performed as above but using a specific broth (No. 323), which is recommended on the NBRC
238 website (NBRC 2016), and GYP broth [glucose 2% (w/v), yeast extract 0.5% (w/v), Bacto
239 peptone 0.5% (w/v)] (Kitahara and Suzuki 1963), respectively. To generate FA solutions for
240 testing against fungi, all dilutions (starting from the stock solution) were performed as above
241 but using Sabouraud broth [polypeptone 1% (w/v) and glucose 4% (w/v)]. The final cell
242 densities were approx. 1×10⁵ CFU ml⁻¹ for bacteria and yeast, and approx. 1×10⁴ conidia ml⁻¹
243 for filamentous fungi in a transparent 96-well culture plate (Corning Inc., NY, USA). MICs for
244 *A. acidoterrestris* and *Sporolact. inulinus* were determined after 24-h incubation at 45°C and
245 48-h incubation at 37°C in an anaerobic chamber with an AnaeroPack Kenki that can reduce

246 the oxygen percentage to <0.1% within 2 h (Mitsubishi Gas Chemical Company, Inc., Tokyo,
247 Japan), respectively. MICs for the other bacteria, yeast, and filamentous fungi were determined
248 after 24-h incubation in ambient air at 37°C, 28°C, and 25°C, respectively.

249 **Photoantimicrobial assay**

250 Photoantimicrobial activity against bacteria and fungi was determined by plating and counting
251 the colony-forming units remaining after treatment of a microbial suspension as reported
252 previously (Shirai *et al.* 2014). Cells of organisms, precultured according to Table S1, were
253 prepared as described in the same report. Conidial suspensions were prepared as described
254 previously (Shirai *et al.* 2005). Cell density was determined as described in the section above.

255 All antimicrobial assays with UV-A irradiation or with no irradiation (in the dark) were
256 performed in an incubator box maintained at 30°C, using suspensions of organisms placed in
257 the individual wells of a transparent 48-well culture plate (AGC Tecno Glass Co. Ltd., Tokyo,
258 Japan). An aliquot of microbial suspension (0.1 ml, approx. 2×10^7 CFU ml⁻¹ for bacteria or
259 approx. 2×10^5 CFU ml⁻¹ for fungi) was added to each well of a 48-well culture plate already
260 containing 0.01 ml of test compound at concentrations of 5, 10, 15, and 100 g l⁻¹ in 80%
261 DMSO and 0.89 ml of sterile water per well (volume of the final tested suspensions 1 ml;
262 depth 13 mm). For UV-A irradiation alone (no test compound), DMSO was added to a
263 concentration of 0.8% to each well of the 48-well plate. The single UV-A LED was placed
264 face-up to permit upward irradiation into the bottom of the 48-well culture plate. The device
265 was set 30 mm (height) from the middle of the tested suspensions. The intensity of was 8.58
266 mW cm⁻² at the bottom of the well, which was measured with a laser power and energy meter
267 (Nova II; Ophir Optronics Solutions Ltd., Saitama, Japan) equipped with a photodiode sensor
268 (PD-300-UV; Ophir Optronics Solutions Ltd.). Total fluence tested (0.5, 0.75, 1.0, 1.5, 3.0, 10,

269 and 30 J cm^{-2}) was calculated based on the irradiance (8.58 mW cm^{-2}) and exposure time (0.97
270 to 58.3 min). After treatments with or without UV-A irradiation for bacteria, aliquots (0.15 ml)
271 of each suspension were diluted 10-fold with SCDLP broth (Nihon Pharmaceutical Co., Ltd.,
272 Tokyo, Japan) followed by serial 10-fold dilutions with 0.8% (w/v) physiological saline
273 containing 0.7% (w/w) Tween 80 (Kanto Chemical Co., Inc., Tokyo, Japan). Viable cell
274 counts (CFU ml^{-1}) after 10-fold serial dilutions were determined by plating on suitable agar
275 plates and incubating them (Table S1). For fungi, 10-fold serial dilutions were performed with
276 Sabouraud broth containing 0.1% (w/w) Tween 80.

277 Antimicrobial activity was expressed as the log survival ratio (log S) according to the
278 equation: $\log S = \log(N_t/N_0)$, where N_0 represents the number of CFUs before bactericidal
279 treatment and N_t represents the number of CFUs after treatment for time t .

280 **Statistical analysis**

281 All antimicrobial experiments were performed as three independent procedures, and results are
282 presented as the mean and SD. Inferential analysis was performed using a two-tailed, unpaired
283 Student's t -test. P -values of <0.05 were considered significant.

284 **Acknowledgements**

285 This work was supported by a Grant-in-Aid for Scientific Research (C) from the Japan Society
286 for the Promotion of Science (No. 15K05632).

287 **Conflict of Interest**

288 No conflict of interest declared.

289 **References**

- 290 Braun, P., Fehlhaber, K., Klug, C. and Kopp, K. (1999) Investigations into the activity of
291 enzymes produced by spoilage-causing bacteria: a possible basis for improved shelf-life
292 estimation. *Food Microbiol* **16**, 531-540.
- 293 Chen, Y.-L., Huang, S.-T., Sun, F.-M., Chiang, Y.-L., Chiang, C.-J., Tsai, C.-M. and Weng,
294 C.-J. (2011) Transformation of cinnamic acid from *trans*- to *cis*-form raises a notable
295 bactericidal and synergistic activity against multiple-drug resistant *Mycobacterium*
296 *tuberculosis*. *Eur J Pharm Sci* **43**, 188-194.
- 297 De Gruijl, F.R. (2002) p53 mutations as a marker of skin cancer risk: comparison of UVA and
298 UVB effects. *Exp Dermatol* **11**, 37-39.
- 299 De Paiva, L.B., Goldbeck, R., Dos Santos, W.D. and Squina, F.M. (2013) Ferulic acid and
300 derivatives: molecules with potential application in the pharmaceutical field. *Braz J Pharm*
301 *Sci* **49**, 395-411.
- 302 Fleet, G.H. (2007) Yeasts in foods and beverages: impact on product quality and safety. *Curr*
303 *Opin Biotechnol* **18**, 170-175.
- 304 Graf, E. (1992) Antioxidant potential of ferulic acid. *Free Radic Biol Med* **13**, 435-448.
- 305 Hamamoto, A., Mori, M., Takahashi, A., Nakano, M., Wakikawa, N., Akutagawa, M., Ikehara,
306 T., Nakaya, Y. and Kinouchi, Y. (2007) New water disinfection system using UVA
307 light-emitting diodes. *J Appl Microbiol* **103**, 2291-2298.
- 308 Hamanaka, D., Atungulu, G.G., Tanaka, F. and Uchino T. (2010) Effect of combining infrared
309 heating with ultraviolet irradiation on inactivation of mold spores. *Food Sci Technol Res*
310 **16**, 279-284.

- 311 Islam, M.S., Patras, A., Pokharel, B., Wu, Y., Vergne, M.J., Shade, L., Xiao, H. and Sasges,
312 M. (2016) UV-C irradiation as an alternative disinfection technique: Study of its effect on
313 polyphenols and antioxidant activity of apple juice. *Innovative Food Sci Emerg Technol*
314 **34**, 344-351.
- 315 JFCRF, The Japan Food Chemical Research Foundation (1996) *List of Existing Food*
316 *Additives*. Available at:
317 <http://www.ffcr.or.jp/zaidan/FFCRHOME.nsf/7bd44c20b0dc562649256502001b65e9/346>
318 [b67d78f8905324925685c0027a6e0/\\$FILE/En2014.1.30.pdf](http://www.ffcr.or.jp/zaidan/FFCRHOME.nsf/7bd44c20b0dc562649256502001b65e9/346). Accessed on 26 Oct. 2016.
- 319 Justé, A., Lievens, B., Frans, I., Klingeberg, M., Michiels, C.W. and Willems, K.A. (2008)
320 Present knowledge of the bacterial microflora in the extreme environment of sugar thick
321 juice. *Food Microbiol* **25**, 831-836.
- 322 Kim, S.-J., Kim, D.-K. and Kang, D.-H. (2016) Inactivating foodborne pathogens by using
323 UVC-LEDs at wavelengths from 266 to 279 nm and application to pasteurize sliced
324 cheese. *Appl Environ Microbiol* **82**, 11-17.
- 325 Kitahara, K. and Suzuki, J. (1963) *Sporolactobacillus* nov. subgen. *J Gen Appl Microbiol* **9**,
326 59-71.
- 327 Nakamura, K., Ishiyama, K., Sheng, H., Ikai, H., Kanno, T. and Niwano, Y. (2015)
328 Bactericidal activity and mechanism of photoirradiated polyphenols against Gram-positive
329 and -negative bacteria. *J Agric Food Chem* **63**, 7707-7713.
- 330 NBRC, National Institute of Technology and Evaluation Biological Resource Center. *Medium*
331 *detail information*. Available at:

- 332 <http://www.nbrc.nite.go.jp/NBRC2/NBRCMediumDetailServlet?NO=323>. Accessed on 26
333 Oct. 2016.
- 334 Shin, J.-Y., Kim, S.-J., Kim, D.-K. and Kang D.-H. (2016) Fundamental characteristics of
335 deep-UV light-emitting diodes and their application to control foodborne pathogens. *Appl*
336 *Environ Microbiol* **82**, 2-10.
- 337 Shirai, A., Aihara, M., Takahashi, A., Maseda, H. and Omasa, T. (2014) Synergistic
338 antimicrobial activity based on the combined use of a gemini-quaternary ammonium
339 compound and ultraviolet-A light. *J Photochem Photobiol B*, **130**, 226-233.
- 340 Shirai, A., Kajiura, M., Matsumura, K. and Omasa, T. (2015a) Improved photobactericidal
341 activity of ultraviolet-A light in combination with isomerizable *p*-coumaric acid
342 derivatives. *Biocontrol Sci* **20**, 231-238.
- 343 Shirai, A., Kajiura, M. and Omasa, T. (2015b) Synergistic photobactericidal activity based on
344 ultraviolet-A irradiation and ferulic acid derivatives. *Photochem Photobiol* **91**, 1422-1428.
- 345 Shirai, A., Maeda, T., Nagamune, H., Matsuki, H., Kaneshina, S. and Kourai, H. (2005)
346 Biological and physicochemical properties of gemini quaternary ammonium compounds in
347 which the positions of a cross-linking sulfur in the spacer differ. *Eur J Med Chem* **40**,
348 113-123.
- 349 Tribst, A.A.L., De Souza Sant'Ana, A. and De Massaguer, P.R. (2009) Review:
350 microbiological quality and safety of fruit juices—past, present and future perspectives.
351 *Crit Rev Microbiol* **35**, 310-339.
- 352 Yen, G.-C., Chen, Y.-L., Sun, F.-M., Chiang, Y.-L., Lu, S.-H. and Weng, C.-J. (2011) A
353 comparative study on the effectiveness of *cis*- and *trans*-form of cinnamic acid treatments

354 for inhibiting invasive activity of human lung adenocarcinoma cells. *Eur J Pharm Sci* **44**,
355 281-287.

356 **Supporting Information**

357 Additional Supporting Information may be found in the online version of this article:

358 **Text S1.** Preparation, separation, and purity of *cis*-FA.

359 **Table S1.** Conditions for preculture and viability assay of **microorganisms** tested.

360 **Figure S1.** Emission spectrum of the UV-A LED as used in this study; the spectrum exhibited
361 a maximum at 365 nm, which was measured at a distance of 30 mm between the illumination
362 source and the UV meter.

363 **Figure S2.** HPLC chromatogram of FA prepared by UV-A exposure: before separation (A),
364 and after separation and purification of the *cis*-form (B).

365 **Figure S3.** UV-visible absorption spectra of FA (*trans*-form, solid line) and isomerized FA
366 (mixture of 93% *cis*-form and 7% *trans*-form, dashed line). Samples were dissolved at 50 μM
367 in 0.4% DMSO.

368 **FIGURE LEGENDS**

369 **Figure 1.** UV-A fluence-dependent changes in *E. coli* survival following irradiation in the
370 absence (A) or presence (B) of FA. Filled (black) symbols, no UV-A exposure; unfilled
371 (white) symbols, UV-A exposure. Cell suspensions were treated with FA at 50 mg l^{-1}
372 (diamond), 100 mg l^{-1} (triangle), or 150 mg l^{-1} (circle). Data are presented as means \pm SD (n =
373 3). Significant differences (** $P < 0.01$) were calculated based on comparison to samples

374 irradiated with UV-A (3.0 J cm^{-2}) in the absence of FA. Samples for which survival was <10
375 CFU ml^{-1} (lower limit of detection) are noted in parentheses as 6-log unit reductions.

376 **Figure 2.** Photobactericidal activity against the six indicated bacterial species of UV-A
377 irradiation (1.0 J cm^{-2}) in the absence (light-gray bars) and presence (unshaded bars) of 100 mg
378 l^{-1} FA. Black bars indicate 100 mg l^{-1} FA treatment in the absence of UV-A (for 1.95 min, the
379 incubation time used to provide a fluence of 1.0 J cm^{-2} UV-A). Data are presented as means \pm
380 SD ($n = 3$). Significant differences ($**P < 0.01$) were calculated based on comparison to
381 viability of the respective bacterium exposed to FA alone and to UV-A exposure without FA.
382 Samples for which survival was $<10 \text{ CFU ml}^{-1}$ (lower limit of detection) are represented as
383 6-log reductions.

384 **Figure 3.** Photoantimicrobial activity against four fungal species of UV-A irradiation (10 J
385 cm^{-2}) in the absence (light-gray bars) and presence (unshaded bars) of 1000 mg l^{-1} FA. Black
386 bars indicate 1000 mg l^{-1} FA treatment in the absence of UV-A (for 19.4 min, the incubation
387 time used to provide a fluence of 10 J cm^{-2} UV-A). Significant differences ($**P < 0.01$) were
388 calculated based on comparison to viability for the respective fungus exposed to FA addition
389 alone and UV-A without FA. Samples for which survival was $<10 \text{ CFU ml}^{-1}$ (lower limit of
390 detection) are noted in parentheses as 4-log unit reductions.

391

392

393

394 **Table 1** MICs of *trans*-FA and isomerized FA

Microorganism	MIC (mg l ⁻¹)*	
	<i>trans</i> -FA	Isomerized FA**
<i>Escherichia coli</i> NBRC12713	>1000	>1000
<i>Salmonella enterica</i> NBRC13245	1000	1000
<i>Staphylococcus aureus</i> NBRC12732	500	1000
<i>Bacillus cereus</i> NBRC15305	1000	1000
<i>Bacillus subtilis</i> ATCC6633	1000	670 ± 290
<i>Kocuria rhizophila</i> NBRC12708	500	500
<i>Alicyclobacillus acidoterrestris</i> NBRC108913	125	125
<i>Sporolactobacillus inulinus</i> NBRC13595	500	500
<i>Saccharomyces cerevisiae</i> NBRC1136	>1000	>1000
<i>Cladosporium cladosporioides</i> IFM63149	>1000	>1000
<i>Byssochlamys fulva</i> NBRC31767	>1000	>1000
<i>Eupenicillium lapidosum</i> NBRC6100	>1000	>1000

395

396 *Values are the mean ± SD obtained from three independent experiments. Values without SDs
 397 were identical in each of the three independent experiments.

398 **Composed of 93% *cis*-form and 7% *trans*-form




Fig 1 UV-A fluence-dependent changes in *E. coli* survival following irradiation in the absence (A) or presence (B) of FA. Filled (black) symbols, no UV-A exposure; unfilled (white) symbols, UV-A exposure. Cell suspensions were treated with FA at 50 mg l⁻¹ (diamond), 100 mg l⁻¹ (triangle), or 150 mg l⁻¹ (circle). Data are presented as means \pm SD (n = 3). Significant differences (**P < 0.01) were calculated based on comparison to samples irradiated with UV-A (3.0 J cm⁻²) in the absence of FA. Samples for which survival was <10 CFU ml⁻¹ (lower limit of detection) are noted in parentheses as 6-log unit reductions.




Fig 2 Photobactericidal activity against the six indicated bacterial species of UV-A irradiation (1.0 J cm^{-2}) in the absence (light-gray bars) and presence (unshaded bars) of 100 mg l^{-1} FA. Black bars indicate 100 mg l^{-1} FA treatment in the absence of UV-A (for 1.95 min, the incubation time used to provide a fluence of 1.0 J cm^{-2} UV-A). Data are presented as means \pm SD ($n = 3$). Significant differences (** $P < 0.01$) were calculated based on comparison to viability of the respective bacterium exposed to FA alone and to UV-A exposure without FA. Samples for which survival was $<10 \text{ CFU ml}^{-1}$ (lower limit of detection) are represented as 6-log reductions.




Fig 3 Photoantimicrobial activity against four fungal species of UV-A irradiation (10 J cm^{-2}) in the absence (light-gray bars) and presence (unshaded bars) of 1000 mg l^{-1} FA. Black bars indicate 1000 mg l^{-1} FA treatment in the absence of UV-A (for 19.4 min, the incubation time used to provide a fluence of 10 J cm^{-2} UV-A). Significant differences (** $P < 0.01$) were calculated based on comparison to viability for the respective fungus exposed to FA addition alone and UV-A without FA. Samples for which survival was $<10 \text{ CFU ml}^{-1}$ (lower limit of detection) are noted in parentheses as 4-log unit reductions.

1 Preparation, separation, and purity of *cis*-FA.

2 *cis*-FA was prepared by isomerization of commercially available FA by UV-A irradiation. FA
3 (250 mg) was dissolved in 5 ml methyl alcohol (HPLC grade) and the mixture was irradiated
4 using the UV-A LED, as used in the photoantimicrobial assay, at 6.18 mW cm^{-2} for 15 min at
5 room temperature (around 25°C). The LED was used to irradiate downward into a standard
6 glass Petri dish (internal diameter 27 mm) without a cover at a distance of 50 mm between the
7 LED and the bottom of the Petri dish.

8 Isolation of the *cis*-isomer from the *cis*- and *trans*-FA mixture was performed using a
9 Shimadzu HPLC system (pump, LC-20AT; auto-sampler, SIL-20AC; UV-detector, SPD-
10 M20A; column oven, CTO-20AC; Kyoto, Japan) equipped with a COSMOSIL column (HILIC,
11 $10 \times 250 \text{ mm}$; Nacalai Tesque Inc.) using acetonitrile/10 mM ammonium acetate (aq.) (9:1) as
12 the mobile phase (flow rate 8 ml min^{-1}). The detection wavelength was 318 nm, which was the
13 maximum absorbance wavelength of the mixture. The resulting HPLC spectrum before
14 separation is shown in Figure S2(A). Two major peaks were detected with retention times of
15 21–38 min and 39–45 min; these peaks were assigned as *trans*-FA and *cis*-FA, respectively, on
16 the basis of the proton nuclear magnetic resonance ($^1\text{H-NMR}$) spectra using a JEM-EX 400
17 spectrometer (JOEL, Tokyo, Japan).

18 After the *cis*-FA eluent was collected, the organic solvent was evaporated. The residual
19 solution was adjusted to pH 3–4 with 6 M HCl (aq.). An ethyl acetate layer extracted from the
20 solution was washed with brine (sat. NaCl sol.), dried over sodium sulfate, and concentrated
21 under reduced pressure. Purification by flash chromatography on silica gel (silica gel 120,

22 mesh 70–230 spherical; Nacalai Tesque Inc.) using elution with chloroform/methyl alcohol
23 (20:1) yielded *cis*-FA (yellowish, oily). Mass analysis (Acquity UPLC-LCT Premier liquid
24 chromatography-mass spectrometry system; Nihon Waters K.K., Tokyo, Japan) yielded a mass
25 for the isomerized FA consistent with the expected value for this compound [ESI-MS (*m/z*) for
26 $C_{10}H_{10}O_4$, calculated: 194.06 $[M]^+$, found: 195.066 $[M+H]^+$].

27 The proportions of the *cis*- and *trans*-forms in the isomerized FA after separation and
28 purification were determined using the Shimadzu HPLC equipped with a COSMOSIL column
29 (HILIC, 4.6×150 mm; Nacalai Tesque Inc.) at a detection wavelength of 318 nm. The elution
30 solvent was acetonitrile/10 mM ammonium acetate (aq.) (8:2) at a flow rate of 0.5 ml min^{-1}
31 [Fig. S2(B)]. To determine the difference in peak area corresponding to *trans*-FA,
32 commercially available *trans*-FA and the isomerized FA were (separately) dissolved at 0.35
33 mg ml^{-1} in 80% DMSO and assayed. The isomerized FA was shown to be composed of 93%
34 *cis*-form and 7% *trans*-form on the basis of the HPLC analysis. Additionally, the isomerized
35 FA was analyzed with $^1\text{H-NMR}$; for this assay, the isomerized FA was dissolved in $\text{DMSO-}d_6$
36 with tetramethylsilane as an internal standard. Integrated values for the α -H of the double bond
37 in the propenoic acid moiety corresponding to *cis*-FA and *trans*-FA were compared. $^1\text{H-NMR}$
38 analysis confirmed that the isomerized FA included both isomers, as demonstrated by detection
39 of the α -H of double bonds in each propenoic acid moiety corresponding to *cis*- and *trans*-FA;
40 the coupling constants were 13.0 Hz (δ , 5.73 ppm; doublet peak) and 15.9 Hz (δ , 6.36 ppm;
41 doublet peak), respectively. These NMR data were consistent with those in a previous report
42 (Guo *et al.* 2015). The proportion of the *cis*-form, calculated from the integrated values for the

43 α -H protons relevant to the *cis*- and *trans*-forms, respectively, was similar to that derived
44 based on HPLC analysis.

45 For determination of UV-visible absorbance spectra, commercially available *trans*-FA and
46 the isomerized FA were (separately) dissolved at 50 μ M in 0.4% (v/v) DMSO. Spectra were
47 measured using a 1-cm pathlength cuvette and a U-3300 spectrophotometer (Hitachi Ltd.,
48 Tokyo, Japan). The UV absorption spectrum of the *cis*-form was distinct from that of the
49 *trans*-form (Fig. S3); the difference was very similar to that observed between resveratrol
50 isomers (Trela and Waterhouse 1996).

51 When the isomerized FA was prepared with 80% DMSO or pure water, the composition
52 was stable at 30°C for at least 98 days, as judged by negligible time-dependent changes in the
53 area of the HPLC peak corresponding to *trans*-FA (data not shown). The isomerized FA,
54 which was highly enriched for the *cis*-form, was used for determination of MICs.

55 **References**

- 56 Guo, J., Zhang, J., Wang, W., Liu, T. and Xin, Z. (2015) Isolation and identification of bound
57 compounds from corn bran and their antioxidant and angiotensin I-converting enzyme
58 inhibitory activities. *Eur Food Res Technol* **241**, 37-47.
- 59 Trela, B.C. and Waterhouse, A.L. (1996) Resveratrol: isomeric molar absorptivities and
60 stability. *J Agric Food Chem* **44**, 1253-1257.

Table S1 Conditions for the preculture and viability assay of the tested **microorganisms**

Microorganism	Preculture condition	Viability assay
	Medium / Temp. / Time of growth	Medium / Temp. / Time of growth
<i>E. coli</i> NBRC12713	LB [*] / 37°C / 17 h with shaking	SCDLP agar ^{††} / 37°C / 48 h
<i>Salm. enterica</i> NBRC13245	LB / 37°C / 17 h with shaking	SCDLP agar / 37°C / 48 h
<i>Staph. aureus</i> NBRC12732	LB / 37°C / 17 h with shaking	SCDLP agar / 37°C / 48 h
<i>B. cereus</i> NBRC15305	LB / 37°C / 17 h with shaking	SCDLP agar / 37°C / 48 h
<i>B. subtilis</i> ATCC6633	LB / 37°C / 17 h with shaking	SCDLP agar / 37°C / 48 h
<i>K. rhizophila</i> NBRC12708	LB / 37°C / 17 h with shaking	SCDLP agar / 37°C / 48 h
<i>A. acidoterrestris</i> NBRC108913	Specific broth (No. 323) ^{**} / 45°C / 17 h with shaking	No. 323 agar / 45°C / 48 h
<i>Sporolact. inulinus</i> NBRC13595	MRS broth ^{***} / 37°C / 24 h in an anaerobic chamber (no shaking)	MRS broth / 37°C / 72 h in an anaerobic chamber
<i>S. cerevisiae</i> NBRC1136	Sabouraud broth ^{**} / 28°C / 24 h with shaking	Sabouraud agar ^{†††} / 28°C / 72 h
<i>C. cladosporioides</i> IFM63149	PDA [†] / 25°C / 12 days	PDA / 25°C / 72 h
<i>B. fulva</i> NBRC31767	PDA / 25°C / 12 days	PDA / 25°C / 72 h
<i>E. lapidosum</i> NBRC6100	PDA / 25°C / 12 days	PDA / 25°C / 72 h

^{*}Luria-Bertani medium (Lennox; Nacalai Tesque Inc., Kyoto, Japan); ^{**}See Materials and Methods; ^{***}MRS broth for microbiology (Sigma-Aldrich Co., LLC, St. Louis, MO, USA); [†]Potato dextrose agar (Nissui Pharmaceutical Co. Ltd., Tokyo, Japan); ^{††}Nihon Pharmaceutical Co., Ltd. Tokyo, Japan; ^{†††}Nissui Pharmaceutical Co. Ltd.




Fig S1 Emission spectrum of the UV-A LED as used in this study; the spectrum exhibited a maximum at 365 nm, which was measured at a distance of 30 mm between the illumination source and the UV meter.




Fig S2 HPLC chromatogram of FA prepared by UV-A exposure: before separation (A), and after separation and purification of the *cis*-form (B).




Fig S3 UV-visible absorption spectra of FA (*trans*-form, solid line) and isomerized FA (mixture of 93% *cis*-form and 7% *trans*-form, dashed line). Samples were dissolved at 50 μ M in 0.4% DMSO.

Authors' answers against Reviewers' comments

I indicated all correction points (changes and edits) by using the highlighter tool to highlight the changes in yellow in our manuscript.

REFeree COMMENTS TO AUTHORS (Reviewer 1)

Without underestimating the author's effort, but taking their experience from previous cited scientific reports, this study would have been more completed if trials of food liquids would had been performed. In that way, it will give us a clue on the effect of the organic content of these foods items (juices) or if ferulic acid would have affected the quality attributes of the juices.

Authors' answer

Thank you for giving me some advices related to next challenge. In this report, the inactivation experiments were carried out in sterile water of high UV transmittance. In future work, we will show the potential use of the combination of UV-A light and FA in complex food surfaces.

Comment of reviewer 1 (in general)

Page 2, line 21. Use the word "microorganisms" instead of "microbes".

Ans. I corrected the term of "microbes" to "microorganisms" in our manuscript, report title, and Tables 1 and S1 (Lines 1, 63, 96, 98, 208 and 359).

Comment of reviewer 1 (in general)

Page 2, line 20. I prefer to read the impact of food/juice industry when using this proposed method of disinfection.

Ans. I rewrote "SIGNIFICANCE AND IMPACT OF THE STUDY" (Line 21-28).

Comment of reviewer 1 (in general)

Page 2, line 29. The term "sterilization" is not the right one. I suggest "decontamination" and/or "disinfection".

Ans. I corrected the term of “sterilization” to “disinfection” in our manuscript (Lines 30, 47).

Comment of reviewer 1 (in general)

Page 2, line 31. The word "killing" is a general term, I would prefer a more "biological" term: "reducing" and/or "eliminating".

Ans. I corrected the term of “killing” (Lines 32, 37, 38, 145, 181, 186, 187 and 200).

Comment of reviewer 1 (in general)

Page 2, line 40. Delete the word "minimum".

Ans. I deleted “minimum” (Line 41).

Comment of reviewer 1 (in general)

Page 19. Table 1. Do not abbreviate scientific names.

Ans. I indicated scientific names for microorganisms in Table 1 (Page 19).

REFeree COMMENTS TO AUTHORS (Reviewer 2)

Major concerns.

In my opinion, the section corresponding to significance and impact of the study is not written according to the instructions given in the author guidelines of the journal; it has been written more as a kind of abstract. Please, check the instructions and modify.

Authors' answer

I rewrote “SIGNIFICANCE AND IMPACT OF THE STUDY” (Line 21-28).

Comment of reviewer 2

Lines 117-118. According to your citation of Graf (1992), is not possible that trans ferulic acid had been isomerized to cis-form by your UV treatment, therefore, you had actually had cis-form in both cases?

Authors' answer

Please see supporting information (Text S1, Page 23-25).

cis-FA was generated by isomerization using UV-A exposure of *trans*-FA, and then the *cis*-form was separated from the parent isomer using HPLC. We confirmed that the separated FA was composed of 93% *cis*-form and 7% *trans*-form on the basis of the HPLC analysis, as demonstrated by detection of the α -H of double bonds in each propenoic acid moiety corresponding to *cis*- and *trans*-FA; the coupling constants were 13.0 Hz (δ , 5.73 ppm; doublet peak) and 15.9 Hz (δ , 6.36 ppm; doublet peak), respectively.

Comment of reviewer 2

Line 155. On complete inactivation. How are authors sure that complete inactivation was achieved without performing enrichment tests that can show that no survivors existed?

Ans. I thought that the word of “complete” was not appropriate, because the inactivation means that viable cell count was below 10 CFU ml⁻¹ (the lower limit of detection). In line 153, I corrected the term of “Complete inactivation” to “Potent inactivation”.

Comment of reviewer 2 (Minor concerns)

Line 40. Delete “Minimum”.

Ans. I deleted “minimum” (line 41).

Line 48. More versatile than what?

Ans. I corrected “a more versatile UV source” to “a versatile UV source” (Line 48).

Line 58. Delete “D.”

Ans. I deleted “D” (Line 58)..

Lines 59-61. Which is the advantage of UV-A over UV-C in this regard?

Ans. UV-A is superior to UV-C in this regard. A previous report has described no differences of Vitamin C content in cabbage tissue before and after UV-A irradiation (Aihara M., et al., J Med Invest (2014) 61, 285-290).

Lines 67, 122. Avoid using very personal terms, write in a more impersonal fashion.

Ans. I wrote in sentences as follows (Lines 67-70):

To increase the inactivation efficiency of UV-A, the synergistic bactericidal activity of the

combination of UV-A light and organic substrates, a quaternary ammonium salt (Shirai et al. 2014), and natural compounds [coumaric acid and ferulic acid (FA)] and their derivatives, has been investigated in our laboratory (Shirai et al. 2015a, 2015b).

Ans. I wrote in sentences as follows (Line 121):

Photoantimicrobial assays were performed at FA concentrations below its MIC.

Line 78. Delete “D” before “Paiva”.

Ans. I deleted “D” (Line 76).

Lines 80-93. Write in few lines the goal of this research at the end of the introduction in order for the reader to catch it quickly and clearly, avoiding mixing them with the justification of testing the microorganisms used in the research, which can be placed before the goals.

Ans. Please see lines 80-92 in manuscript.

Lines 132 and elsewhere. It is not necessary to repeat each time data on concentration, fluences, etc. It is enough to do it at first mentioning.

Ans. I rewrote sentences as follows (Lines 130-131):

the combination of FA with UV-A yielded significant ($P < 0.01$) decreases in the viable cell count.

Line 139 and elsewhere. Use a single letter for generic names.

Ans. The abbreviation for those generic names, *Salm.*, *Staph.*, and *Sporolact.*, complies with “common generic names” indicated in Author Guidelines for this journal (Lines 137 and elsewhere).

Line 183. Use only two decimals.

Ans. I corrected 0.007 to 0.01 (Line 181).

Line 198. Separate conclusions in a single paragraph for better exposure to the reader.

Ans. I separated the conclusions in a single paragraph (Line 197).

Line 212. Microbial strains must be included in this section. You could cite table 1 to this end.

Ans. I wrote in sentence as follows:

Those strains are listed in Table 1 (Line 210).

REFEREE COMMENTS TO AUTHORS (Reviewer 3)

This manuscript is useful and helpful to develop new types of antimicrobial reagents, especially combination with UV-A without hazardous UV-C is interesting. On the other hand, why does UV-A enhance the antimicrobial activity of FA? This question is a major issue to the next research. I hope you experimentally solve this question soon.

Ans. I am going to solve the synergistic bactericidal mechanism in a next report.