

EXPANDED ABSTRACT

AVP-stimulated nucleotide secretion in perfused mouse medullary thick ascending limb and cortical collecting duct

Elvin Odgaard, Helle A Praetorius, and Jens Leipziger

Department of Physiology and Biophysics, The Water and Salt Research Center, Aarhus University, Aarhus, Denmark

Summary : Extracellular nucleotides are local, short-lived signaling molecules that inhibit renal tubular transport via both luminal and basolateral P2 receptors (1, 2). Apparently, the renal epithelium itself is able to release nucleotides (3, 4). The mechanism and circumstances under which epithelia nucleotide release is stimulated remains elusive (5, 6). Here, we investigate the phenomenon of nucleotide secretion in intact perfused mouse medullary thick ascending limb (mTAL) and cortical collecting duct (CCD). The nucleotide secretion was monitored by a biosensor cell placed to register nucleotides in the tubular out-flow. $[Ca^{2+}]_i$ was measured simultaneously in the biosensor cells and the renal tubule with fluo-4. We were able to identify spontaneous tubular nucleotide secretion in resting perfused mTAL. This was seen as lively $[Ca^{2+}]_i$ oscillations in the nucleotide biosensor cells when the tubular outflow fluid engulfed the sensing cells. In mouse mTAL 10 nM AVP and dDAVP induced robust $[Ca^{2+}]_i$ oscillations, whereas AVP in the CCD induced large, slow and transient $[Ca^{2+}]_i$ elevations. Importantly, we identify that AVP/dDAVP triggers tubular secretion of nucleotides in mTAL. After addition of AVP/dDAVP the biosensor cells registered bursts of nucleotides originating from the tubular perfusate. The approximated tubular nucleotide concentration reached peak values of $\sim 0.2\text{-}0.3 \mu\text{M}$. A very similar response was observed after AVP stimulation of CCDs. Thus, AVP stimulated tubular secretion of nucleotides in a burst like pattern with peak tubular nucleotide concentrations in the low micromolar range. Luminal nucleotides are prone to activate luminal P2 receptors (1) which in turn are well described to inhibit AVP-augmented aquaporin-2-dependent water absorption (7) or ENaC-mediated Na^+ transport (8). Therefore, we speculate that local nucleotide signaling is an intrinsic feed-back element of hormonal control of renal tubular transport (9). *J. Med. Invest.* 56 Suppl. : 262-263, December, 2009

Keywords : purinergic signaling, ATP release, ATP biosensor, renal epithelium

Received for publication November 2, 2009 ; accepted November 9, 2009.

Address correspondence and reprint requests to Jens Leipziger, Department of Physiology and Biophysics, The Water and Salt Research Center, Aarhus University, Ole Worms Allé 1160, 8000 Aarhus C, Denmark and Fax : +45-86-129065.

REFERENCES

1. Leipziger J : Control of epithelial transport via luminal P2 receptors. *Am J Physiol Renal Physiol* 284 : F419-F432, 2003
2. Vallon V : P2 receptors in the regulation of renal transport mechanisms. *Am J Physiol Renal*

- Physiol 294 : F10-F27, 2008
3. Geyti CS, Odgaard E, Overgaard MT, Jensen ME, Leipziger J, Praetorius HA : Slow spontaneous $[Ca^{2+}]_i$ oscillations reflect nucleotide release from renal epithelia. *Pflügers Arch* 455 : 1105-1117, 2008
 4. Praetorius HA, Frokiaer J, Leipziger J : Trans-epithelial pressure pulses induce nucleotide release in polarized MDCK cells. *Am J Physiol Renal Physiol* 288 : F133-F141, 2005
 5. Praetorius HA, Leipziger J : Fluid flow sensing and triggered nucleotide release in epithelia. *J Physiol* 586 : 2669, 2008.
 6. Lazarowski ER, Boucher RC, Harden TK : Mechanisms of release of nucleotides and integration of their action as P2X- and P2Y-receptor activating molecules. *Mol Pharmacol* 64 : 785-795, 2003
 7. Kishore BK, Chou C-L, Knepper MA : Extracellular nucleotide receptor inhibits AVP-stimulated water permeability in inner medullary collecting duct. *Am J Physiol* 269 : F863-F869, 1995
 8. Lehrmann H, Thomas J, Kim SJ, Jacobi C, Leipziger J : Luminal P2Y₂ receptor-mediated inhibition of Na⁺ absorption in isolated perfused mouse CCD. *J Am Soc Nephrol* 13 : 10-18, 2002
 9. Odgaard E, Praetorius HA, Leipziger J : AVP-stimulated nucleotide secretion in perfused mouse medullary thick ascending limb and cortical collecting duct. *Am J Physiol Renal Physiol* 297 : F341-F349, 2009