

PROCEEDING

Role of protein kinase C- δ in isoproterenol-induced amylase release in rat parotid acinar cells

Hiroshi Sugiya^{1,2}, Keitaro Satoh¹, Miwako Matsuki-Fukushima¹, Bing Qi¹,
Ming-Yu Guo¹, and Junko Fujita-Yoshigaki^{1,2}

¹Department of Physiology and ²Research Institute of Oral Science, Nihon University School of Dentistry at Matsudo, Matsudo, Japan

Abstract : In parotid acinar cells, β -adrenergic receptor activation results in accumulation of intracellular cAMP. Subsequently, cAMP-dependent protein kinase (PKA) is activated and consequently amylase release is provoked. In this paper, we investigated involvement of protein kinase C- δ (PKC δ), a novel isoform of PKC, in amylase release induced by β -adrenergic receptor stimulation. Amylase release stimulated with the β -agonist isoproterenol (IPR) was inhibited by rottlerin, an inhibitor of PKC δ . IPR activated PKC δ and the effect of IPR were inhibited by a PKA inhibitor, H89. Myristoylated alanine-rich C kinase substrate (MARCKS), a major cellular substrate for PKC, was detected in rat parotid acinar cells, and a MARCKS inhibitor, MARCKS-related peptide, inhibited the IPR-induced amylase release. IPR stimulated MARCKS phosphorylation, which was found to be inhibited by H89 and rottlerin. These observations suggest that PKC δ activation is a downstream pathway of PKA activation and is involved in amylase release *via* MARCKS phosphorylation in rat parotid acinar cells stimulated with β -adrenergic agonist. *J. Med. Invest.* 56 Suppl. : 368-370, December, 2009

Keywords : PKC δ , amylase release, MARCKS, parotid acinar cells

INTRODUCTION

In parotid acinar cells, stimulation of β -adrenergic receptors provokes amylase release (1). The amylase release is depended on the increase in intracellular cAMP levels stimulated with β -adrenergic receptors but occurs independent of elevation of intracellular calcium ion (Ca²⁺) levels. Activation of cAMP-dependent protein kinase (PKA) followed by the increase in intracellular cAMP levels is thought to be an essential step in this process (1). On the other hand, it has also been thought that protein

kinase C (PKC) activation is involved in amylase release in parotid acinar cells (2-4). We here investigated the role of PKC on amylase release induced by the β -adrenergic agonist isoproterenol (IPR) in rat parotid acinar cells.

INVOLVEMENT OF PKC δ IN IPR-INDUCED AMYLASE RELEASE

The PKC family consists of at least 10 members, which are classified into three subgroups based on their structural characteristics and cofactor requirement. They include the conventional cPKC (α , β I, β II and γ), stimulated by Ca²⁺ and diacylglycerol (DAG) or phorbol esters; the novel nPKC (δ , ϵ , η and θ), activated by DAG and phorbol esters but Ca²⁺ independent; and the atypical aPKC (ζ and ι/λ),

Received for publication October 13, 2009; accepted October 20, 2009.

Address correspondence and reprint requests to Hiroshi Sugiya, Department of Physiology, Nihon University School of Dentistry at Matsudo, 2-870-1 Sakaecho-nishi, Matsudo, Chiba 271-8587, Japan and Fax : +81-47-360-9325.

unresponsive to Ca^{2+} , DAG and phorbol esters (5, 6). Of those, the expression of PKC α , δ , ϵ and ζ has been reported in rat parotid acinar cells (7, 8). In the presence of calphostin C, an inhibitor of DAG-sensitive PKC, IPR-induced amylase release was reduced. Since β -adrenergic receptor activation does not elicit Ca^{2+} mobilization, this observation suggests that a Ca^{2+} -independent and DAG-sensitive PKC, i.e. a novel PKC such as PKC δ , is involved in IPR-induced amylase release. Then, the effect of the PKC δ inhibitor rottlerin was examined. The inhibitor mimicked the effect of calphostin C. Although rottlerin has been used widely as a PKC- δ inhibitor, this polyphenolic compound has also been reported to cause a general inhibition of the mitochondrial metabolism which correlated with a depletion of the cellular ATP concentration (9, 10). Then we examined the effect of rottlerin on cellular ATP level in rat parotid acinar cells. However, the cellular ATP level was scarcely reduced by rottlerin. Therefore, it is unlikely that the inhibition of IPR-induced amylase release by rottlerin is caused by a toxic effect.

ACTIVATION OF PKC δ BY IPR

It is known that activated isoforms of PKC, including PKC δ , are phosphorylated and translocated from the cytosol to the cell membrane (11). When parotid acinar cells were stimulated with IPR, the increase in the amount of phosphorylated-PKC δ in the membrane fraction was observed by Western blotting using anti-phosphorylated-PKC δ antibody. This observation suggests that stimulation of β -adrenergic receptor induces PKC δ activation in parotid acinar cells. Further, we directly examined kinase activity of PKC δ using PKC δ -specific immunoprecipitates. When the kinase activity was measured with PKC δ purified from cell lysate by immunoprecipitation, the activity was shown to be increased in the acinar cells by treatment with IPR. Next we examined the effect of the PKA inhibitor H89 on IPR-induced PKC δ activation by detection of phosphorylated-PKC δ in the membrane fraction. In the presence of H89, IPR failed to increase the amount of phosphorylated-PKC δ in the membrane fraction, indicating that this PKA inhibitor inhibits IPR-induced PKC δ activation. These results suggest that PKC δ is activated via PKA activation in the IPR-stimulated rat parotid acinar cells.

MARCKS PHOSPHORYLATION INDUCED BY IPR

Myristoylated alanine-rich C kinase substrate (MARCKS) is a major cellular substrate for PKC (12). The involvement of MARCKS in the IPR-induced amylase release was examined in parotid acinar cells. The N-terminal sequence peptide of MARCKS (MANS) has been shown to be a useful inhibitor of MARCKS function because this peptide suppresses the secretory function *in vivo* (13, 14) and in cultured cells (15). In parotid acinar cells pretreated with the MANS peptide, IPR-induced amylase release was reduced, suggesting the involvement of MARCKS in IPR-induced amylase release.

Involvement of phosphorylation of MARCKS by PKC in secretory function has been demonstrated in chromaffin cells (16, 17), insulin-producing INS-1 cells (18), bovine luteal cells (19), and SH-SY5Y human neuroblastoma cells (20). When we examined the effect of IPR on MARCKS phosphorylation by Western blotting using anti-phosphorylated MARCKS antibody, this β -agonist induced MARCKS phosphorylation in parotid acinar cells. The IPR-induced MARCKS phosphorylation was inhibited in the presence of a PKA inhibitor, H89, implying that MARCKS phosphorylation stimulated by IPR is evoked via PKA activation in parotid acinar cells. In the presence of the PKC δ -specific inhibitor, rottlerin, the MARCKS phosphorylation was also inhibited. These results suggest that PKC δ activated by IPR stimulation via PKA activation provokes MARCKS phosphorylation.

Taken together, it is most likely that PKA-PKC δ -MARCKS pathway is involved in IPR-induced amylase release in rat parotid acinar cells.

ACKNOWLEDGEMENTS

This study was supported by a Nihon University Multidisciplinary Research Grant for 2008-2009 and a Grant for Supporting Project for Strategic Research by MEXT, 2008-2012.

REFERENCES

1. Quissell DO, Watson E, Dowd FJ : Signal transduction mechanisms involved in salivary gland regulated exocytosis. *Crit Rev Oral Biol Med* 3 : 83-107, 1992

2. Putney JW Jr, McKinney JS, Aub DL, Leslie BA : Phorbol ester-induced protein secretion in rat parotid gland. *Mol Pharmacol* 26 : 261-266, 1984
3. Takuma T, Ichida T : Phorbol ester stimulates amylase secretion from rat parotid cells. *FEBS Lett* 199 : 53-56, 1986
4. McKinney JS, Rubin DP : Enhancement of cyclic AMP modulated salivary amylase secretion by protein kinase C activators. *Biochem Pharmacol* 37 : 4433-4438, 1988
5. Breitzkreutz D, Braiman-Wiksman L, Daum N, Denning MF, Tennenbaum T : Protein kinase C family : on the crossroads of cell signaling in skin and tumor epithelium. *J Cancer Res Clin Oncol* 133 : 793-808, 2007
6. Steinberg SF : Structural basis of protein kinase C isoform function. *Physiol Rev* 88 : 1341-1378, 2008
7. Terzian AR, Rubin RP : Translocation of the α -isozyme of protein kinase C during stimulation of rat parotid acinar cells by phorbol ester and carbachol. *Arch Oral Biol* 38 : 1051-1056, 1993
8. Terzian AR, Zhang X, Rubin RP : Differential modulation of protein kinase C isozymes in rat parotid acinar cells. *Biochem Pharmacol* 52 : 569-577, 1996
9. Soltoff SP : Rottlerin is a mitochondrial uncoupler that decreases cellular ATP levels and indirectly blocks protein kinase C δ tyrosine phosphorylation. *J Biol Chem* 276 : 37986-37992, 2001
10. Tapia JA, Jensen RT, García-Marín LJ : Rottlerin inhibits stimulated enzymatic secretion and several intracellular signaling transduction pathways in pancreatic acinar cells by a non-PKC- δ -dependent mechanism. *Biochim Biophys Acta* 1763 : 25-38, 2006
11. Shirai Y, Saito N : Activation mechanisms of protein kinase C : maturation, catalytic activation, and targeting. *J Biochem* 132 : 663-668, 2002
12. Stumpo DJ, Graff JM, Albert KA, Greengard P, Blackshear PJ : Molecular cloning, characterization, and expression of a cDNA encoding the "80- to 87-kDa" myristoylated alanine-rich C kinase substrate : a major cellular substrate for protein kinase C. *Proc Natl Acad Sci USA* 86 : 4012-4016, 1989
13. Agrawal A, Rengarajan S, Adler KB, Ram A, Ghosh B, Fahim M, Dickey BF : Inhibition of mucin secretion with MARCKS-related peptide improves airway obstruction in a mouse model of asthma. *J Appl Physiol* 102 : 399-405, 2007
14. Singer M, Martin LD, Vargaftig BB, Park J, Gruber AD, Li Y, Adler KB : A MARCKS-related peptide blocks mucus hypersecretion in a mouse model of asthma. *Nat Med* 10 : 193-196, 2004
15. Li Y, Martin LD, Spizz G, Adler KB : MARCKS protein is a key molecule regulating mucin secretion by human airway epithelial cells *in vitro*. *J Biol Chem* 276 : 40982-40990, 2001
16. Rosé SD, Lejen T, Zhang L, Trifaró JM : Chromaffin cell F-actin disassembly and potentiation of catecholamine release in response to protein kinase C activation by phorbol esters is mediated through myristoylated alanine-rich C kinase substrate. *J Biol Chem* 276 : 36757-36763, 2001
17. Trifaró JM, Rosé SD, Lejen T, Elzagallaai A : Two pathway control chromaffin cell cortical F-actin dynamics during exocytosis. *Biochimie* 82 : 339-352, 2000
18. Suzuki Y, Zhang H, Saito N, Kojima I, Urano T, Mogami H : Glucagon-like peptide 1 activates protein kinase C through Ca²⁺-dependent activation of phospholipase C in insulin-secreting cells. *J Biol Chem* 281 : 28499-28507, 2006
19. Salli U, Saito N, Stormshak F : Spatiotemporal interactions of myristoylated alanine-rich C kinase substrate (MARCKS) protein with the actin cytoskeleton and exocytosis of oxytocin upon prostaglandin F₂ α stimulation of bovine luteal cells. *Biol Reprod* 69 : 2053-2058, 2003
20. Goodall AR, Turner NA, Walker JH, Ball SG, Vaughan PF : Activation of protein kinase C- α and translocation of the myristoylated alanine-rich C kinase substrate correlate with phorbol ester-enhanced noradrenaline release from SH-SY5Y human neuroblastoma cells. *J Neurochem* 68 : 392-401, 1997