

This is a post-peer-review, pre-copyedit version of an article published in *Inflammation*.  
The final authenticated version is available online at: <https://doi.org/10.1007/s10753-018-0764-0>

*Research Article*

**Distinct regulation of CXCL10 production by cytokines in human salivary gland ductal and acinar cells**

Keiko Aota<sup>1,\*</sup>, Koichi Kani<sup>1</sup>, Tomoko Yamanoi<sup>1</sup>, Koh-ichi Nakashiro<sup>2</sup>,  
Naozumi Ishimaru<sup>3</sup>, and Masayuki Azuma<sup>1</sup>

<sup>1</sup>Department of Oral Medicine, Tokushima University Graduate School of Biomedical Sciences, Tokushima, Japan

<sup>2</sup>Department of Oral and Maxillofacial Surgery, Ehime University Graduate School of Medicine, Ehime, Japan

<sup>3</sup>Department of Oral Molecular Pathology, Tokushima University Graduate School of Biomedical Sciences, Tokushima, Japan

**\*Corresponding author:** Dr. Keiko Aota, Department of Oral Medicine, Tokushima University Graduate School of Biomedical Sciences, 3-18-15 Kuramoto-cho, Tokushima 770-8504, Japan. Tel: +81-88-633-7352; Fax: +81-88-633-7388.  
Email: [aota.keiko@tokushima-u.ac.jp](mailto:aota.keiko@tokushima-u.ac.jp)

**ORCID iD:** 0000-0001-5413-5750

**Financial support:** This work was supported by a Grant-in-Aid for Scientific Research from the Japan Ministry of Education, Culture, Sports, Science and Technology (No. 17K11842, No. 17K17258).

**ABSTRACT (224 words)**

1  
2  
3 CXCL10, a CXC chemokine induced by interferon-gamma [IFN- $\gamma$ ], has been observed in a  
4  
5  
6 wide variety of chronic inflammatory disorders and autoimmune conditions. Although  
7  
8  
9 CXCL10 is known to be overexpressed in the salivary glands of individuals with primary  
10  
11  
12 Sjögren's syndrome (pSS), it is unclear which cells produce CXCL10 under what types of  
13  
14  
15 stimulations. Here we investigated the precise molecular mechanisms by which CXCL10 was  
16  
17  
18 produced in human salivary gland ductal (NS-SV-DC) and acinar (NS-SV-AC) cell lines. Our  
19  
20  
21 results demonstrated that NS-SV-DC cells produced higher levels of CXCL10 compared to  
22  
23  
24 NS-SV-AC cells. In addition, our findings demonstrated that the regulator of the enhancement  
25  
26  
27 of CXCL10 was different between NS-SV-DC and NS-SV-AC cells; i.e., interferon-gamma  
28  
29  
30 (IFN- $\gamma$ ) had more potential than interferon-alpha (IFN- $\alpha$ ), tumor necrosis factor (TNF)- $\alpha$ , and  
31  
32  
33 interleukin (IL)1- $\beta$  in the induction of CXCL10 production in NS-SV-DC cells, whereas  
34  
35  
36 TNF- $\alpha$  had potential to induce CXCL10 production in NS-SV-AC cells. A Western blot  
37  
38  
39 analysis demonstrated that IFN- $\gamma$  enhanced the production of CXCL10 via both the  
40  
41  
42 JAK/STAT1 pathway and the NF- $\kappa$ B pathway in NS-SV-DC cells, whereas TNF- $\alpha$  enhanced  
43  
44  
45 the production of CXCL10 via the NF- $\kappa$ B pathway in NS-SV-AC cells. The results of study  
46  
47  
48 suggest that the CXCL10 overexpression in the salivary glands is caused mainly by  
49  
50  
51 IFN- $\gamma$ -stimulated salivary gland ductal cells. The enhanced production of CXCL10 by IFN- $\gamma$   
52  
53  
54 from ductal cells may result in the inflammation of pSS lesions.  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**KEY WORDS:**

CXCL10, IFN- $\gamma$ , TNF- $\alpha$ , primary Sjögren's syndrome, salivary gland ductal cells, salivary gland acinar cells

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## INTRODUCTION

1  
2  
3 Chemokines are a specialized family of cytokines that function as potent mediators of  
4  
5  
6 inflammation based on their ability to recruit and activate specific leucocyte subpopulations  
7  
8  
9 by modifying the expression and affinity status of adhesion molecules on the leucocyte  
10  
11  
12 surface [1]. CXCL10 is a non-ELR (lacking the Glu-Leu-Arg tripeptide motif) CXC  
13  
14  
15 chemokine [2]. CXCL9, CXCL10, and CXCL11 utilize the same chemokine receptor,  
16  
17  
18 CXCR3 [2], a G protein-coupled receptor. CXCR3 is induced on naïve T cells following  
19  
20  
21 activation, and preferentially remains highly expressed on type-1 helper (Th1)-type CD4<sup>+</sup> T  
22  
23  
24 cells, effector CD8<sup>+</sup> T cells, and innate-type lymphocytes, such as natural killer (NK) and  
25  
26  
27 NKT cells [2].

28  
29  
30 Sjögren's syndrome (SS), one of the most common autoimmune diseases [3], is  
31  
32  
33 characterized by the eventual total replacement of the acinar structure by marked lymphocytic  
34  
35  
36 infiltrates in the salivary and lacrimal glands [4]. Histological studies show that CD4<sup>+</sup> T cells  
37  
38  
39 predominantly infiltrate the salivary glands at an early stage of primary SS (pSS) [5]. It has  
40  
41  
42 been reported that CXCL9, CXCL10, and CXCL11 were involved in the accumulation of  
43  
44  
45 T-cell infiltrates in the salivary glands of pSS patients [6, 7]. In addition, the inoculation of a  
46  
47  
48 CXCL10 antagonist into MRL/*lpr* mice during the early stage of sialadenitis significantly  
49  
50  
51 reduced the mononuclear cell infiltration and parenchymal destruction [8].  
52

53  
54  
55 Interferons (IFNs) play significant roles in the pathogenesis of autoimmune diseases,  
56  
57  
58 including SS. IFNs are subdivided into three distinct types: type I IFN (IFN-I; mainly  
59  
60

1 IFN- $\alpha/\beta$ ), type II IFN (IFN-II; mainly IFN- $\gamma$ ), and type III IFN (IFN-III; mainly  
2  
3 IFN- $\lambda$ /interleukin [IL]-28/IL-29) [9]. Although the genes involved in SS were originally  
4  
5 considered to be mediated via only type I interferons, recent findings demonstrated that genes  
6  
7 upregulated by both type I and type II interferons are involved in pSS [10]. In addition,  
8  
9 microarray studies in the labial salivary glands (LSGs) of patients with pSS revealed that  
10  
11 several interferon-regulated genes were up-regulated [11,12]. Several other cytokines,  
12  
13 including IL-6, IL-7, IL-10, IL-12, IL-17, IL-21, tumor necrosis factor-alpha (TNF- $\alpha$ ) [13],  
14  
15 and IL-1 $\beta$  [14, 15], have been suggested to play roles in the pathogenesis of pSS.  
16  
17  
18  
19  
20  
21  
22  
23

24 It has thus been hypothesized that several cytokines cause autoimmune salivary  
25  
26 diseases, and that CXCL10 plays an important role in the pathogenesis of SS. However, it is  
27  
28 not yet known which cells produce CXCL10 by what types of cytokines. To the best of our  
29  
30 knowledge, no study has evaluated the effects of IFN- $\alpha$ , IFN- $\gamma$ , or several cytokines on the  
31  
32 secretion of CXCL9, CXCL10, and CXCL11 by human salivary ductal cells or acinar cells.  
33  
34 We therefore examined the effects of IFN- $\alpha$ , IFN- $\gamma$ , TNF- $\alpha$ , and IL-1 $\beta$  stimulation on the  
35  
36 expressions of CXCL9, CXCL10, and CXCL11 by using immortalized human salivary gland  
37  
38 cell clones in an *in vitro* experiment. We also investigated the signal transduction pathways  
39  
40 that induced the production of CXCL10.  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53

## 54 **MATERIALS AND METHODS**

### 57 **Cell culture**

1 The characteristics of the cell lines NS-SV-DC (immortalized human salivary gland ductal  
2 cells) and NS-SV-AC (immortalized human salivary gland acinar cells) are described in detail  
3 elsewhere [16]. These cell clones were cultured at 37°C in serum-free keratinocyte medium  
4  
5 (Gibco Laboratories, Gaithersburg, MD) in an incubator with an atmosphere containing 5%  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 18 **Reagents**

21 Recombinant human IFN- $\alpha$ , IFN- $\gamma$ , TNF- $\alpha$  and IL-1 $\beta$  were purchased from R&D Systems  
22  
23 (Minneapolis, MN). Bay11-7082 was purchased from Wako (Osaka, Japan).  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 42 **Quantitative real-time reverse transcription-polymerase chain reaction (qRT-PCR)**

45 We treated NS-SV-DC and NS-SC-AC cells with 1000 IU/mL IFN- $\alpha$ , 10 ng/mL IFN- $\gamma$ , 10  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 an ABI Prism 7000 Sequence Detection System (Applied Biosystems Japan, Tokyo) and  
2  
3 TaqMan® Universal PCR Master Mix (Applied Biosystems) and Assays-on-Demand™ Gene  
4  
5 Expression Products (Applied Biosystems) according to the manufacturer's recommendations.  
6  
7  
8  
9 The thermal cycler protocol was 95°C for 10 min, followed by 40 cycles of 95°C for 5 s and  
10  
11 60°C for 30 s.  
12

13  
14 We performed an analysis of the relative gene expression data using the  $2^{-\Delta\Delta CT}$  method  
15  
16 on Sequence Detection System Software (Applied Biosystems). We calculated the fold  
17  
18 change in the studied gene expression, normalized to an endogenous control, using the  
19  
20 formula  $RQ = 2^{-\Delta\Delta CT}$ . The relative expression levels of CXCR3 ligand mRNAs are expressed  
21  
22 as the fold increase in the GAPDH mRNA expression.  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

### 33 **Reverse transcription-polymerase chain reaction (RT-PCR)**

34  
35  
36 The total cellular RNA was isolated from unstimulated NS-SV-DC and NS-SC-DC cells  
37  
38 using TRIzol reagent. The cDNA was synthesized according to the above method. Specific  
39  
40 primers were custom synthesized (Sigma, Deisenhofen, Germany). The sense and anti-sense  
41  
42 primers for IFNGR1, IFNGR2, TNFR1, TNFR2 and GAPDH, respectively, were as follows:  
43  
44  
45 primers for IFNGR1, IFNGR2, TNFR1, TNFR2 and GAPDH, respectively, were as follows:  
46  
47  
48 5'-GCTGTATGCCGAGATGGAAAA-3' and 5'-AGGAAAATGGCTGGTATGACG-3',  
49  
50  
51 5'-CGACAGTAAATGGTTCACGGC-3' and  
52  
53  
54 5'-TGGACATAATAACAAAAAAGGC-3', 5'-GCCTCTGCCTCAATGGGACCG-3' and  
55  
56  
57 5'-GGGGTGAAGCCTGGAGTGGGA-3', 5'-GCCCACTCGGAACCAGCCACA-3' and  
58  
59  
60  
61  
62  
63  
64  
65

1 5'-TCAGGCACTCCAAGGGGCAGG-3', 5'-ACGCATTTGGCTGTATTGGG-3' and  
2  
3 5'-TGATTTTGGAGGGATCTCGC-3'. The PCR reactions were conducted in a DNA  
4  
5 Thermal Cycler (model LifeECO, Nippon Genetics, Tokyo). The reaction conditions were as  
6  
7 follows: IFNGR1, after 1 min of denaturation at 94°C, 35 cycles of PCR were performed  
8  
9 (94°C for 30 s and 68°C for 6 min), followed by a final 6 min extension at 68°C; IFNGR2,  
10  
11 TNFR1, TNFR2, and GAPDH: after 1 min of denaturation at 94°C, 35 cycles of PCR were  
12  
13 performed (94°C for 30 s and 68°C for 3 min), followed by a final 3 min extension at 68°C.  
14  
15 The amplified products were electrophoresed in ethidium bromide-stained 1.5% agarose gel.  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26

### 27 **Enzyme-linked immunosorbent assay (ELISA)**

28  
29 We treated NS-SV-DC and NS-SC-AC cells with 1000 IU/mL IFN- $\alpha$ , 10 ng/mL IFN- $\gamma$ , 10  
30  
31 ng/mL TNF- $\alpha$ , and 1 ng/mL IL-1 $\beta$  for 6, 12, or 24 h. The concentrations of CXCL9, CXCL10,  
32  
33 and CXCL11 were each determined with commercially available ELISA kits (R&D Systems)  
34  
35 according to the manufacturer's instructions. Optimal absorbance was read at 450 nm in a  
36  
37 microtiter plate reader. The CXCR3 ligand protein levels were determined using the standard  
38  
39 curve prepared for each assay.  
40  
41  
42  
43  
44  
45  
46  
47

48 We performed blocking experiments using specific transcription factors. NS-SV-DC  
49  
50 cells were incubated with IFN- $\gamma$  (10 ng/mL) in the presence or absence of the STAT1  
51  
52 inhibitor, MTA (1 mM), or the NF- $\kappa$ B inhibitor, Bay11-7082 (10  $\mu$ M), for 24 h. NS-SV-AC  
53  
54 cells were incubated with TNF- $\alpha$  (10 ng/mL) in the presence or absence of Bay11-7082 (10  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3  $\mu\text{M}$ ) or MTA (1 mM) for 24 h. CXCL10 secretion was analyzed by ELISA kits. Optimal  
4  
5  
6  
7  
8  
9 absorbance was read at 450 nm in a microtiter plate reader.

### 10 **Protein isolation and Western blot analysis**

11  
12 NS-SV-DC cells were treated with 10 ng/mL IFN- $\gamma$  for 1, 5, 10, 30, 60, 120, or 360 min, and  
13  
14  
15 NS-SV-AC cells were treated with 10 ng/mL TNF- $\alpha$  for 5, 15, 30, 60, or 120 min. Whole-cell  
16  
17  
18 lysates were prepared using M-PER lysis solution (Thermo Fisher Scientific, Waltham, MA)  
19  
20  
21 supplemented with Halt<sup>TM</sup> Protease Inhibitor Cocktail (Thermo Fisher Scientific). Nuclear  
22  
23  
24 extracts were prepared using NE-PER Nuclear and Cytoplasmic Extraction Reagent (Thermo  
25  
26  
27 Fisher Scientific). Cytosolic extracts (20  $\mu\text{g}$ ) were subjected to electrophoresis on 10%  
28  
29  
30 sodium dodecyl sulfate (SDS)-polyacrylamide gels (Bio-Rad, Hercules, CA), and then  
31  
32  
33 transferred onto nitrocellulose membranes. The membranes were blocked with 3% bovine  
34  
35  
36 serum albumin and incubated with each of the following antibodies (all from Cell Signaling  
37  
38  
39 Technology, Beverly, MA; diluted at 1:1000): anti-janus kinase1 (JAK1), anti-Phospho-JAK1,  
40  
41  
42 anti-JAK2, anti-Phospho-JAK2, anti-STAT1, anti- Phospho-STAT1, anti-p65, anti-inhibitor of  
43  
44  
45  $\kappa\text{B}$  ( $\text{I}\kappa\text{B}$ )- $\alpha$ , anti-Phospho- $\text{I}\kappa\text{B}$ - $\alpha$ , and anti- $\beta$ -actin. After intervening rinses with TBS-T, the  
46  
47  
48 IgG secondary antibodies (Cell Signaling Technology; diluted at 1:1000) were used for the  
49  
50  
51 respective primary antibodies. The immune complexes were visualized by enhanced  
52  
53  
54 chemiluminescence (ECL) Western Blotting Detection Reagent (GE Healthcare,  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 Buckinghamshire, UK). The density of the visualized immune complexes was digitized using  
2  
3 an Amersham Imager 600 (GE Healthcare).  
4  
5  
6  
7

## 8 **Statistical analysis**

9  
10  
11 The data were analyzed with the use of the non-parametric Mann-Whitney U-test. SPSS  
12  
13 Statistics-15.0 J software was used for the analyses. Statistical tests were two-sided, and  
14  
15  
16  
17  
18 significance was defined as a *p*-value <0.05.  
19  
20  
21  
22  
23

## 24 **RESULTS**

### 25 26 27 **The expressions of CXCL9, CXCL10, and CXCL11 mRNA and protein following IFN- $\alpha$ , 28 29 30 IFN- $\gamma$ , TNF- $\alpha$ , or IL-1 $\beta$ stimulation in NS-SV-DC and NS-SV-AC cells**

31  
32  
33 We attempted to identify the factors that regulate the expressions of CXCL9, CXCL10, and  
34  
35  
36 CXCL11, using NS-SV-DC and NS-SV-AC cells. The CXCL10 gene has been reported to  
37  
38  
39 contain response elements for STAT1 and NF- $\kappa$ B [19]. We therefore selected IFN- $\alpha$ , IFN- $\gamma$ ,  
40  
41  
42 TNF- $\alpha$ , and IL-1 $\beta$ . Indeed, these cytokines were reported to be associated with pSS [10, 13–  
43  
44  
45 15]. We conducted a qRT-PCR to examine the expression levels of CXCL9, CXCL10, and  
46  
47  
48 CXCL10 mRNAs following IFN- $\alpha$ , IFN- $\gamma$ , TNF- $\alpha$ , or IL-1 $\beta$  stimulation. As shown in Figure  
49  
50  
51 1a, significant increases in the expressions of CXCL9, CXCL10, and CXCL11 mRNAs were  
52  
53  
54 observed in NS-SV-DC cells after treatment with IFN- $\gamma$  compared to the unstimulated control  
55  
56  
57 cells. At 12 h of treatment, the mRNA levels of CXCL9, CXCL10, and CXCL11 were  
58  
59  
60  
61  
62  
63  
64  
65

1 24352.8-, 3026.4-, and 712.5-fold higher than the basal levels, respectively. In the case of  
2  
3 TNF- $\alpha$  treatment, an increase in the expression of CXCL10 mRNA was detected only in  
4  
5 NS-SV-AC cells (Fig. 1a). On the other hand, IFN- $\alpha$  had a marginal effect on the expression  
6  
7 of only CXCL11 mRNAs in NS-SV-DC cells, and no effect in NS-SV-AC cells. IL-1 $\beta$  had no  
8  
9 effect on the expression of these CXCR3 ligand mRNAs in either cell line (Fig. 1a).  
10  
11  
12  
13

14 To further evaluate the expressions of CXCL9, CXCL10, and CXCL11 proteins, we  
15  
16 detected the protein levels by ELISA using the conditioned medium derived from NS-SV-DC  
17  
18 and NS-SV-AC cells treated with IFN- $\alpha$ , IFN- $\gamma$ , TNF- $\alpha$ , or IL-1 $\beta$ . The NS-SV-DC cells  
19  
20 showed a significantly increased expression of the CXCL10 proteins after IFN- $\gamma$  stimulation  
21  
22 (Fig. 1b). The NS-SV-AC cells showed an increase in the expression of CXCL10 proteins  
23  
24 after TNF- $\alpha$  stimulation (Fig. 1b). IFN- $\alpha$  and IL-1 $\beta$  had no significant effect on the  
25  
26 expressions of these CXCR3 ligand proteins in either cell line (Fig. 1b). These results indicate  
27  
28 that ductal cells play an important role in the secretion of CXCR3 ligand chemokines,  
29  
30 especially CXCL10, in salivary gland cells. In addition, our findings demonstrated that IFN- $\gamma$   
31  
32 and TNF- $\alpha$  are the key cytokines in CXCL10 production in salivary gland cells.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

#### 48 **The detection of IFN- $\gamma$ receptor and TNF- $\alpha$ receptor mRNA in NS-SV-DC and** 49 **NS-SV-AC cells** 50 51

52 NS-SV-DC and NS-SV-AC cells were examined for the presence of mRNA for IFN- $\gamma$   
53  
54 receptor (IFNGR) and TNF- $\alpha$  receptor (TNFR). IFNGR consists of two subunits: IFNGR1  
55  
56  
57  
58  
59  
60

1 (also known as the IFN- $\gamma$  receptor  $\alpha$  chain) and IFNGR2 (also known as the IFN- $\gamma$  receptor  $\beta$   
2 chain) [20]. TNFR consists of two subunits: a 55 kDa type1 receptor (TNFR1) and a 75 kDa  
3 type2 receptor (TNFR2) [21]. As shown in Figure 2, NS-SV-DC and NS-SV-AC cells  
4  
5 expressed the mRNAs for IFNGR1, IFNGR2, TNFR1 and TNFR2 almost same at the intense  
6  
7 level in both cells.  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17

### 18 **The effect of IFN- $\gamma$ on the CXCL10 production in NS-SV-DC cells and the effect of** 19 20 **TNF- $\alpha$ on CXCL10 production in NS-SV-AC cells**

21  
22 We next conducted an ELISA assay to investigate the effect by the density of cytokines on the  
23  
24 CXCL10 production in each cell line. As shown in Figure 3a, the CXCL10 production in  
25  
26 NS-SV-DC cells was increased by the treatment with IFN- $\gamma$  in a dose-dependent manner.  
27  
28 Although the CXCL10 production in NS-SV-AC cells was increased by the treatment with  
29  
30 TNF- $\alpha$  (>10 ng/mL), there was no dose-dependent increase (Fig. 3b). These results suggested  
31  
32 that NS-SV-DC cells had a greater ability to produce CXCL10 compared to NS-SV-AC cells.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44

### 45 **The involvement of STAT1 and NF- $\kappa$ B in CXCL10 production in NS-SV-DC and** 46 47 **NS-SV-AC cells**

48  
49 The CXCL10 gene has been reported to contain response elements for STAT1 and NF- $\kappa$ B  
50  
51 [19]. Accordingly, we next examined the involvement of the two transcription factors STAT1  
52  
53 and NF- $\kappa$ B in the production of CXCL10. We did this by using a STAT1 inhibitor, MTA [18],  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 and an NF- $\kappa$ B inhibitor, Bay11-7082 [17]. As shown in Figure 4a, the addition of MTA (1  
2 mM) resulted in a significant decrease in the IFN- $\gamma$ -induced CXCL10 production in  
3 NS-SV-DC cells ( $p < 0.05$ ). Bay11-7082 also significantly decreased the IFN- $\gamma$ -induced  
4 CXCL10 secretion ( $p < 0.05$ ) (Fig. 4a). In contrast, the TNF- $\alpha$ -induced CXCL10 production in  
5 NS-SV-AC cells was significantly reduced by Bay11-7082 ( $p < 0.05$ ) but not by MTA (Fig.  
6 4b). These data suggest that in NS-SV-DC cells, IFN- $\gamma$  enhanced the CXCL10 production via  
7 both the JAK/STAT1 pathway and the NF- $\kappa$ B pathway, whereas in NS-SV-AC cells, TNF- $\alpha$   
8 enhanced the CXCL10 production via the NF- $\kappa$ B pathway but not the JAK/STAT1 pathway.  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26

**IFN- $\gamma$  promoted the phosphorylations of JAK1, STAT1, and I $\kappa$ B- $\alpha$  in NS-SV-DC cells,  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65**

To clarify the signal transduction pathway activated by IFN- $\gamma$  in NS-SV-DC cells, we  
examined the expression levels of JAK1, JAK2, STAT1, phospho-JAK1, phospho-JAK2,  
phospho-STAT1, p65, I $\kappa$ B- $\alpha$ , and phospho-I $\kappa$ B- $\alpha$ . The Western blot analysis demonstrated  
that IFN- $\gamma$  induced the phosphorylations of JAK2 and STAT1 (Fig. 5a); however, JAK1 was  
not observed in NS-SV-DC cells (data not shown). In addition, IFN- $\gamma$  promoted the  
phosphorylation of I $\kappa$ B- $\alpha$ , and p65 was shifted to the nucleus in NS-SV-DC cells (Fig. 5b).

We then investigated the signal transduction pathway activated by TNF- $\alpha$  in  
NS-SV-AC cells. The expression levels of p65, I $\kappa$ B- $\alpha$ , and phospho-I $\kappa$ B- $\alpha$  were examined.  
The Western blot analysis showed that I $\kappa$ B- $\alpha$  was degraded by TNF- $\alpha$ , and p65 was shifted to

1 the nucleus in NS-SV-AC cells (Fig. 5c). These results indicate that IFN- $\gamma$  promotes the  
2 phosphorylation of JAK2 and I $\kappa$ B- $\alpha$ , thereby resulting in the activation of both the  
3  
4  
5 JAK/STAT1 pathway and the NF- $\kappa$ B pathway in NS-SV-DC cells, whereas TNF- $\alpha$  promotes  
6  
7  
8 the phosphorylation of I $\kappa$ B- $\alpha$ , thereby resulting in the activation of the NF- $\kappa$ B pathway in  
9  
10  
11 NS-SV-AC cells.  
12  
13  
14  
15  
16  
17

## 18 **DISCUSSION**

19  
20  
21 SS is a chronic autoimmune disease characterized by lymphocytic infiltration and the  
22  
23  
24 destruction of salivary and lacrimal glands [3, 4]. Histologically, SS is characterized by an  
25  
26  
27 extensive target tissue infiltration of Th cells, predominantly CD4<sup>+</sup> Th cells but also CD8<sup>+</sup> Th  
28  
29  
30 cells [5]. Based on the expression patterns of cytokines, it has been shown that CD4<sup>+</sup> Th cells  
31  
32  
33 are distributed in Th1 and Th2 cells. Accumulated evidence indicates a close relationship  
34  
35  
36 between the cytokine expression in salivary gland tissues and the development and  
37  
38  
39 progression of pSS. There is supportive evidence that leukocyte activation and elicitation  
40  
41  
42 through a cytokine-driven network is one of the key processes in the autoimmune response  
43  
44  
45 [22]. A 2001 review indicated that chemokines are one of the families of effector molecules  
46  
47  
48 under the control of the cytokine-driven network [22].  
49  
50

51 Chemokines are a superfamily of cytokines that regulate immune cell migration under  
52  
53  
54 both inflammatory and normal physiological conditions [23]. The interactions between  
55  
56  
57 chemokines and their receptors play important roles in the induction of a selective local  
58  
59  
60

1 infiltration of specific cells in various diseases [24]. The CXCR3 ligand chemokines (CXCL9,  
2 CXCL10, and CXCL11) are dual-function chemokines induced by IFN- $\gamma$ , and they are  
3 produced during Th1-type immune responses [2] by diverse cell types, including neutrophils,  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

The expression of CXCR3 and CXCR3 targeting chemokines has been demonstrated to be involved in several diseases and pathological states, including sarcoidosis, viral infections, allograft rejection, rheumatoid arthritis, diabetes, and glomerulonephritis [26–29]. Regarding SS, it was reported that CXCR3 ligand chemokines were expressed in the salivary glands of pSS [6, 7], and that the downregulation of CXCL10 reduced the mononuclear cell infiltration in MRL/lpr mice [8]. Although CXCR3 ligand chemokines and CXCR3 appear to contribute to the pathogenesis of pSS, the clinical significance of chemokines and their receptors and the underlying molecular mechanisms are not yet known.

In the present study, we performed an *in vitro* experiment using NS-SV-DC and NS-SV-AC cells to identify the precise molecular mechanism involved in the regulation of CXCL9, CXCL10, and CXCL11 in salivary gland cells. We selected IFN- $\alpha$ , IFN- $\gamma$ , TNF- $\alpha$ , and IL-1 $\beta$  as reagents that may regulate the expression of CXCR3 ligands. The CXCL10 gene has been reported to contain NF- $\kappa$ B and STAT1-responsible elements in the promoter regions [19]. Interestingly, our study demonstrated that the regulator of the enhancement of CXCR3 ligand chemokines was different between NS-SV-DC and NS-SV-AC cells. Although both cells expressed IFN- $\gamma$  receptor and TNF- $\alpha$  receptor, we detected a significant increase in the

1 expression of CXCL10 in NS-SV-DC cells after IFN- $\gamma$  treatment, whereas the increase in the  
2  
3 expression of CXCL10 in NS-SC-AC cells was detected after TNF- $\alpha$  treatment. These results  
4  
5 indicated that IFN- $\gamma$  and TNF- $\alpha$  are the key cytokines in CXCL10 production in salivary  
6  
7 gland cells. Our findings also demonstrated that NS-SV-DC cells had a greater ability to  
8  
9 produce CXCL10 compared to NS-SV-AC cells. These findings are in accord with the  
10  
11 histopathological images of pSS that show the periductal infiltration of T cells. The CXCL10  
12  
13 secretion from ductal cells by IFN- $\gamma$  stimulation may be caused by an infiltration of CXCR3<sup>+</sup>  
14  
15 T cells in pSS lesion.  
16  
17  
18  
19  
20  
21  
22  
23

24 The blocking experiments using specific transcription factors revealed that a specific  
25  
26 inhibitor of the STAT1 pathway (MTA) decreased the CXCL10 in IFN- $\gamma$ -treated NS-SC-DC  
27  
28 cells. The specific inhibitor of the NF- $\kappa$ B pathway (Bay11-7082) also decreased the CXCL10  
29  
30 in IFN- $\gamma$ -treated NS-SV-DC cells. These results are consistent with those of a study reporting  
31  
32 that IFN- $\gamma$  potentiates the TNF- $\alpha$ -induced CXCL10 production in human monocytes by  
33  
34 increasing the activation of STAT1 and NF- $\kappa$ B through the JAK1 and JAK2 pathways (i.e.,  
35  
36 by crosstalk between IFN- $\gamma$  and TNF- $\alpha$  signal transduction) [30]. In contrast, the specific  
37  
38 inhibitor of the NF- $\kappa$ B pathway (Bay11-7082) but not the specific inhibitor of the STAT1  
39  
40 pathway (MTA) decreased the CXCL10 in TNF- $\alpha$ -treated NS-SV-AC cells. Our findings  
41  
42 demonstrated that the IFN- $\gamma$ -induced CXCL10 expression was mediated by the JAK2/STAT1  
43  
44 and NF- $\kappa$ B pathways in NS-SV-DC cells, whereas the TNF- $\alpha$ -induced CXCL10 expression  
45  
46 was transduced by the NF- $\kappa$ B pathway in NS-SV-AC cells.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3 In conclusion, the results of this study demonstrated that the regulator of the  
4  
5 enhancement of CXCL10 was different between salivary ductal cells and acinar cells, and that  
6  
7 salivary ductal cells had a higher ability to produce CXCL10 compared to acinar cells. The  
8  
9 enhanced production of CXCL10 by IFN- $\gamma$  from ductal cells may result in the migration of  
10  
11 CXCR3<sup>+</sup> T cells. This may be one of the key processes in the inflammation of pSS lesions.  
12  
13  
14  
15  
16  
17  
18  
19  
20

## 21 **ACKNOWLEDGMENTS**

22  
23  
24 This work was supported by Grants-in-Aid for Scientific Research from the Japan Ministry of  
25  
26 Education, Culture, Sports, Science and Technology (nos. 17K11842 and 17K17258).  
27  
28  
29  
30  
31  
32

## 33 **COMPLIANCE WITH ETHICAL STANDARDS**

### 34 35 36 **Conflicts of interest:**

37  
38  
39 None of the authors has any potential financial conflict of interest related to this manuscript.  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## REFERENCES

1. Adams DH, Lloyd AR. 1997. Chemokines: Leucocyte recruitment and activation cytokines. *Lancet* 15: 490–495.
2. Groom JR, Luster AD. 2011. CXCR3 ligands: Redundant, collaborative and antagonistic functions. *Immunol Cell Biol* 89: 207–215.
3. Alspaugh MA, Whaley K. 1981. Sjögren's syndrome, *Textbook of Rheumatology*, Edited by Kelley WN, Harris ED, Ruddy S, Sledge CB. Philadelphia, W.B. Saunders: 971–999.
4. Daniel TE. 1984. Labial salivary gland biopsy in Sjögren's syndrome: Assessment as a diagnostic criterion in 362 suspected cases. *Arthritis Rheum* 27: 147–156.
5. Adamson TC 3rd, Fox RI, Frisman DM, Howell FV. 1983. Immunohistologic analysis of lymphoid infiltrates in primary Sjögren's syndrome using monoclonal antibodies. *J Immunol* 130: 203–208.
6. Ogawa N, Ping L, Zhenjun L, Takada Y, Sugai S. 2002. Involvement of the interferon- $\gamma$ -induced T cell-attracting chemokines, interferon- $\gamma$ -inducible 10-kD protein (CXCL10) and monokine induced by interferon- $\gamma$  (CXCL9), in the salivary gland lesions of patients with Sjögren's syndrome. *Arthritis Rheum* 46: 2730–2741.
7. Ogawa N, Kawanami T, Shimoyama K, Ping L, Sugai S. 2004. Expression of interferon-inducible T cell alpha chemoattractant (CXCL11) in the salivary glands of patients with Sjögren's syndrome. *Clin Immunol* 112: 235–238.
8. Hasegawa H, Inoue A, Kohno M, Muraoka M, Miyazaki T, Terada M, Nakayama T, Yoshie O, Nose M, Yasukawa M. 2006. Antagonist of interferon-inducible protein

10/CXCL10 ameliorates the progression of autoimmune sialadenitis in MRL/lpr mice.

1  
2  
3 Arthritis Rheum 54: 1174–1183.  
4

5  
6 9. Rauch I, Rosebrock F, Hainzl E, Heider S, Majoros A, Wienerroither S, Strobl B,  
7

8  
9 Stockinger S, Kenner L, Müller M, Decker T. 2015. Noncanonical effects of IRF9 in  
10

11  
12 intestinal inflammation: More than Type I and Type III interferons. Mol Cell Bio 35: 2332–  
13

14  
15 2343.  
16

17  
18 10. Hall JC, Baer AN, Shah AA, Criswell LA, Shiboski CH, Rosen A, Casciola-Rosen L.  
19

20  
21 2015. Molecular subsetting of interferon pathways in Sjögren's syndrome. Arthritis  
22

23  
24 Rheumatol 67: 2437–2446.  
25

26  
27 11. Hjelmervik TO, Petersen K, Jonassen I, Jonsson R, Bolstad AI. 2005. Gene expression  
28

29  
30 profiling of minor salivary glands clearly distinguishes primary Sjogren's syndrome  
31

32  
33 patients from healthy control subjects. Arthritis Rheum 52: 1534–1544.  
34

35  
36 12. Pérez P, Anaya JM, Aguilera S, Urzúa U, Munroe D, Molina C, Hermoso MA, Cherry JM,  
37

38  
39 Alliende C, Olea N, Ruiz-Narváez E, González MJ. 2009. Gene expression and  
40

41  
42 chromosomal location for susceptibility to Sjögren's syndrome. J Autoimmun 33: 99–108.  
43

44  
45 13. Nocturne G, Mariette X. 2013. Advances in understanding the pathogenesis of primary  
46

47  
48 Sjögren's syndrome. Nat Rev Rheumatol 9: 544–556.  
49

50  
51 14. Fox RI, Kang HI, Ando D, Abrams J, Pisa E. 1994. Cytokine mRNA expression in  
52

53  
54 salivary gland biopsies of Sjögren's syndrome. J Immunol 152: 5529–5532.  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
15. Yamada A, Arakaki R, Kudo Y, Ishimaru N. 2013. Targeting IL-1 in Sjögren's syndrome. *Expert Opin Ther Targets* 17: 393–401.
  16. Azuma M, Tamatani T, Kasai Y, Sato M. 1993. Immortalization of normal human salivary gland cells with duct-, myoepithelial-, acinar-, or squamous phenotype by transfection of SV40 ori-mutant DNA. *Lab Invest* 69: 24–42.
  17. Pierce JW, Schoenleber R, Jesmok G, Best J, Moore SA, Collins T, Gerritsen ME. 1997. Novel inhibitors of cytokine-induced I $\kappa$ B $\alpha$  phosphorylation and endothelial cell adhesion molecule expression show anti-inflammatory effects in vivo. *J Biol Chem* 272: 21096–21103.
  18. Mowen KA, Tang J, Zhu W, Schurter BT, Shuai K, Herschman HR, David M. 2001. Arginine methylation of STAT1 modulates IFN $\alpha$ / $\beta$ -induced transcription. *Cell* 104: 731–741.
  19. Majumder S, Zhou LZ, Chaturvedi P, Babcock G, Aras S, Ransohoff RM. 1998. p48/STAT-1 $\alpha$ -containing complexes play a predominant role in induction of IFN- $\gamma$ -inducible protein, 10 kDa (IP-10) by IFN- $\gamma$  alone or in synergy with TNF- $\alpha$ . *J Immunol* 161: 4736–4744.
  20. Bach EA, Aguet M, Schreiber RD. 1997. The IFN gamma receptor: a paradigm for cytokine receptor signaling. *Annu Rev Immunol* 15: 563–591.
  21. Dembic Z, Loetscher H, Gubler U, Pan YC, Lahm HW, Gentz R, Brockhaus M, Lesslauer W. 1990. Two human TNF receptors have similar extracellular, but distinct intracellular, domain sequences. *Cytokine* 2: 231–237.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
22. Godessart N, Kunkel SL. 2001. Chemokines in autoimmune disease. *Curr Opin Immunol* 13: 670–675.
23. Kelner GS, Kennedy J, Bacon KB, Kleyensteuber S, Largaespada DA, Jenkins NA, Copeland NG, Bazan JF, Moore KW, Schall TJ, et al. 1994. Lymphotoctin: A cytokine that represents a new class of chemokine. *Science* 266: 1395–1399.
24. Murdoch C, Finn A. 2000. Chemokine receptors and their role in inflammation and infectious diseases. *Blood* 95: 3032–3041.
25. Shin SY, Nam JS, Lim Y, Lee YH. 2010. TNF $\alpha$ -exposed bone marrow-derived mesenchymal stem cells promote locomotion of MDA-MB-231 breast cancer cells through transcriptional activation of CXCR3 ligand chemokines. *J Biol Chem* 285: 30731–30740.
26. Agostini C, Cassatella M, Zambello R, Trentin L, Gasperini S, Perin A, Piazza F, Siviero M, Facco M, Dziejman M, Chilosi M, Qin S, Luster AD, Semenzato G. 1998. Involvement of the IP-10 chemokine in sarcoid granulomatous reactions. *J Immunol* 161: 6413–6420.
27. Agostini C, Facco M, Siviero M, Carollo D, Galvan S, Cattelan AM, Zambello R, Trentin L, Semenzato G. 2000. CXC chemokines IP-10 and mig expression and direct migration of pulmonary CD8<sup>+</sup>/CXCR3<sup>+</sup> T cells in the lungs of patients with HIV infection and T-cell alveolitis. *Am J Respir Crit Care Med* 162: 1466–1473.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
28. Agostini C, Calabrese F, Rea F, Facco M, Tosoni A, Loy M, Binotto G, Valente M, Trentin L, Semenzato G. 2001. CXCR3 and its ligand CXCL10 are expressed by inflammatory cells infiltrating lung allografts and mediate chemotaxis of T cells at sites of rejection. *Am J Pathol* 158: 1703–1711.
  29. Ruschpler P, Lorenz P, Eichler W, Koczan D, Hänel C, Scholz R, Melzer C, Thiesen HJ, Stiehl P. 2003. High CXCR3 expression in synovial mast cells associated with CXCL9 and CXCL10 expression in inflammatory synovial tissues of patients with rheumatoid arthritis. *Arthritis Res Ther* 5: R241–R252.
  30. Qi XF, Kim DH, Yoon YS, Jin D, Huang XZ, Li JH, Deung YK, Lee KJ. 2009. Essential involvement of cross-talk between IFN- $\gamma$  and TNF- $\alpha$  in CXCL10 production in human THP-1 monocytes. *J Cell Physiol* 220: 690–697.

## FIGURE LEGENDS

1  
2  
3 **Fig. 1. The expressions of CXCL9, CXCL10, and CXCL11 mRNA and protein following**

4  
5 **IFN- $\alpha$ , IFN- $\gamma$ , TNF- $\alpha$  or IL-1 $\beta$  stimulation in NS-SV-DC and NS-SV-AC cells. a:**

6  
7  
8  
9 NS-SV-DC and NS-SC-AC cells were treated with IFN- $\alpha$  (1000 IU/mL), IFN- $\gamma$  (10 ng/mL),

10  
11  
12 TNF- $\alpha$  (10 ng/mL) or IL-1 $\beta$  (1 ng/mL) for 6, 12, or 24 h. Total RNAs were prepared, and the

13  
14  
15 quantification of mRNA levels was evaluated by qRT-PCR. Unstimulated cells were used as a

16  
17  
18 control. Bar: mean $\pm$ SD. **b:** NS-SV-DC and NS-SC-AC cells were treated with IFN- $\alpha$  (1000

19  
20  
21 IU/mL), IFN- $\gamma$  (10 ng/mL), TNF- $\alpha$  (10 ng/mL) or IL-1 $\beta$  (1 ng/mL) for 6, 12, or 24 h. An

22  
23  
24 ELISA was performed to measure CXCL9, CXCL10, and CXCL11 protein in the conditioned

25  
26  
27 medium. Unstimulated cells were used as a control. Bar: mean $\pm$ SD.  
28  
29  
30  
31  
32

33 **Fig. 2. The detection of IFN- $\gamma$  receptor and TNF- $\alpha$  receptor mRNA in NS-SV-DC and**

34  
35  
36 **NS-SV-AC cells.** NS-SV-DC and NS-SV-AC cells were examined for the presence of

37  
38  
39 mRNAs for IFNGR1, IFNGR2, TNFR1, and TNFR2 by RT-PCR.

40  
41  
42 Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) was used as housekeeping gene. The

43  
44  
45 mRNAs of IFN- $\gamma$  receptor and TNF- $\alpha$  receptor were detected in both cells.  
46  
47  
48  
49  
50

51 **Fig. 3. The effect of IFN- $\gamma$  on the CXCL10 production in NS-SV-DC cells and the effect**

52  
53  
54 **of TNF- $\alpha$  on the CXCL10 production in NS-SV-AC cells. a:** NS-SV-DC cells were treated

55  
56  
57 with 1, 10, 50, or 100 ng/ml of IFN- $\gamma$  for 24 h. An ELISA was performed to measure  
58  
59  
60  
61  
62  
63  
64  
65

1 CXCL10 protein in the conditioned medium. Bar: mean±SD. \* $p$ <0.05. **b:** NS-SV-AC cells  
2  
3 were treated with 1, 10, 50, or 100 ng/ml of TNF- $\alpha$  for 24 h. An ELISA was performed to  
4  
5 measure CXCL10 protein in the conditioned medium. Bar: mean±SD. \* $p$ <0.05.  
6  
7  
8  
9  
10


11 **Fig. 4. The involvement of STAT1 and NF- $\kappa$ B in the CXCL10 production in NS-SV-DC**  
12 **and NS-SV-AC cells. a:** NS-SV-DC cells were incubated with IFN- $\gamma$  (10 ng/mL) in the  
13  
14 presence or absence of the STAT1 inhibitor MTA (1 mM) or the NF- $\kappa$ B inhibitor Bay11-7082  
15  
16 (10  $\mu$ M) for 24 h. CXCL10 secretion was analyzed with ELISA kits. Optimal absorbance was  
17  
18 read at 450 nm in a microtiter plate reader. The IFN- $\gamma$ -induced CXCL10 production in  
19  
20 NS-SV-DC cells was significantly reduced by Bay11-7082 and MTA. Bar: mean±SD.  
21  
22 \* $p$ <0.05. **b:** NS-SV-AC cells were incubated with TNF- $\alpha$  (10 ng/mL) in the presence or  
23  
24 absence of Bay11-7082 (10  $\mu$ M) or MTA (1 mM) for 24 h. The TNF- $\alpha$ -induced CXCL10  
25  
26 production in NS-SV-AC cells was significantly reduced by Bay11-7082, but not by MTA.  
27  
28 CXCL10 secretion was analyzed with ELISA kits. Optimal absorbance was read at 450 nm in  
29  
30 a microtiter plate reader. Bar: mean±SD. \* $p$ <0.05. ND: not determined.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

48 **Fig. 5. IFN- $\gamma$  promoted the phosphorylations of JAK1, STAT1, and I $\kappa$ B- $\alpha$  in NS-SV-DC**  
49 **cells, whereas TNF- $\alpha$  promoted the phosphorylation of I $\kappa$ B- $\alpha$  in NS-SV-AC cells. a:**  
50  
51 NS-SV-DC cells were treated with 10 ng/mL of IFN- $\gamma$ . The Western blot analysis showed that  
52  
53 phospho-JAK2 and phospho-STAT1 were significantly increased after IFN- $\gamma$  stimulation. **b:**  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65




1 NS-SV-DC cells were treated with 10 ng/mL of IFN- $\gamma$ . IFN- $\gamma$  promoted the phosphorylation  
2  
3 of I $\kappa$ B- $\alpha$ , and p65 was shifted to the nucleus in NS-SV-DC cells. **c:** NS-SV-AC cells were  
4  
5 treated with 10 ng/mL of TNF- $\alpha$ . The Western blot analysis showed that I $\kappa$ B- $\alpha$  was degraded  
6  
7  
8  
9 by TNF- $\alpha$ , and p65 was shifted to the nucleus in NS-SV-AC cells.  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65


**Figure 1**  
**a**



**b**




**Figure 2**




# Figure 3

a




b




# Figure 4

**a**



IFN- $\gamma$	-	+	+	+
MTA	-	-	+	-
Bay11-7082	-	-	-	+


**b**




TNF- $\alpha$	-	+	+	+
Bay11-7082	-	-	+	-
MTA	-	-	-	+

# Figure 5

**a**



**b**



**c**

