

Polyphenol oxidase from *Pectobacterium atrosepticum*: identification and cloning of gene and characteristics of the enzyme

Gorshkov V., Tarasova N., Gogoleva N., Osipova E., Petrova O., Kovtunov E., Gogolev Y.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2017 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim In the present study, we attempted to elucidate if the harmful phytopathogenic bacteria of *Pectobacterium* genus (*P. atrosepticum*) possess the enzymes for oxidation of phenolic compounds. Polyphenol oxidase (laccase) activity was revealed in *P. atrosepticum* cell lysates. Using bioinformatic analysis, an ORF encoding a putative copper-containing polyphenol oxidase of 241 amino acids with a predicted molecular mass of 25.9 kDa was found. This protein (named Pal1) shares significant level of identity with laccases of a new type described for several bacterial species. Cloning and expression of the *pal1* gene and the analysis of corresponding recombinant protein confirmed that Pal1 possessed laccase activity. The recombinant Pal1 protein was characterized in terms of substrate specificity, kinetic parameters, pH and temperature optimum, sensitivity to inhibitors and metal content. Pal1 demonstrated alkali- and thermo-tolerance. The kinetic parameters K_m and k_{cat} for 2,6-dimethoxyphenol were 0.353 ± 0.062 mM and 98.79 ± 4.9 s⁻¹, respectively. The protein displayed high tolerance to sodium azide, sodium fluoride, NaCl, SDS and cinnamic acid. The transcript level of the *pal1* gene in *P. atrosepticum* was shown to be induced by plant-derived phenolic compound (ferulic acid) and copper sulfate.

<http://dx.doi.org/10.1002/jobm.201700413>

Keywords

laccase, *Pectobacterium atrosepticum*, polyphenol oxidase

References

- [1] Jeon JP, Baldrian P, Murugesan K, Chang YS. Laccase-catalysed oxidations of naturally occurring phenols: from in vivo biosynthetic pathways to green synthetic applications. *Microbiol Biotechnol* 2012; 5:318-32.
- [2] Min K, Feeman C, Kang H, Choi SU. The regulation by phenolic compounds of soil organic matter dynamics under a changing environment. *Biomed Res. Int. Volume* 2015; 2015:1-11.
- [3] Burke RM, Cairney JWG. Laccases and other polyphenol oxidases in ecto- and ericoid mycorrhizal fungi. *Mycorrhiza* 2002; 12:105-16.
- [4] Claus H. Laccases and their occurrence in prokaryotes. *Arch Microbiol* 2003; 179:145-50.
- [5] Feng BZ, Li PQ, Fu L, Yu XM. Exploring laccase genes from plant pathogen genomes: a bioinformatic approach. *Genet Mol Res* 2015; 14:14019-36.
- [6] Alahmad K, Gasmalla MAA, Alyousef H. Efficient methods for polyphenol oxidase production. *NJNFS* 2015; 4:656-9.

- [7] Sharma P, Goel R, Capalash N. Bacterial laccases. *World J Microbiol Biotechnol* 2007; 23:823–32.
- [8] Chandra R, Chowdhary P. Properties of bacterial laccases and their application in bioremediation of industrial wastes. *Environ Sci Processes Impacts* 2015; 17:326–42.
- [9] Bhattacharya A, Sood P, Citovsky V. The roles of plant phenolics in defence and communication during *Agrobacterium* and *Rhizobium* infection. *Mol Plant Pathol* 2010; 11:705–19.
- [10] Daayf F, El Hadrami A, El-Bebany A, Henriquez M. Phenolic compounds in plant defense and pathogen counter-defense mechanisms. In: Cheyner V, Sarni-Manchado P, Quideau S, editors. *Recent advances in polyphenol research*. West Sussex, UK: Wiley-Blackwell; 2012. pp 191–208.
- [11] O'Brien J, Daudi A, Butt V, Bolwell G. Reactive oxygen species and their role in plant defence and cell wall metabolism. *Planta* 2012; 236:765–79.
- [12] Das K, Roychoudhury A. Reactive oxygen species (ROS) and response of antioxidants as ROS-scavengers during environmental stress in plants. *Front Environ Sci* 2014; 2:1–3.
- [13] Gianfreda L, Mora M, Diez M. Restoration of polluted soils by means of microbial and enzymatic processes. *RC Suelo Nutr* 2006; 6:20–40.
- [14] Petrova O, Gorshkov V, Daminova A, Ageeva M. Stress response in *Pectobacterium atrosepticum* SCRI1043 under starvation conditions: adaptive reactions at a low population density. *Res Microbiol* 2014; 165:119–27.
- [15] Petrova O, Gorshkov V, Sergeeva I, Daminova A. Alternative scenarios of starvation-induced adaptation in *Pectobacterium atrosepticum*. *Res Microbiol* 2016; 167:254–61.
- [16] Charkowski A, Blanco C, Condemine G, Expert D. The role of secretion systems and small molecules in soft-rot *Enterobacteriaceae* pathogenicity. *Ann Rev Phytopathol* 2012; 50:425–49.
- [17] Tarasova N, Gorshkov V, Petrova O, Gogolev Y. Potato signal molecules that activate pectate lyase synthesis in *Pectobacterium atrosepticum* SCRI1043. *WJMB* 2013; 29:1189–96.
- [18] Norman-Setterblad C. Interacting signal pathways control defense gene expression in *Arabidopsis* in response to cell wall-degrading enzymes from *Erwinia carotovora*. *Mol Plant Microbe Interact* 2000; 13:430–8.
- [19] Davidsson R. Pathogenicity of and plant immunity to soft rot *pectobacteria*. *Front Plant* 2013; 191:1–3.
- [20] Gorshkov V, Daminova A, Mikshina P, Petrova O. Pathogen-induced conditioning of the primary xylem vessels – a prerequisite for the formation of bacterial emboli by *Pectobacterium atrosepticum*. *Plant Biol* 2016; 18:609–17.
- [21] Joshi J, Burdman S, Lipsky A, Yedidia I. Effects of plant antimicrobial phenolic compounds on virulence of the genus *Pectobacterium*. *Res Microbiol* 2015; 166:535–45.
- [22] Bell KS, Sebahia M, Pritchard L, Holden M. Genome sequence of the enterobacterial phytopathogen *Erwinia carotovora* subsp. *atroseptica* and characterization of virulence factors. *Microbiology* 2004; 101:11105e10.
- [23] Gorshkov V, Daminova A, Ageeva M, Petrova O. Dissociation of a population of *Pectobacterium atrosepticum* SCRI1043 in tobacco plants: formation of bacterial emboli and dormant cells. *Protoplasma* 2014; 251:499–510.
- [24] Altschul SF, Madden TL, Schaffer AA, Zhang J. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 1997; 25:3389–402.
- [25] Tamura K, Stecher G, Peterson D, Filipski A. MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* 2013; 30:2725–9.
- [26] Nei M, Kumar S. *Molecular evolution and phylogenetics*. New York: Oxford University Press; 2000.
- [27] Henikoff S, Henikoff JG. Amino acid substitution matrices from protein blocks. *Proc Natl Acad Sci USA* 1992; 89:10915–9.
- [28] Yang J, Yan R, Roy A, Xu D. The I-TASSER Suite: protein structure and function prediction. *Nat Methods* 2015; 12:7–8.
- [29] Yang J, Zhang Y. I-TASSER server: new development for protein structure and function predictions. *Nucleic Acids Res* 2015; 43:W174–81.
- [30] Roy A, Kucukural A, Zhang Y. I-TASSER: a unified platform for automated protein structure and function prediction. *Nat Protoc* 2010; 5:725–38.
- [31] Zhang Y. I-TASSER server for protein 3D structure prediction. *BMC Bioinf* 2008; 9:40.
- [32] Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970; 227:680–2.
- [33] Reiss R, Ihssen J, Richter M, Eichhorn E. Laccase versus laccase-like multi-copper oxidase: a comparative study of similar enzymes with diverse substrate spectra. *PLoS ONE* 2013; 8:e65633.
- [34] Finn RD, Coggill P, Eberhardt RY, Eddy SR. The Pfam protein families database: towards a more sustainable future. *Nucleic Acids Res* 2016; 44:D279–85.
- [35] Marchler-Bauer A, Derbyshire MK, Gonzales NR, Lu S. CDD: NCBI's conserved domain database. *Nucleic Acids Res* 2015; 43:D222–6.

- [36] Beloqui A, Pita M, Polaina J, Martinez-Arias A. Novel polyphenol oxidase mined from a metagenome expression library of bovine rumen: biochemical properties, structural analysis, and phylogenetic relationships. *J Biol Chem* 2006; 281:22933-42.
- [37] Guo X, Sh Zhou, Wang Y, Song J. Characterization of a highly thermostable and organic solvent-tolerant copper-containing polyphenol oxidase with dye-decolorizing ability from *Kurthia huakuii* LAM0618T. *PLoS ONE* 2016; 11:e0164810.
- [38] Berman HM, Bhat TN, Bourne PE, Feng Z. The protein data bank and the challenge of structural genomics. *Nat Struct Biol* 2000; 7:957-9.
- [39] Chen CY, Hsieh ZSh, Cheepudom J, Yang CH. A 24.7-kDa copper-containing oxidase, secreted by *Thermobifida fusca*, significantly increasing the xylanase/cellulase-catalyzed hydrolysis of sugarcane bagasse. *Appl Microbiol Biotechnol* 2013; 97:8977-86.
- [40] Kim Y, Maltseva N, Dementieva I, Collart F. Crystal structure of hypothetical protein YfiH from *Shigella flexneri* at 2 Å resolution. *Proteins* 2006; 63:1097-101.
- [41] Minasov G, Shuvalova L, Mondragon A, Taneja B. 1.8 Å crystal structure of an uncharacterized *B. stearrowthermophilus* protein. RCSB PDB. Protein Data Bank 2011; <https://doi.org/10.2210/pdb1t8h/pdb>
- [42] Fernandez E, Sanchez-Amat A, Solano A. Location and catalytic characteristics of a multipotent bacterial polyphenol oxidase. *Pigment Cell Res* 1999; 12:331-9.
- [43] Lu Y. Cupredoxins. In: McCleverty J, Meyer T, editors. *Comprehensive coordination chemistry II: from biology to nanotechnology*. New York: Elsevier Ltd; 2003. pp 91-122.
- [44] Messerschmidt A, Robert HR. The blue oxidases, ascorbate oxidase, laccase and ceruloplasmin. Modelling and structural relationships. *Eur J Biochem* 1990; 187:341-52.
- [45] Solano F, Lucas-Elio P, Lopez-Serrano D, Fernandez E. Dimethoxyphenol oxidase activity of different microbial blue multicopper proteins. *FEMS Microb Lett* 2001; 204:175-81.
- [46] Solano F, Sanchez-Amat A. Studies on the phylogenetic relationships of melanogenic marine bacteria: proposal of *Marinomonas mediterranea* sp. nov. *Int J Syst Bacteriol* 1999; 49:1241-6.
- [47] Sherif M, Waung D, Korbeci B, Mavisakalyan V. Biochemical studies of the multicopper oxidase (small laccase) from *Streptomyces coelicolor* using bioactive phytochemicals and site-directed mutagenesis. *Microbiol Biotechnol* 2013; 6:588-97.
- [48] Ruijsenaars HJ, Hartmans S. A cloned *Bacillus halodurans* multicopper oxidase exhibiting alkaline laccase activity. *Appl Microbiol Biotechnol* 2004; 65:177-82.
- [49] Reiss R, Ihssen J, Thony-Meyer L. *Bacillus pumilus* laccase: a heat stable enzyme with a wide substrate spectrum. *BMC Biotechnol* 2011; 11:9.
- [50] Brander S, Mikkelsen JD, Kepp KP, Mikkelsen JD. Characterization of an alkali- and halide-resistant laccase expressed in *E. coli*: CotA from *Bacillus clausii*. *PLoS ONE* 2014; 9:e99402.
- [51] Miyazaki K. A hyperthermophilic laccase from *Thermus thermophilus* HB27. *Extremophiles* 2005; 9:415-25.