

Cerebellar atrophy and changes in cytokines associated with the CACNA1A R583Q mutation in a Russian familial hemiplegic migraine type 1 family

Khaiboullina S., Mendelevich E., Shigapova L., Shagimardanova E., Gazizova G., Nikitin A., Martynova E., Davidyuk Y., Bogdanov E., Gusev O., Van Den Maagdenberg A., Giniatullin R., Rizvanov A.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2017 Khaiboullina, Mendelevich, Shigapova, Shagimardanova, Gazizova, Nikitin, Martynova, Davidyuk, Bogdanov, Gusev, van den Maagdenberg, Giniatullin and Rizvanov. Background: Immune mechanisms recently emerged as important contributors to migraine pathology with cytokines affecting neuronal excitation. Therefore, elucidating the profile of cytokines activated in various forms of migraine, including those with a known genetic cause, can help in diagnostic and therapeutic approaches. Methods: Here we (i) performed exome sequencing to identify the causal gene mutation and (ii) measured, using Bio-Plex technology, 22 cytokines in serum of patients with familial migraine (two with hemiplegic migraine and two with migraine with aura) from a Russian family that ethnically belongs to the Tatar population. MRI scanning was used to assess cerebellar atrophy associated with migraine in mutation carriers. Results: Whole-exome sequencing revealed the R583Q missense mutation in the CACNA1A gene in the two patients with hemiplegic migraine and cerebellar ataxia with atrophy, confirming a FHM1 disorder. Two further patients did not have the mutation and suffered from migraine with aura. Elevated serum levels of pro-inflammatory and pro-nociceptive IL-6 and IL-18 were found in all four patients (compared to a reference panel), whereas pro-apoptotic SCGF- β and TRAIL were higher only in the patients with the FHM1 mutation. Also, cytokines CXCL1, HGF, LIF, and MIF were found particularly high in the two mutation carriers, suggesting a possible role of vascular impairment and neuroinflammation in disease pathogenesis. Notably, some "algesic" cytokines, such as β -NGF and TNF β , remained unchanged or even were down-regulated. Conclusion: We present a detailed genetic, neurological, and biochemical characterization of a small Russian FHM1 family and revealed evidence for higher levels of specific cytokines in migraine patients that support migraine-associated neuroinflammation in the pathology of migraine.

<http://dx.doi.org/10.3389/fncel.2017.00263>

Keywords

Cytokines, FHM1, Inflammation, Migraine, Nociception

References

- [1] Ahola-Olli, A. V., Würtz, P., Havulinna, A. S., Aalto, K., Pitkänen, N., Lehtimäki, T., et al. (2017). Genome-wide association study identifies 27 loci influencing concentrations of circulating cytokines and growth factors. *Am. J. Hum. Genet.* 100, 40–50. doi: 10.1016/j.ajhg.2016.11.007
- [2] Arumugam, M., and Parthasarathy, V. (2016). Reduction of CD4 CD25 regulatory T-cells in migraine: is migraine an autoimmune disorder? *J. Neuroimmunol.* 290, 54–59. doi: 10.1016/j.jneuroim.2015.11.015
- [3] Bo, S. H., Davidsen, E. M., Gulbrandsen, P., Dietrichs, E., Bovim, G., Stovner, L. J., et al. (2009). Cerebrospinal fluid cytokine levels in migraine, tension-type headache and cervicogenic headache. *Cephalgia* 29, 365–372. doi: 10.1111/j.1468-2982.2008.01727.x
- [4] Cantarella, G., Pignataro, G., Di Benedetto, G., Anzilotti, S., Vinciguerra, A., Cuomo, O., et al. (2014). Ischemic tolerance modulates TRAIL expression and its receptors and generates a neuroprotected phenotype. *Cell Death Dis.* 5, e1331. doi: 10.1038/cddis.2014.286
- [5] Cao, D. L., Qian, B., Zhang, Z. J., Gao, Y. J., and Wu, X. B. (2016). Chemokine receptor CXCR2 in dorsal root ganglion contributes to the maintenance of inflammatory pain. *Brain Res. Bull.* 127, 219–225. doi: 10.1016/j.brainresbull.2016.09.016
- [6] Chamorro, Á., Meisel, A., Planas, A. M., Urra, X., van de Beek, D., and Veltkamp, R. (2012). The immunology of acute stroke. *Nat. Rev. Neurol.* 8, 401–410. doi: 10.1038/nrneurol.2012.98
- [7] Cuesta, M. C., Quintero, L., Pons, H., and Suarez-Roca, H. (2002). Substance P and calcitonin gene-related peptide increase IL-1 beta, IL-6 and TNF alpha secretion from human peripheral blood mononuclear cells. *Neurochem. Int.* 40, 301–306. doi: 10.1016/S0097-0186(01)00094-8
- [8] Dorgans, K., Salvi, J., Bertaso, F., Bernard, L., Lory, P., Doussau, F., et al. (2017). Characterization of the dominant inheritance mechanism of Episodic Ataxia type 2. *Neurobiol. Dis.* 106, 110–123. doi: 10.1016/j.nbd.2017.07.004
- [9] Du, X., Wang, J., Zhu, H., Rinaldo, L., Lamar, K. M., Palmenberg, A. C., et al. (2013). Second cistron in CACNA1A gene encodes a transcription factor mediating cerebellar development and SCA6. *Cell* 154, 118–133. doi: 10.1016/j.cell.2013.05.059
- [10] Ducros, A., Denier, C., Joutel, A., Cecillon, M., Lescoat, C., Vahedi, K., et al. (2001). The clinical spectrum of familial hemiplegic migraine associated with mutations in a neuronal calcium channel. *N. Engl. J. Med.* 345, 17–24. doi: 10.1056/NEJM200107053450103
- [11] Ferrari, M. D., Klever, R. R., Terwindt, G. M., Ayata, C., and van den Maagdenberg, A. M. (2015). Migraine pathophysiology: lessons from mouse models and human genetics. *Lancet Neurol.* 14, 65–80. doi: 10.1016/S1474-4422(14)70220-0
- [12] Fletcher, C. F., Lutz, C. M., O'Sullivan, T. N., Shaughnessy, J. D., Jr., Hawkes, R., Frankel, W. N., et al. (1996). Absence epilepsy in tottering mutant mice is associated with calcium channel defects. *Cell* 87, 607–617. doi: 10.1016/S0092-8674(00)81381-1
- [13] Gerring, Z. F., Powell, J. E., Montgomery, G. W., and Nyholt, D. R. (2017). Genome-wide analysis of blood gene expression in migraine implicates immune-inflammatory pathways. *Cephalgia*. doi: 10.1177/0333102416686769. [Epub ahead of print].
- [14] Giunti, P., Mantuano, E., Frontali, M., and Veneziano, L. (2015). Molecular mechanism of Spinocerebellar Ataxia type 6: glutamine repeat disorder, channelopathy and transcriptional dysregulation. The multifaceted aspects of a single mutation. *Front. Cell. Neurosci.* 9:36. doi: 10.3389/fncel.2015.00036
- [15] Gormley, P., Anttila, V., Winsvold, B. S., Palta, P., Esko, T., Pers, T. H., et al. (2016). Meta-analysis of 375,000 individuals identifies 38 susceptibility loci for migraine. *Nat. Genet.* 48, 856–866. doi: 10.1038/ng.3598
- [16] Grinberg, Y. Y., Dibbern, M. E., Levasseur, V. A., and Kraig, R. P. (2013). Insulin-like growth factor-1 abrogates microglial oxidative stress and TNFalpha responses to spreading depression. *J. Neurochem.* 126, 662–672. doi: 10.1111/jnc.12267
- [17] Gyotoku, E., Morita, E., Kameyoshi, Y., Hiragun, T., Yamamoto, S., and Hide, M. (2001). The IL-6 family cytokines, interleukin-6, interleukin-11, oncostatin M, and leukemia inhibitory factor, enhance mast cell growth through fibroblast-dependent pathway in mice. *Arch. Dermatol. Res.* 293, 508–514. doi: 10.1007/PL00007465
- [18] Headache Classification Committee of the International Headache Society (IHS). (2013). The international classification of headache disorders, 3rd edition (beta version). *Cephalgia* 33, 629–808. doi: 10.1177/0333102413485658
- [19] Hiragun, T., Morita, E., Mihara, S., Tanaka, T., Gyotoku, E., Kameyoshi, Y., et al. (2000). Leukemia inhibitory factor enhances mast cell growth in a mast cell/fibroblast co-culture system through stat3 signaling pathway of fibroblasts. *FEBS Lett.* 487, 219–223. doi: 10.1016/S0014-5793(00)02355-3
- [20] Jacobs, B., and Dussor, G. (2016). Neurovascular contributions to migraine: moving beyond vasodilation. *Neuroscience* 338, 130–144. doi: 10.1016/j.neuroscience.2016.06.012
- [21] Jeng, C. J., Chen, Y. T., Chen, Y. W., and Tang, C. Y. (2006). Dominantnegative effects of human P/Q-type Ca₂₊ channel mutations associated with episodic ataxia type 2. *Am. J. Physiol. Cell Physiol.* 290, C1209–C1220. doi: 10.1152/ajpcell.00247.2005

- [22] Khaibullin, T., Ivanova, V., Martynova, E., Cherepnev, G., Khabirov, F., Granatov, E., et al. (2017). Elevated levels of proinflammatory cytokines in cerebrospinal fluid of multiple sclerosis patients. *Front. Immunol.* 8:531. doi: 10.3389/fimmu.2017.00531
- [23] Kichev, A., Rousset, C. I., Baburamani, A. A., Levison, S. W., Wood, T. L., Gressens, P., et al. (2014). Tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) signaling and cell death in the immature central nervous system after hypoxia-ischemia and inflammation. *J. Biol. Chem.* 289, 9430–9439. doi: 10.1074/jbc.M113.512350
- [24] Kilinc, E., Guerrero-Toro, C., Zakharov, A., Vitale, C., Gubert-Olive, M., Koroleva, K., et al. (2017). Serotonergic mechanisms of trigeminal meningeal nociception: implications for migraine pain. *Neuropharmacology* 116, 160–173. doi: 10.1016/j.neuropharm.2016.12.024
- [25] Kraus, R. L., Sinnegger, M. J., Koschak, A., Glossmann, H., Stenirri, S., Carrera, P., et al. (2000). Three new familial hemiplegic migraine mutants affect P/Q-type Ca₂ channel kinetics. *J. Biol. Chem.* 275, 9239–9243. doi: 10.1074/jbc.275.13.9239
- [26] Levite, M. (1998). Neuropeptides, by direct interaction with T cells, induce cytokine secretion and break the commitment to a distinct T helper phenotype. *Proc. Natl. Acad. Sci. U.S.A.* 95, 12544–12549. doi: 10.1073/pnas.95.21.12544
- [27] Levy, D. (2009). Migraine pain, meningeal inflammation, and mast cells. *Curr. Pain Headache Rep.* 13, 237–240. doi: 10.1007/s11916-009-0040-y
- [28] Levy, D., Burstein, R., and Strassman, A. M. (2006). Mast cell involvement in the pathophysiology of migraine headache: a hypothesis. *Headache* 46(Suppl 1), S13–S18. doi: 10.1111/j.1526-4610.2006.00485.x
- [29] Mark, M. D., Krause, M., Boele, H. J., Kruse, W., Pollok, S., Kuner, T., et al. (2015). Spinocerebellar ataxia type 6 protein aggregates cause deficits in motor learning and cerebellar plasticity. *J. Neurosci.* 35, 8882–8895. doi: 10.1523/JNEUROSCI.0891-15.2015
- [30] Nair, A., Simonetti, M., Birsa, N., Ferrari, M. D., van den Maagdenberg, A. M., Giniatullin, R., et al. (2010). Familial hemiplegic migraine Ca(v)2.1 channel mutation R192Q enhances ATP-gated P2X3 receptor activity of mouse sensory ganglion neurons mediating trigeminal pain. *Mol. Pain* 6:48. doi: 10.1186/1744-8069-6-48
- [31] Neeb, L., Hellen, P., Boehnke, C., Hoffmann, J., Schuh-Hofer, S., Dirnagl, U., et al. (2011). IL-1beta stimulates COX-2 dependent PGE(2) synthesis and CGRP release in rat trigeminal ganglia cells. *PLoS ONE* 6:e17360. doi: 10.1371/journal.pone.0017360
- [32] Ophoff, R. A., Terwindt, G. M., Vergouwe, M. N., van Eijk, R., Oefner, P. J., Hoffman, S. M., et al. (1996). Familial hemiplegic migraine and episodic ataxia type-2 are caused by mutations in the Ca channel gene CACNL1A4. *Cell* 87, 543–552. doi: 10.1016/S0092-8674(00)81373-2
- [33] Perini, F., D'Andrea, G., Galloni, E., Pignatelli, F., Billo, G., Alba, S., et al. (2005). Plasma cytokine levels in migraineurs and controls. *Headache* 45, 926–931. doi: 10.1111/j.1526-4610.2005.05135.x
- [34] Pietrobon, D., and Moskowitz, M. A. (2013). Pathophysiology of migraine. *Annu. Rev. Physiol.* 75, 365–391. doi: 10.1146/annurev-physiol-030212-183717
- [35] Pusic, K. M., Pusic, A. D., Kemme, J., and Kraig, R. P. (2014). Spreading depression requires microglia and is decreased by their M2a polarization from environmental enrichment. *Glia* 62, 1176–1194. doi: 10.1002/glia.22672
- [36] Roy, M., Richard, J. F., Dumas, A., and Vallieres, L. (2012). CXCL1 can be regulated by IL-6 and promotes granulocyte adhesion to brain capillaries during bacterial toxin exposure and encephalomyelitis. *J. Neuroinflammation* 9:18. doi: 10.1186/1742-2094-9-18
- [37] Schiller, J. H., Storer, B. E., Witt, P. L., Alberti, D., Tombes, M. B., Arzumanian, R., et al. (1991). Biological and clinical effects of intravenous tumor necrosis factor-alpha administered three times weekly. *Cancer Res.* 51, 1651–1658.
- [38] Tamura, S., Morikawa, Y., Miyajima, A., and Senba, E. (2003). Expression of oncostatin M receptor beta in a specific subset of nociceptive sensory neurons. *Eur. J. Neurosci.* 17, 2287–2298. doi: 10.1046/j.1460-9568.2003.02681.x
- [39] Terwindt, G., Kors, E., Haan, J., Vermeulen, F., Van den Maagdenberg, A., Frants, R., et al. (2002). Mutation analysis of the CACNA1A calcium channel subunit gene in 27 patients with sporadic hemiplegic migraine. *Arch. Neurol.* 59, 1016–1018. doi: 10.1001/archneur.59.6.1016
- [40] Tfelt-Hansen, P., and Olesen, J. (2011). Possible site of action of CGRP antagonists in migraine. *Cephalgia* 31, 748–750. doi: 10.1177/0333102411398403
- [41] Tozzi, A., de Iure, A., Di Filippo, M., Costa, C., Caproni, S., Pisani, A., et al. (2012). Critical role of calcitonin gene-related peptide receptors in cortical spreading depression. *Proc. Natl. Acad. Sci. U.S.A.* 109, 18985–18990. doi: 10.1073/pnas.1215435109
- [42] Uzar, E., Evliyaoglu, O., Yucel, Y., Ugur Cevik, M., Acar, A., Guzel, I., et al. (2011). Serum cytokine and pro-brain natriuretic peptide (BNP) levels in patients with migraine. *Eur. Rev. Med. Pharmacol. Sci.* 15, 1111–1116. Available online at: <http://www.europeanreview.org/wp/wp-content/uploads/1043.pdf>

- [43] van den Maagdenberg, A.M., Haan, J., Terwindt, G.M., and Ferrari, M. D. (2007). Migraine: gene mutations and functional consequences. *Curr. Opin. Neurol.* 20, 299–305. doi: 10.1097/WCO.0b013e3281338d1f
- [44] van den Maagdenberg, A. M., Pietrobon, D., Pizzorusso, T., Kaja, S., Broos, L. A., Cesetti, T., et al. (2004). A Cacna1a knockin migraine mouse model with increased susceptibility to cortical spreading depression. *Neuron* 41, 701–710. doi: 10.1016/S0896-6273(04)00085-6
- [45] Vera, P. L., and Meyer-Siegler, K. L. (2003). Anatomical location of macrophage migration inhibitory factor in urogenital tissues, peripheral ganglia and lumbosacral spinal cord of the rat. *BMC Neurosci.* 4:17. doi: 10.1186/1471-2202-4-17
- [46] Wang, J. G., Strong, J. A., Xie, W., Yang, R. H., Coyle, D. E., Wick, D. M., et al. (2008). The chemokine CXCL1/growth related oncogene increases sodium currents and neuronal excitability in small diameter sensory neurons. *Mol. Pain* 4:38. doi: 10.1186/1744-8069-4-38
- [47] Yan, J., Melemedjian, O. K., Price, T. J., and Dussor, G. (2012). Sensitization of dural afferents underlies migraine-related behavior following meningeal application of interleukin-6 (IL-6). *Mol. Pain* 8:6. doi: 10.1186/1744-8069-8-6
- [48] Yang, L. H., Xu, G.M., and Wang, Y. (2016). Up-regulation of CXCL1 and CXCR2 contributes to remifentanil-induced hypernociception via modulating spinal NMDA receptor expression and phosphorylation in rats. *Neurosci. Lett.* 626, 135–141. doi: 10.1016/j.neulet.2015.12.044
- [49] Zhang, X., Burstein, R., and Levy, D. (2012). Local action of the proinflammatory cytokines IL-1 β and IL-6 on intracranial meningeal nociceptors. *Cephalalgia* 32, 66–72. doi: 10.1177/0333102411430848
- [50] Zhuchenko, O., Bailey, J., Bonnen, P., Ashizawa, T., Stockton, D. W., Amos, C., et al. (1997). Autosomal dominant cerebellar ataxia (SCA6) associated with small polyglutamine expansions in the alpha 1A-voltage-dependent calcium channel. *Nat. Genet.* 15, 62–69. doi: 10.1038/ng0197-62