

PLoS ONE, 2017, vol.12, N9

Optimization of the expression, purification and polymerase activity reaction conditions of recombinant human PrimPol

Boldinova E., Stojkovič G., Khairullin R., Wanrooij S., Makarova A.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2017 Boldinova et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. Human PrimPol is a DNA primase/polymerase involved in DNA damage tolerance and prevents nuclear genome instability. PrimPol is also localized to the mitochondria, but its precise function in mitochondrial DNA maintenance has remained elusive. PrimPol works both as a translesion (TLS) polymerase and as the primase that restarts DNA replication after a lesion. However, the observed biochemical activities of PrimPol vary considerably between studies as a result of different reaction conditions used. To reveal the effects of reaction composition on PrimPol DNA polymerase activity, we tested the polymerase activity in the presence of various buffer agents, salt concentrations, pH values and metal cofactors. Additionally, the enzyme stability was analyzed under various conditions. We demonstrate that the reaction buffer with pH 6–6.5, low salt concentrations and 3 mM Mg²⁺ or 0.3–3 mM Mn²⁺ cofactor ions supports the highest DNA polymerase activity of human PrimPol in vitro. The DNA polymerase activity of PrimPol was found to be stable after multiple freeze-thaw cycles and prolonged protein incubation on ice. However, rapid heat-inactivation of the enzyme was observed at 37°C. We also for the first time describe the purification of human PrimPol from a human cell line and compare the benefits of this approach to the expression in *Escherichia coli* and in *Saccharomyces cerevisiae* cells. Our results show that active PrimPol can be purified from *E. coli* and human suspension cell line in high quantities and that the activity of the purified enzyme is similar in both expression systems. Conversely, the yield of full-length protein expressed in *S. cerevisiae* was considerably lower and this system is therefore not recommended for expression of full-length recombinant human PrimPol.

<http://dx.doi.org/10.1371/journal.pone.0184489>

References

- [1] Bianchi J, Rudd SG, Jozwiakowski SK, Bailey LJ, Soura V, Taylor E, et al. PrimPol bypasses UV photo-products during eukaryotic chromosomal DNA replication. *Mol Cell*. 2013; 52(4):566–573. <https://doi.org/10.1016/j.molcel.2013.10.035> PMID: 24267451
- [2] García-Gómez S, Reyes A, Martínez-Jiménez MI, Chocrón ES, Mourón S, Terrados G, et al. PrimPol, an archaic primase/polymerase operating in human cells. *Mol Cell*. 2013; 52(4):541–553. <https://doi.org/10.1016/j.molcel.2013.09.025> PMID: 24207056

- [3] Wan L, Lou J, Xia Y, Su B, Liu T, Cui J, et al. hPrimpol1/CCDC111 is a human DNA primase-polymerase required for the maintenance of genome integrity. *EMBO Rep.* 2013; 14:1104-1112. <https://doi.org/10.1038/embor.2013.159> PMID: 24126761
- [4] Iyer LM, Koonin EV, Leipe DD, Aravind L. Origin and evolution of the archaeo-eukaryotic primase superfamily and related palm-domain proteins: structural insights and new members. *Nucleic Acids Res.* 2005; 33(12):3875-3896. <https://doi.org/10.1093/nar/gki702> PMID: 16027112
- [5] Rudd SG, Glover L, Jozwiakowski SK, Horn D, Doherty AJ. PPL2 translesion polymerase is essential for the completion of chromosomal DNA replication in the African trypanosome. *Mol Cell.* 2013; 52 (6):554-565.
- [6] Zafar MK, Ketkar A, Lodeiro MF, Cameron CE, Eoff RL. Kinetic analysis of human PrimPol DNA polymerase activity reveals a generally error-prone enzyme capable of accurately bypassing 7,8-dihydro-8-ox-2'-deoxyguanosine. *Biochemistry.* 2014; 53(41):6584-6594. <https://doi.org/10.1021/bi501024u> PMID: 25255211
- [7] Mourón S, Rodríguez-Acebes S, Martínez-Jiménez MI, García-Gómez S, Chocrón S, Blanco L, et al. Repriming of DNA synthesis at stalled replication forks by human PrimPol. *Nat Struct Mol Biol.* 2013; 20 (12):1383-1389. <https://doi.org/10.1038/nsmb.2719> PMID: 24240614
- [8] Mislak AC, Anderson KS. Insights into the Molecular Mechanism of Polymerization and Nucleoside Reverse Transcriptase Inhibitor Incorporation by Human PrimPol. *Antimicrob Agents Chemother.* 2015; 60(1):561-569. <https://doi.org/10.1128/AAC.02270-15> PMID: 26552983
- [9] Schiavone D, Jozwiakowski SK, Romanello M, Guilbaud G, Guillian TA, Bailey LJ, et al. PrimPol Is Required for Replicative Tolerance of G Quadruplexes in Vertebrate Cells. *Mol Cell.* 2016; 61(1):161-169. <https://doi.org/10.1016/j.molcel.2015.10.038> PMID: 26626482
- [10] Stojkovič G, Makarova AV, Wanrooij PH, Forslund J, Burgers PM, Wanrooij S. Oxidative DNA damage stalls the human mitochondrial replisome. *Sci Rep.* 2016; 6:28942. <https://doi.org/10.1038/srep28942> PMID: 27364318
- [11] Keen BA, Bailey LJ, Jozwiakowski SK, Doherty AJ. Human PrimPol mutation associated with high myopia has a DNA replication defect. *Nucleic Acids Res.* 2014; 42(19):12102-12111. <https://doi.org/10.1093/nar/gku879> PMID: 25262353
- [12] Bylund G, Majka J, Burgers PM. Overproduction and purification of RFC-related clamp loaders and PCNA-related clamps from *Saccharomyces cerevisiae*. *Methods in Enzymology.* 2006; 409:1-11. [https://doi.org/10.1016/S0076-6879\(05\)09001-4](https://doi.org/10.1016/S0076-6879(05)09001-4) PMID: 16793392
- [13] Keen BA, Jozwiakowski SK, Bailey LJ, Bianchi J, Doherty AJ. Molecular dissection of the domain architecture and catalytic activities of human PrimPol. *Nucleic Acids Res.* 2014; 42(9):5830-5845. <https://doi.org/10.1093/nar/gku214> PMID: 24682820
- [14] Martínez-Jiménez MI, García-Gómez S, Bebenek K, Sastre-Moreno G, Calvo PA, Díaz-Talavera A. Alternative solutions and new scenarios for translesion DNA synthesis by human PrimPol. *DNA Repair (Amst).* 2015; 29:127-138.
- [15] Costa S, Almeida A, Castro A, Domingues L. Fusion tags for protein solubility, purification and immunogenicity in *Escherichia coli*: the novel Fh8 system. *Front Microbiol.* 2014; 5:63. <https://doi.org/10.3389/fmicb.2014.00063> PMID: 24600443
- [16] Holz C, Prinz B, Bolotina N, Sievert V, Büsow K, Simon B. Establishing the yeast *Saccharomyces cerevisiae* as a system for expression of human proteins on a proteome-scale. *J Struct Funct Genomics.* 2003; 4(2-3):97-108. PMID: 14649293
- [17] Kazachenko KY, Miropolskaya NA, Gening LV, Tarantul VZ, Makarova AV. Alternative splicing at exon 2 results in the loss of the catalytic activity of mouse DNA polymerase ι in vitro. *DNA Repair.* 2017; 50:77-82. <https://doi.org/10.1016/j.dnarep.2017.01.001> PMID: 28077248
- [18] Makarova AV, Stodola JL, Burgers PM. A four-subunit DNA polymerase ζ complex containing Pol δ accessory subunits is essential for PCNA-mediated mutagenesis. *Nucleic Acids Res.* 2012; 40 (22):11618-11626. <https://doi.org/10.1093/nar/gks948> PMID: 23066099
- [19] Makarova AV, Nick McElhinny SA, Watts BE, Kunkel TA, Burgers PM. Ribonucleotide incorporation by yeast DNA polymerase ζ . *DNA Repair.* 2014; 18:63-67. <https://doi.org/10.1016/j.dnarep.2014.02.017> PMID: 24674899
- [20] Lee YS, Gregory MT, Yang W. Human Pol ζ purified with accessory subunits is active in translesion DNA synthesis and complements Pol η in cisplatin bypass. *Proc Natl Acad Sci U S A.* 2014; 111 (8):2954-2959. <https://doi.org/10.1073/pnas.1324001111> PMID: 24449906
- [21] Guillian TA, Jozwiakowski SK, Ehlinger A, Barnes RP, Rudd SG, Bailey L.J. et al. Human PrimPol is a highly error-prone polymerase regulated by single-stranded DNA binding proteins. *Nucleic Acids Res.* 2015; 43(2):1056-1068. <https://doi.org/10.1093/nar/gku1321> PMID: 25550423
- [22] Brown JA, Pack LR, Fowler JD, Suo Z. Pre-steady-state kinetic analysis of the incorporation of anti-HIV nucleotide analogs catalyzed by human X- and Y-family DNA polymerases. *Antimicrob Agents Chemother.* 2011; 55(1):276-283. <https://doi.org/10.1128/AAC.01229-10> PMID: 21078938
- [23] Klenow H, Henningsen I. Effect of monovalent cations on the activity of the DNA polymerase of *Escherichia coli*. *Eur J Biochem.* 1969; 9(1):133-141. PMID: 4891612

- [24] Garg P, Stith CM, Majka J, Burgers PM. Proliferating cell nuclear antigen promotes translesion synthesis by DNA polymerase zeta. *J Biol Chem.* 2005; 280(25):23446–13450. <https://doi.org/10.1074/jbc.C500173200> PMID: 15879599
- [25] Nishimoto N, Suzuki M, Izuta S. Effect of pH on the Misincorporation Rate of DNA Polymerase η . *Biol Pharm Bull.* 2016; 39(6):953–958. <https://doi.org/10.1248/bpb.b15-00900> PMID: 27251497
- [26] Shimazaki N, Yoshida K, Kobayashi T, Toji S, Tamai K, Koiwai O. Over-expression of human DNA polymerase lambda in *E. coli* and characterization of the recombinant enzyme. *Genes Cells.* 2002; 7(7) 639–651. PMID: 12081642
- [27] Copeland WC, Wang TS. Mutational analysis of the human DNA polymerase alpha. The most conserved region in alpha-like DNA polymerases is involved in metal-specific catalysis. *J Biol Chem.* 1993; 268(15)11028–11040. PMID: 8496164
- [28] Baneyx F, Mujacic M. Recombinant protein folding and misfolding in *Escherichia coli*. *Nat Biotechnol.* 2004; 22(11)1399–1408. <https://doi.org/10.1038/nbt1029> PMID: 15529165
- [29] Schlieker C, Bukau B, Mogk A. Prevention and reversion of protein aggregation by molecular chaperones in the *E. coli* cytosol: implications for their applicability in biotechnology. *J Biotechnol.* 2002; 96(1) 13–21. PMID: 12142139
- [30] Schumann W, Ferreira LCS. Production of recombinant proteins in *Escherichia coli*. *Genet Mol Biol.* 2004; 27(3):442–453.
- [31] Porowińska D, Czarnecka J, Komoszyński M. Chaperones are necessary for the expression of catalytically active potato apyrases in prokaryotic cells. *Appl Biochem Biotechnol.* 2014; 173(6): 1349–1359. <https://doi.org/10.1007/s12010-014-0858-6> PMID: 24801402
- [32] Johnson RE, Prakash L, Prakash S. Yeast and human translesion DNA synthesis polymerases: expression, purification, and biochemical characterization. *Methods Enzymol.* 2006; 408:390–407. [https://doi.org/10.1016/S0076-6879\(06\)08024-4](https://doi.org/10.1016/S0076-6879(06)08024-4) PMID: 16793382
- [33] Zhang Y, Yuan F, Wu X, Wang Z. Preferential incorporation of G opposite template T by the low-fidelity human DNA polymerase iota. *Mol Cell Biol.* 2000; 20(19):7099–7108. PMID: 10982826
- [34] Zhang Y, Yuan F, Xin H, Wu X, Rajpal DK, Yang D, et al. Human DNA polymerase kappa synthesizes DNA with extraordinarily low fidelity. *Nucleic Acids Res.* 2000; 28(21):4147–4156. PMID: 11058111
- [35] Rechkoblit O, Gupta YK, Malik R, Rajashankar KR, Johnson RE, Prakash L, et al. Structure and mechanism of human PrimPol, a DNA polymerase with primase activity. *Sci Adv.* 2016; 2(10)e1601317. <https://doi.org/10.1126/sciadv.1601317> PMID: 27819052