

Rare meteorites common in the Ordovician period

Heck P., Schmitz B., Bottke W., Rout S., Kita N., Cronholm A., Defouilloy C., Dronov A., Terfelt F.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2017 Macmillan Publishers Limited, part of Springer Nature. All rights reserved. Most meteorites that fall today are H and L type ordinary chondrites, yet the main belt asteroids best positioned to deliver meteorites are LL chondrites 1,2. This suggests that the current meteorite flux is dominated by fragments from recent asteroid breakup events 3,4 and therefore is not representative over longer (100-Myr) timescales. Here we present the first reconstruction of the composition of the background meteorite flux to Earth on such timescales. From limestone that formed about one million years before the breakup of the L-chondrite parent body 466 Myr ago, we have recovered relict minerals from coarse micrometeorites. By elemental and oxygen-isotopic analyses, we show that before 466 Myr ago, achondrites from different asteroidal sources had similar or higher abundances than ordinary chondrites. The primitive achondrites, such as Iodranites and acapulcoites, together with related ungrouped achondrites, made up ~15-34% of the flux compared with only ~0.45% today. Another group of abundant achondrites may be linked to a 500-km cratering event on (4) Vesta that filled the inner main belt with basaltic fragments a billion years ago 5. Our data show that the meteorite flux has varied over geological time as asteroid disruptions create new fragment populations that then slowly fade away from collisional and dynamical evolution. The current flux favours disruption events that are larger, younger and/or highly efficient at delivering material to Earth.

<http://dx.doi.org/10.1038/s41550-016-0035>

References

- [1] Vernazza, P., et al. Compositional differences between meteorites and near-Earth asteroids. *Nature* 454, 858-860 (2008).
- [2] Dunn, T. L., Burbine, T. H., Bottke, W. F. Jr & Clark, J. P. Mineralogies and source regions of near-Earth asteroids. *Icarus* 222, 273-282 (2014).
- [3] Nesvorný, D., Vokrouhlický, D., Morbidelli, A., Bottke, W. F. Asteroidal source of L chondrite meteorites. *Icarus* 200, 698-701 (2009).
- [4] Bottke, W. F., et al. in *Asteroids IV* (eds Michel, P., et al.) 701-724 (Univ. Arizona Press, 2015).
- [5] Marchi, S., et al. High-velocity collisions from the lunar cataclysm recorded in asteroidal meteorites. *Nat. Geosci.* 6, 303-307 (2013).
- [6] Schmitz, B., Häggström, T., Tassinari, M. Sediment-dispersed extraterrestrial chromite traces a major asteroid disruption event. *Science* 300, 961-964 (2003).
- [7] Heck, P. R., et al. A search for H-chondritic chromite grains in sediments that formed immediately after the breakup of the L-chondrite parent body 470 Ma ago. *Geochim. Cosmochim. Acta* 177, 120-129 (2016).
- [8] Heck, P. R., et al. A single asteroidal source for extraterrestrial Ordovician chromite grains from Sweden and China: High-precision oxygen three-isotope SIMS analysis. *Geochim. Cosmochim. Acta* 74, 497-509 (2010).
- [9] Heck, P. R., Schmitz, B., Baur, H., Halliday, A. N., Wider, R. Fast delivery of meteorites to Earth after a major asteroid collision. *Nature* 430, 323-325 (2004).

- [10] Schmitz, B. Extraterrestrial spinels and the astronomical perspective on Earth's geological record and evolution of life. *Chem. Erde* 73, 117-145 (2013).
- [11] Heck, P. R., Schmitz, B., Baur, H., Wider, R. Noble gases in fossil micrometeorites and meteorites from 470 Myr old sediments from southern Sweden, and new evidence for the L-chondrite parent body breakup event. *Meteorit. Planel Sci.* 43, 517-528 (2008).
- [12] Alwmark, C., Schmitz, B., Meier, M. M. M., Baur, H., Wider, R. A global rain of micrometeorites following breakup of the L-chondrite parent body: Evidence from solar wind-implanted Ne in fossil extraterrestrial chromite grains from China. *Meteorit. Planel Sci.* 47, 1297-1304 (2012).
- [13] Meier, M. M. M., Schmitz, B., Lindskog, A., Maden, C., Wider, R. Cosmic-ray exposure ages of fossil micrometeorites from mid-Ordovician sediments at Lynna River, Russia. *Geochim. Cosmochim. Acta* 125, 338-350 (2014).
- [14] Van Ginneken, M., Folco, L., Cordier, C., Rochette, P. Chondritic micrometeorites from the Transantarctic Mountains. *Meteorit. Planel Sci.* 47, 228-247 (2012).
- [15] Prasad, M. S., Rudraswami, N. G., Dc Araujo, A., Babu, E. V. S. S. K., Kumar, T. V. Ordinary chondritic micrometeorites from the Indian Ocean. *Meteorit. Planel Sci.* 50, 1013-1031 (2015).
- [16] Desnoyers, C., et al. Mineralogy of the Bocaiuva iron meteorite: A preliminary study. *Meteoritics* 20, 113-124 (1985).
- [17] Clayton, R. N., Mayeda, T. K. Oxygen isotope studies of achondrites. *Geochim. Cosmochim. Acta* 60, 1999-2017 (1996).
- [18] Schmitz, B., et al. A new type of solar-system material recovered from Ordovician marine limestone. *NuL Commun.* 7, 11851 (2016).
- [19] Mittlefehldt, D. W. Asteroid (4) Vesta: I. The howardite eucrite diogenite (HED) clan of meteorites. *Chem. Erde* 75, 155-183 (2015).
- [20] Bogard, D. D. K-Ar ages of meteorites: Clues to parent-body thermal histories. *Chem. Erde* 71, 207-226 (2011).
- [21] Swindle, T. D., Kring, D. A., Weirich, J. R. in *Advances in 4Ar/3Ar Dating: From Archaeology to Planetary Sciences* Vol. 378 (eds Jourdan, E, Mark, D. F., Verati, C.) 333-347 (2014).
- [22] Jourdan, F., Andreoli, M. A. G., McDonald, I., Maier, W D. ArP9Ar thermochronology of the fossil LL6-chondrite from the Morokweng crater, South Africa. *Geochim. Cosmochim. Acta* 74, 1734-1747 (2010).
- [23] Dykhuis, M. J., Molnar, L., Van Kooten, S. J., Greenberg, R. Defining the Flora family: Orbital properties, reflectance properties and age. *Icarus* 243, 111-128 (2014).
- [24] Cordier, C., Folco, L. Oxygen isotopes in cosmic spherules and the composition of the near Earth interplanetary dust complex. *Geochim. Cosmochim. Acta* 146, 18-26 (2014).
- [25] Suavet, C., et al. Ordinary chondrite-related giant (800 tm) cosmic spherules from the Transantarctic Mountains, Antarctica. *Geochim. Cosmochim. Acta* 75, 6200-6210 (2011).
- [26] Cordier, C., et al. HED-like cosmic spherules from the Transantarctic Mountains, Antarctica: Major and trace element abundances and oxygen isotopic compositions. *Geochim. Cosmochim. Acta* 77, 515-529 (2012).
- [27] Lindskog, A., Schmitz, B., Cronholm, A., Dronov, A. A Russian record of a Middle Ordovician meteorite shower: Extraterrestrial chromite at Lynna River, St. Petersburg region. *Meteorit. Planel Sci.* 47, 1274-1290 (2012).
- [28] Kita, N. T., Ushikubo, T., Fu, B., Valley, J. W High precision SIMS oxygen isotope analysis and the effect of sample topography. *Chem. Geol.* 264, 43-57 (2009).
- [29] Schmitz, B., Haggstrom, T. Extraterrestrial chromite in Middle Ordovician marine limestone at Kinnekulle, southern Sweden-traces of a major asteroid breakup event. *Meteorit. Planel Sci.* 41, 455-466 (2006).
- [30] Bottke, W E, Nesvorný, D., Grimm, E. R., Morbidelli, A., O'Brien D. P. Iron meteorites as remnants of planetesimals formed in the terrestrial planet region. *Nature* 439, 821-824 (2006).
- [31] Burkhardt C., et al. NWA 5363!NWA 5400 and the Earth: Isotopic twins or just distant cousins? *Planel Inst.* 46, 2732 (2015).