

Bulletin of the Academy of Sciences of the USSR Division of Chemical Science 1972 vol.21 N8,
pages 1649-1651

Investigation of conformational equilibrium in a series of some 1,3-dioxa-2-phosphorinanes

Raevskii O., Vereshchagin A., Khalitov F., Donskaya Y., Cherkasov R., Ovchinnikov V.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

1. The IR spectra of certain 1,3-dioxa-2-phosphorinanes were studied under conditions of various polarities of the medium and temperatures, and it was shown that 2,4-dimethyl-2-thiono- and 2-chloro-2-thiono-1,3-dioxa-2-phosphorinanes are characterized by stabilization of one conformational form, while for 2-methyl-2-thiono-1,3-dioxa-2-phosphorinane a dynamic equilibrium of two conformers is realized with an appreciable dependence of their amounts on the dielectric permeability of the medium. 2. An equilibrium of three conformers was detected for 2-chloro-4-methyl-2-thiono-1,3-dioxa-2-phosphorinane. It was hypothesized that, together with the "chair" conformation of the ring with axial and equatorial positions of the P=S bond, the conformer with an equatorial arrangement of the P=S bond and a "boat" form of the ring participates in the equilibrium. © 1973 Consultants Bureau.

<http://dx.doi.org/10.1007/BF00851171>
