

Bulletin of the Academy of Sciences of the USSR Division of Chemical Science 1987 vol.36 N2,  
pages 381-384

---

## Synthesis and vibrational spectra of some 2-alkoxy-5-6-benzo-1,3,2-dioxastibepines

Arbuzov B., Shagidullin R., Vinogradova V., Mareev Y., Shakirov I., Fedotova N.  
*Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia*

---

### Abstract

1) Tranesterification of the respective antimony trialkoxides with phthalyl alcohol have 2-methoxy-, 2-ethoxy-, and 2-tert-butoxy-5,6-benzo-1,3,2-dioxastibepines. 2) According to the IR and Raman spectra, 2-ethoxy- and 2-tert-butoxy-5,6-benzo-1,3,2-dioxastibepines are polymerized in the crystalline phase; the  $\nu_{\text{Sb-O}}$  frequencies are at least 100  $\text{cm}^{-1}$  higher in melt and in solution than in the crystalline phase; they exist both as dimer and monomer. 2-tert-butoxy-5,6-benzo-1,3,2-dioxastibepine exists predominantly in the chair conformation. © 1987 Plenum Publishing Corporation.

<http://dx.doi.org/10.1007/BF00959387>

---