

Chemistry of Heterocyclic Compounds 2002 vol.38 N4, pages 406-416

Condensation of 2-pyridylmethylithium nucleophiles and pyridine electrophiles as a convenient synthetic route to polydentate chelating N-donor ligands

Vedernikov N., Miftakhov R., Borisoglebski S., Caulton K., Solomonov B.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

Condensation of 2-pyridylmethylithium or (6-methyl-2-pyridyl)methylithium nucleophiles and pyridine, 2-picoline, or 4-tert-butylpyridine as electrophiles leads to new polydentate N-donor ligands, methyl-, tert-butyl-substituted tripyridinedimethanes, or to tripyridinedimethane itself, in good or high yields. Depending on the reagent ratio, solvent used, and reaction conditions, the corresponding intermediate dipyridinemethanes can be minor by-product or major products of the condensation. In contrast to 2-pyridylmethylithium, lithiated 2-isopropylpyridine does not react with pyridine electrophiles. Vice versa, nucleophilic substitution at the C(2)-pyridine carbon of 2,2-bis(2-pyridyl)propane with 2-pyridylmethylithium takes place to produce products of condensation of 2-isopropylpyridine and dipyridylmethylithium. DFT calculations of the Gibbs free energies of reactions combined with pKa values of the CH-acids involved help to explain the reactivity observed.

<http://dx.doi.org/10.1023/A:1016023103898>

Keywords

(6-methyl-2-pyridyl) methylithium, 2,6-lutidine, 2-picoline, 2-pyridylmethylithium, 4-tert-butylpyridine, C-C bond cleavage, Condensation, DFT calculations, Dipyridinemethanes, Gibbs free energy of reaction, Pyridine, tripyridinedimethanes