

An accurate simplified data treatment for the initial adsorption kinetics in conditions of laminar convection in a slit: Application to protein adsorption

Déjardin P., Vasina E.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

We present the derivation of a simple approximation for the original expression of the adsorption rate [Langmuir 10 (1994) 3898] in conditions of laminar flow in a slit, to relate the measured initial kinetic constant k with the interfacial kinetic constant k_a and the transport-limited Lévêque constant k_{Lev} . The same method of derivation is applied here to get a simple approximation of the average kinetic constant \bar{k} [Biomaterials 20 (1999) 1621]. For the local value, at distance x from the entrance of the slit, we propose $k(x)/k_a = (u-1)(au-1)/(bu+1)$, where $u = k(x)/k_{Lev}$, $a = 0.452$, $b = -0.625$, with a maximal error of 1% in comparison with the exact solution. For the average value over the length of the slit, we propose $\bar{k}/k_a = (U-1)(A-1)/(BU+1)$, where $U = \bar{k}/k_{Lev}$, $A = 0.203$, $B = -0.273$, with a maximal error of 0.03%. These approximations lead to an easy determination of the adsorption constant and diffusion coefficient D of the solute, as appropriate plots of experimental data provide k_a and $D^{2/3}$ as the intercepts of the curve with the ordinate and abscissa axes, respectively. It is pointed out that the linear approximation $k^{-1} = k_a^{-1} + k_{Lev}^{-1}$ would lead to the overestimation of both the diffusion coefficient and adsorption kinetic constant. As an example, the application to the analysis of experimental data for adsorption of α -chymotrypsin onto mica plates is provided. © 2003 Elsevier B.V. All rights reserved.

<http://dx.doi.org/10.1016/j.colsurfb.2003.09.005>

Keywords

Convection-diffusion, Electrochemistry, Interfacial reaction, Protein adsorption