Heteroatom Chemistry 2004 vol.15 N3, pages 225-232

## Disilyl dithiophosphonates in the synthesis of open chain and cyclic organothiophosphorus compounds

Nizamov I., Sergeenko G., Popovich Y., Khaibullin R., Al'metkina L., Abalonin B., Batyeva E., Krivolapov D., Litvinov I.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

## **Abstract**

O-(Trimethylsiloxy)alkyl S-trimethylsilyl aryldithiophosphonates 7a-d were obtained by the reaction of 2,4-diaryl-1,3,2,4-dithiadiphosphetane-2,4- disulfides 5a,b with disilyl derivatives of glycols 6a,c and salicyl alcohol 6b. The reactions of mixed O,S-bis(trimethylsilyl) 2,4-di(3,5-i-tert-butyl-4- hydroxyphenyl)dithiophosphonate 1 and S-silyl aryldithiophosphonates 7a,b with S,S-diethyldithiodiphenylgermane 2, dichlorodiphenylgermane 8a, and dichlorodiphenylstannane 8b were studied. The structure of hexaphenyl-2,4,6,1,3, 5-trithiatrigerminane 11 was established by X-ray single crystal diffraction. © 2004 Wiley Periodicals, Inc.

http://dx.doi.org/10.1002/hc.20009