

Molecular Crystals and Liquid Crystals Science and Technology Section A: Molecular Crystals and Liquid Crystals 1999 vol.332, pages 109-118

Liquid Crystalline Complexes of Cu(II) and Pd(II) with Ferrocene-Containing Ligands

Kadkin O., Galyametdinov Y., Rakhmatullin A.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

The new liquid crystalline heteronuclear complexes Cu(II) 1a and Pd(II) 1b with a non-mesogenic ferrocene-containing β -enaminoketone L1 of the formula $C_5H_5FeC_5H_4-C_6H_4-NH-C_2H_2(-O)-C_6H_4OC_{12}H_{25}$, showing monotropic nematic and smectic A phases, are studied. The ortho-palladated Cl-bridged dimeric complex 3 with a non-mesogenic ferrocene-containing Schiff's base L2 of the formula $C_5H_5FeC_5H_4-C_6H_4-N=CH-C_6H_4OC_{10}H_{21}$, forming a stable smectic A phase, has been synthesized. The novel mixed-ligand heteronuclear complexes 4-7 have been obtained by treatment of the ortho-palladated complex 3 with appropriate ligands. Some of the resulting products exhibit the rather low-melting (below 100deg;C) smectic A phases. Structures of the compounds are studied by elemental analysis, 1H and ^{13}C NMR, and ESR spectroscopies.

Keywords

Cyclopalladated complexes, Ferrocene derivatives, Heteronuclear complexes, Metallomesogens