

Physical Review E - Statistical, Nonlinear, and Soft Matter Physics 2002 vol.66 N3, pages 031508/1-031508/10

Low-temperature phase separation of a binary liquid mixture in porous materials studied by cryoporometry and pulsed-field-gradient NMR

Valiullin R., Furó I.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

The low-temperature liquid-liquid phase separation of the partially miscible hexane-nitrobenzene mixture imbibed in porous glasses of different pore sizes from 7 to 130 nm has been studied using ^1H NMR (nuclear magnetic resonance) cryoporometry and pulse field gradient NMR methods. The mixture was quenched below both its upper critical solution temperature (T_{cr}) and the freezing point of nitrobenzene. The size distribution of frozen nitrobenzene domains was derived through their melting point suppression according to the Gibbs-Thompson relation. The obtained data reveal small initial droplets of nitrobenzene surrounded by hexane, which are created as the temperature is decreased below T_{cr} and which thereafter coalesce by a droplet-diffusion mechanism. The inter-relation between the pore size and the found size distribution and shapes of nitrobenzene domains is discussed, as well as several aspects of molecular self-diffusion. © 2002 The American Physical Society.

<http://dx.doi.org/10.1103/PhysRevE.66.031508>
