



Journal of Molecular Structure (Theochem) 343 (1995) 195–198

**THEO
CHEM**

The structures of dimeric stereoisomeric tartrates of iron(III) as determined by molecular mechanics calculations

V.V. Chevela*, S.N. Matveev, V.E. Semenov, S.G. Bezryadin, T.V. Savitskaya,
I.R. Gromova, G.A. Shamov

V.I. Ul'yanov-Lenin Kazan State University, Lenin str., 18, Kazan 420008, Russia

Received 15 December 1994

Abstract

The stereospecific formation of iron(III) complexes with *d*- and *dl*-tartaric acids (*dH*₄L, *dlH*₄L respectively) is discussed. Molecular mechanics calculations (MIND program, model of Dashevsky–Plyamovaty) are applied to modelling of dimeric tartrate Fe₂(*d*-L)₂²⁻ and Fe₂(*d*-L)(*l*-L)₂²⁻ (II) structures. Stereospecific formation is explained by lesser values of energy of rigid iron(III) octahedral coordination in (I).

1. Introduction

The formation of iron(III) tartrates has been reported in a considerable number of works [1–4]. At pH 2–5, complex forms of iron(III) tartrates undergo interconversion in the conventional manner. A dimeric form Fe₂L₂²⁻ (where L⁴⁻ is the rest of the tartaric acid H₄L, C₄H₆O₆) occurs in aqueous solutions in this pH region. The same scheme was used by us in our investigations of stereoselective and stereospecific effects of formation of iron(III) with stereoisomeric *d*- and *dl*-tartaric acids.

The *dl*-dimeric tartrate Fe₂(*d*-L)(*l*-L)₂²⁻ was demonstrated by methods of proton magnetic relaxation and pH measurements with mathematic simulations as in Ref. [3]. From stability data it is shown that *d*-dimeric tartrate is not formed, in spite of the *dl*-dimeric tartrate. To explain this fact we used molecular mechanics calculations in the

program MIND whose results models are described in Ref. [5].

2. Experiments and results

The molecular mechanics study was accomplished using the program MIND. In the present work we used the molecular mechanics method which takes into consideration interactions between heteroatom electron lone pairs [6]. This method, called the Dashevsky–Plyamovaty model [5,7], and in the force field of that model, the steric energy *E* of the system is presented in the expression in Eq. (1).

$$E = \frac{1}{2} \sum_i K_l (l_i - l_0)^2 + \frac{1}{2} \sum_i K_{\alpha_i} (\alpha_i - \alpha_0)^2 + \frac{1}{2} \sum_i V_i [1 + x \cos(n_i \varphi_i)] + \sum_{i>j} [-ar_{ij}^{-b} + b \exp(-cr_{ij})] \quad (1)$$

* Corresponding author.