

Polymer 2004 vol.45 N1, pages 165-170

---

## Self-diffusion of low-generation PAMAM dendrimers with hydroxyl surface groups in solutions: A general regularity

Sagidullin A., Fritzing B., Scheler U., Skirda V.  
*Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia*

---

### Abstract

An experimental study of the self-diffusion and nuclear magnetic relaxation of poly(amidoamine) dendrimers with hydroxyl surface groups (PAMAM-OH) dissolved in methanol over a wide range of concentration ( $\psi$ ) is reported. It is shown that experimental concentration dependences of PAMAM-OH self-diffusion coefficients ( $D$ ) can be reduced to the so-called generalized  $\psi$  dependence. Over macromolecular concentration range studied, the generalized concentration dependence of PAMAM-OH  $D$  coincides with analogous curve obtained for poly(allylcarbosilane) dendrimers of high generations. This result confirms the existence of common regularities of the dendritic macromolecule self-diffusion, and their independence of the individual physicochemical and structural properties of dendrimer, solvent, and features of their interactions in the given systems. The concentration dependence of the PAMAM-OH diffusion also exhibits a clear signature of an inflexible molecule. © 2003 Elsevier Ltd. All rights reserved.

<http://dx.doi.org/10.1016/j.polymer.2003.10.087>

---

### Keywords

Dendrimer, Scaling, Self-diffusion