

Astronomy and Astrophysics 2008 vol.491 N1, pages 267-277

---

## Modeling of non-stationary accretion disks in X-ray novae A 0620-00 and GRS 1124-68 during outburst

Suleimanov F., Lipunova V., Shakura I.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

---

### Abstract

**Aims.** We address the task of modeling soft X-ray and optical light curves of X-ray novae in the high/soft state. **Methods.** The analytic model of viscous evolution of an externally truncated accretion  $\alpha$ -disk is used. Relativistic effects near a Kerr black hole and self-irradiation of an accretion disk are taken into account. **Results.** The model is applied to the outbursts of X-ray nova Monocerotis 1975 (A 0620-00) and X-ray nova Muscae 1991 (GRS 1124-68). Comparison of observational data with the model yields constraints on the angular momentum (the Kerr parameter) of the black holes in A 0620-00 and GRS 1124-68: 0,3-0,6 and  $\leq .$ , and on the viscosity parameter  $\alpha$  of the disks: 0.7-0.95 and 0.55-0.75. We also conclude that the accretion disks should have an effective geometrical thickness 1.5-2 times greater than the theoretical value of the distance between the photometric layers. © 2008 ESO.

<http://dx.doi.org/10.1051/0004-6361:200810155>

---

### Keywords

Binaries: close, Stars: individual: Nova Mon 1975, Stars: individual: Nova Mus 1991, X-rays: binaries