

Available online at www.sciencedirect.com

Journal of Molecular Structure 828 (2007) 147–153

Journal of
**MOLECULAR
STRUCTURE**www.elsevier.com/locate/molstruc

Conformational isomerism in 3,5,8-trioxabicyclo[5.1.0]octane and its diastereomeric 4-methyl derivatives. A combined IR, X-ray and ab initio study

A.E. Klimovitskii^a, E.E. Zvereva^b, G.A. Chmutova^a, A.B. Dobrynin^b, V.Yu. Fedorenko^a, Yu.G. Shtyrlin^a, I.A. Litvinov^b, T.V. Bulygina^a, E.N. Klimovitskii^{a,*}

^a A.M. Butlerov Chemical Institute, Kazan State University, Kazan 420111, Box 115, Russian Federation

^b A.E. Arbusov Institute of Organic and Physical Chemistry, Russian Academy of Sciences, Arbuzov Str. 8, Kazan 420088, Russian Federation

Received 27 March 2006; received in revised form 25 May 2006; accepted 30 May 2006

Available online 4 August 2006

Abstract


IR spectroscopy and ab initio calculations have been applied to the title compounds. Bicyclic acetals were found to exist as mixture of chair and twist-boat conformations, a parent epoxide being in ternary equilibrium. X-ray data on 4-methyl (exo) derivative display a twist-boat form.

© 2006 Elsevier B.V. All rights reserved.

Keywords: Seven-membered bicyclic acetals; IR spectroscopy; X-ray crystallography; ab initio calculations

1. Introduction

Investigations of 1,3-dioxanes with substituent at C⁵ (polar especially) show a number of interesting trends [1]. 3,5-Dioxabicyclo[5.1.0]octanes with exo- and endo chair-like conformations relate closely to above 1,3-dioxanes if one assumes that three-membered cycles in these molecules are equivalent to substituents at C⁵ in the stereochemical sense (axial ≡ endo and equatorial ≡ exo). The noticeable feature of bicyclic acetals is the real propensity to exist in both chair conformations and a twist-boat form in addition [2–9]. It is worth noting that seven-membered acetals with a planar fragment have been inspected thoroughly and untraditional chair–twist-boat (C ⇌ TB) equilibrium has been clarified [10].



One can expect that bicyclic acetals with X = C, O, S, N may serve as suitable models for the investigation of polar and steric interactions between acetal moiety and three-membered rings. Encouraged by available stereochemical information implying both a broad range of

* Corresponding author. Fax: +7 8432 9272 78.

E-mail address: evgenii.klimovitskii@ksu.ru (E.N. Klimovitskii).