

Molecular Biology 2013 vol.47 N2, pages 293-298

C-terminus of transcription factor TnrA from *Bacillus subtilis* controls DNA-binding domain activity but is not required for dimerization

Fedorova K., Scharafutdinov I., Turbina E., Bogachev M., Ilinskaja O., Kayumov A.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

The transcription factor TnrA, which belongs to the MerR transcription regulators, in *Bacillus subtilis* controls genes of nitrogen metabolism during nitrogen limiting conditions. As all the DNA-binding proteins, it is active as a dimer in cells, but the dimerization site is still unknown. The multiple alignment of TnrA homologs from other Bacilli allowed to identify the putative dimerization sites. Using the C-terminal truncated TnrA proteins it is established, that, in contrast to other MerR-proteins, the TnrA C-terminus does not participate in dimerization. Surface plasmon resonance has revealed that C-terminus truncations of TnrA do not inactivate its DNA-binding activity. Contrary, increased the affinity towards DNA, confirming that C-terminus controls the DNA-binding activity in a full-length TnrA. © 2013 Pleiades Publishing, Ltd.

<http://dx.doi.org/10.1134/S0026893313020052>

Keywords

Bacillus subtilis, dimerization, transcription factor TnrA