ADIOACTIVITY

Journal of Environmental Radioactivity 107 (2012) 78-85



Contents lists available at SciVerse ScienceDirect

Journal of Environmental Radioactivity

journal homepage: www.elsevier.com/locate/jenvrad

Review

Fallout radionuclide-based techniques for assessing the impact of soil conservation measures on erosion control and soil quality: an overview of the main lessons learnt under an FAO/IAEA Coordinated Research Project

G. Dercon^{a,*}, L. Mabit^a, G. Hancock^b, M.L. Nguyen^a, P. Dornhofer^a, O.O.S. Bacchi^c, M. Benmansour^d, C. Bernard^{a,1}, W. Froehlich^e, V.N. Golosov^f, S. Haciyakupoglu^g, P.S. Hai^h, A. Klikⁱ, Y. Li^j, D.A. Lobb^k, Y. Onda¹, N. Popa^m, M. Rafiqⁿ, J.C. Ritchie^o, P. Schuller^p, A. Shakhashiro^q, P. Wallbrink^b, D.E. Walling^r, F. Zapata^a, X. Zhang^s

^a Joint FAO/IAEA Division of Nuclear Techniques in Food and Agriculture, International Atomic Energy Agency, IAEA, Department of Nuclear Sciences and Applications, Wagramerstrasse 5, A-1400 Vienna, Austria

^b Division of Land and Water, Commonwealth Scientific and Industrial Research Organisation (CSIRO), Canberra, Australia

^c Center for Nuclear Energy in Agriculture, Universidade de Sao Paulo, Piracicaba, Brazil

^d Centre National de l'Energie, des Sciences et des Techniques Nucléaires, Rabat, Morocco

e Polish Academy of Sciences, Institute of Geography and Spatial Organization, Homerka Laboratory of Fluvial Processes, Nawojowa, Malopolska, Poland

^fLaboratory for Soil Erosion and Fluvial Processes, Faculty of Geography, Moscow State University, Moscow, and Kazan Federal University, Kazan, Russia

^g Istanbul Technical University, Energy Institute, Istanbul, Turkey

^h Center for Environment Research & Monitoring, Dalat Nuclear Research Institute, Dalat, Viet Nam

¹Institute of Hydraulics and Rural Water Management, University of Natural Resources and Applied Life Sciences, Vienna, Austria

^j Chinese Academy of Agricultural Sciences, Institute of Environment and Sustainable Development in Agriculture, Beijing, China

^k University of Manitoba, Winnipeg, Canada

¹Laboratory of Integrative Environmental Sciences, School of Life and Environmental Sciences, University of Tsukuba, Tsukuba, Japan

^m Research and Development Center for Soil Erosion Control, Perieni, Romania

ⁿ Pakistan Institute of Nuclear Science and Technology, Islamabad, Pakistan

^o US Department of Agriculture, Agricultural Research Service, Hydrology and Remote Sensing Laboratory, Beltsville, USA

^p Universidad Austral de Chile, Facultad de Ciencias, Instituto de Ciencas Químicas, Valdivia, Chile

^q International Atomic Energy Agency, Chemistry Unit, Terrestrial Environment Laboratory, IAEA Laboratories Seibersdorf, Austria

^r University of Exeter, School of Geography, Exeter, Devon, United Kingdom

^s Chinese Academy of Sciences, Institute of Mountain Hazards and Environment, Chengdu, Sichuan, China

ARTICLE INFO

Article history: Received 3 September 2011 Received in revised form 9 January 2012 Accepted 16 January 2012 Available online 14 February 2012

Keywords: Caesium-137 Lead-210 Beryllium-7 FRN Land degradation Agricultural landscapes

ABSTRACT

This paper summarizes key findings and identifies the main lessons learnt from a 5-year (2002–2008) coordinated research project (CRP) on "Assessing the effectiveness of soil conservation measures for sustainable watershed management and crop production using fallout radionuclides" (D1.50.08), organized and funded by the International Atomic Energy Agency through the Joint FAO/IAEA Division of Nuclear Techniques in Food and Agriculture. The project brought together nineteen participants, from Australia, Austria, Brazil, Canada, Chile, China, Japan, Morocco, Pakistan, Poland, Romania, Russian Federation, Turkey, United Kingdom, United States of America and Vietnam, involved in the use of nuclear techniques *and, more particularly*, fallout radionuclides (FRN) to assess the relative impacts of different soil conservation measures on soil erosion and land productivity. The overall objective of the CRP was to develop improved land use and management strategies for sustainable watershed management through effective soil erosion control practices, by the use of ¹³⁷Cs (half-life of 30.2 years), ²¹⁰Pb_{ex} (half-life of 22.3 years) and ⁷Be (half-life of 53.4 days) for measuring soil erosion over several spatial and temporal scales.

The environmental conditions under which the different research teams applied the tools based on the use of fallout radionuclides varied considerably -a variety of climates, soils, topographies and land uses. Nevertheless, the achievements of the CRP, as reflected in this overview paper, demonstrate that fallout

* Corresponding author.

- E-mail address: g.dercon@iaea.org (G. Dercon).
- ¹ Present address: Ministère de l'Agriculture, des Pêcheries et de l'Alimentation

du Québec, Québec, Canada.

⁰²⁶⁵⁻⁹³¹X/\$ – see front matter \odot 2012 Elsevier Ltd. All rights reserved. doi:10.1016/j.jenvrad.2012.01.008