

Human Ecology 2015 vol.2015 N1, pages 28-35

Efficiency of new pyrimidine derivativs influence on physical working capacity of rats in the test 'swimming to failure'

Zobov V., Nazarov N., Vyshtakalyuk A., Galyametdinova I., Semenov V., Reznik V.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© Northern State Medical University. There have been studied effects of xymedon and six new pyrimidine derivatives, that are close and distant analogs of xymedon, on rats' working capacity in the test "swimming to failure". It has been shown that a single administration of the studied compounds did not have a statistically significant effect in the test. In the conditions of multiple intraperitoneal administration of the studied pyrimidine derivatives, the compound L-ascorbate, 1-(2-hydroxyethyl)-4.6-dimethyl-1.2-dihydropyrimidine-2-one had the lowest toxicity and the most pronounced actoprotective effect. Introduction in the dose of 20 mg/kg caused a statistically significant increase 440 % in the duration of swimming of rats on the 14th day of the experiment compared with the control group. Multiple administration of the compound in the conditions of physical load did not affect leucopoiesis, but stimulates erythropoiesis resulting in an increase in the number of erythrocytes and a hemoglobin level. The substance introduction under mixed exhausting loads prevented such changes of blood biochemical parameters as reduction of glucose, increased of urea and lactic acid levels, what indicates improvement in the animals' tolerability of loads and an anti-catabolic effect of the compound. Absence of hepato- and cardiotoxic effects of the substance has been shown.

Keywords

Physical working capacity, Pyrimidine derivatives, Stimulation efficiency, Swimming to failure, Xymedon