

Available online at www.sciencedirect.com

ScienceDirect

<http://www.elsevier.com/locate/biombioe>

Effects of the reduction of the hydraulic retention time to 1.5 days at constant organic loading in CSTR, ASBR, and fixed-bed reactors – Performance and methanogenic community composition

Thomas Schmidt ^{a,c,*}, Ayrat M. Ziganshin ^b, Marcell Nikolausz ^d,
Frank Scholwin ^c, Michael Nelles ^{a,c}, Sabine Kleinsteuber ^d, Jürgen Pröter ^a

^a Department of Biochemical Conversion, Deutsches Biomasseforschungszentrum gemeinnützige GmbH, Leipzig 04347, Germany

^b Department of Microbiology, Kazan (Volga Region) Federal University, Kazan 420008, The Republic of Tatarstan, Russia

^c Faculty of Agricultural and Environmental Sciences, University of Rostock, Rostock 18059, Germany

^d Department of Environmental Microbiology, UFZ-Helmholtz Centre for Environmental Research, Leipzig 04318, Germany

ARTICLE INFO

Article history:

Received 22 October 2013

Received in revised form

3 June 2014

Accepted 17 July 2014

Available online 12 August 2014

Keywords:

Biogas

Retention time

Reactors

Methanogenic community

Washout

ABSTRACT

The hydraulic retention time (HRT) is one of the key parameters in biogas processes and often it is postulated that a minimum HRT of 10–25 days is obligatory in continuous stirred tank reactors (CSTR) to prevent a washout of slow growing methanogens. In this study the effects of the reduction of the HRT from 6 to 1.5 days on performance and methanogenic community composition in different systems with and without immobilization operated with simulated thin stillage (STS) at mesophilic conditions and constant organic loading rates (OLR) of 10 g L⁻¹d⁻¹ of volatile solids were investigated. With the reduction of the HRT process instability was first observed in the anaerobic sequencing batch reactor (ASBR) (at HRT of 3 days) followed by the CSTR (at HRT of 2 days). The fixed bed reactor (FBR) was stable until the end of the experiment, but the reduction of the HRT to 1.5 days caused a decrease of the specific biogas production to about 450 L kg⁻¹ of VS compared to about 600 L kg⁻¹ of VS at HRTs of 4–5 days. *Methanoculleus* and *Methanosarcina* were the dominant genera under stable process conditions in the CSTR and the ASBR and members of *Methanosaeta* and *Methanospirillum* were only present at HRT of 4 days and lower. In the effluent of the FBR *Methanosarcina* spp. were not detected and *Methanosaeta* spp. were more abundant than in the other reactors.

© 2014 Elsevier Ltd. All rights reserved.

* Corresponding author. Department Biochemical Conversion, Deutsches Biomasseforschungszentrum gemeinnützige GmbH, Torgauer Straße 116, 04347 Leipzig, Germany. Tel.: +49 341 2434 516.

E-mail address: thomas.schmidt@dbfz.de (T. Schmidt).

<http://dx.doi.org/10.1016/j.biombioe.2014.07.021>

0961-9534/© 2014 Elsevier Ltd. All rights reserved.