


Contents lists available at [SciVerse ScienceDirect](http://SciVerse.ScienceDirect.com)

# NeuroImage

journal homepage: [www.elsevier.com/locate/ynimg](http://www.elsevier.com/locate/ynimg)

## Review

# Spontaneous activity in developing sensory circuits: Implications for resting state fMRI

Matthew Colonnese<sup>a,b,c,\*</sup>, Rustem Khazipov<sup>a,b,d</sup>

<sup>a</sup> INSERM, U29, INMED, Marseille 13009, France

<sup>b</sup> Aix Marseille Université, Faculté des Sciences, Marseille, F-13000, France

<sup>c</sup> Department of Pharmacology and Physiology, Institute for Neuroscience, George Washington University, Washington DC 20037, USA

<sup>d</sup> Kazan Federal University, Kazan, Russia

## ARTICLE INFO

### Article history:

Accepted 13 February 2012

Available online 24 February 2012

### Keywords:

Development

Resting state

Preterm

Neonatal

Spindle burst

Spontaneous activity

## ABSTRACT

The immature brain spontaneously expresses unique patterns of electrical activity that are believed to contribute to the development of neuronal networks. Certain electrographic features of this activity, particularly modulation on an infraslow time scale, resemble activity patterns observed in the mature brain at 'rest', loosely defined as the absence of an investigator imposed task. However, it is not clear whether the immature activity patterns observed at rest are precursors of the spontaneous neuronal activity that forms resting state networks in the adult. Here, we review recent studies that have explored the generative mechanisms of resting state activity during development in the primary sensory systems of premature human neonates and neonatal rodents. The remarkable hypothesis suggested by this work is that while resting state activity during the pre- and possibly near-term period can bear superficial resemblance to adult activity it is fundamentally different in terms of function and origin. During early development spontaneous thalamocortical activity in primary sensory regions is determined largely by transitory generators in the sensory periphery. This is in contrast to the adult, where spontaneous activity generated within thalamocortex, particularly by cortico-cortical connections, dominates. We therefore suggest a conservative interpretation of developmental mapping studies which are based on indirect measurement of activity (e.g. fMRI), or on the partitioning of EEG frequency using bands derived from adult studies. The generative mechanisms for brain activity at early ages are likely different from those of adults, and may play very different roles; for example in circuit formation as opposed to attention.

© 2012 Elsevier Inc. All rights reserved.

## Contents

Resting state activity in neonates and adults . . . . .	2213
Discontinuous temporal organization of early activity . . . . .	2214
Unique activity patterns in preterm infants . . . . .	2214
Appearance . . . . .	2215
Modulation. . . . .	2215
Developmental separation. . . . .	2215
Generative mechanisms. . . . .	2216
Peripheral dominance of resting activity during early development . . . . .	2216
Development of intrinsic thalamocortical activities . . . . .	2217
Early cortical function and implications for resting state fMRI . . . . .	2219
Summary . . . . .	2219
References . . . . .	2220

\* Corresponding author at: Ross Hall suite 640, 2300 Eye Street NW, Washington DC 20037, USA.  
E-mail address: [colonnese@gwu.edu](mailto:colonnese@gwu.edu) (M. Colonnese).