

Russian Journal of General Chemistry 2009 vol.79 N1, pages 24-30

Template synthesis in the Cu(II)-dihydrazinomethanethione-acetone ternary system

Mikhailov O., Kazymova M., Shumilova T., Solov'Eva S.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

Complex formation in the Cu(II)-dihydrazinomethanethione (H₂NHN-CS-NHNH₂)-acetone ternary system in an ethanol solution containing CuCl₂, dihydrazinomethanethione, and acetone, as well as on contact of gelatin-immobilized copper(II) hecacyanoferrate(II) with alkaline (pH >10) aqueous solutions containing the above organic compounds was studied. It was found that template synthesis is realized in both cases but gives different products: in the first case, a heteroligand chelate of CuL₁(OH) with 9-hydrazino-9-mercapto-4,6,6-trimethyl-2,3,7,8-tetraazanona-3,8-dienethiohydrazide and hydroxide ion is formed, while in the second, a chelate of CuL₂ with 2,8,10,10,16-pentamethyl-3,4,6,7,11, 12,14,15-octaazaheptadeca-2,5,7,12,15-pentaene-5,13-dithiol. In both cases, dihydrazinomethanethione and acetone function as ligands. A scheme of involved processes is suggested. © 2009 Pleiades Publishing, Ltd. Reerernces:.

<http://dx.doi.org/10.1134/S1070363209010046>
