

Optical spectroscopy of PrFe₃(BO₃)₄: Crystal-field and anisotropic Pr-Fe exchange interactions

Popova M., Stanislavchuk T., Malkin B., Bezmaternykh L.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

High-resolution polarized optical absorption spectra of PrFe₃(BO₃)₄ in the paramagnetic and antiferromagnetic phases are reported. The measured energies of the crystal-field (CF) levels within the 4f² configuration of Pr³⁺ in the paramagnetic PrFe₃(BO₃)₄ are described by the CF model that involves the 4f²/4f⁵d and 4f²/4f⁶p configuration interactions. Ordering of Fe spins along the crystalline c axis below T_N = 32K is confirmed by the analysis of the spectra of Er³⁺ introduced as a probe into PrFe₃(BO₃)₄. To account for the observed changes in the optical spectra of Pr³⁺ at temperatures below T_N, in particular, for the shift of the CF levels, splitting of the CF doublets, and the appearance of forbidden lines, the Pr-Fe exchange Hamiltonian defined by seven parameters is considered. The theoretical approach has been tested by calculating the temperature dependence of the magnetic susceptibility. A good agreement between theory and optical and magnetic experimental data is found demonstrating the validity of the model used. The obtained results confirm that the model of the iron dimers inside the spiral chains of Fe³⁺(O₂)₆ octahedrons introduced by us earlier for NdFe₃(BO₃)₄ and modified in the present work may serve as a basis for analyzing the low-temperature properties of other rare-earth iron borates. © 2009 The American Physical Society.

<http://dx.doi.org/10.1103/PhysRevB.80.195101>
