

Contents lists available at [SciVerse ScienceDirect](http://SciVerse.Sciencedirect.com)

Journal of Molecular Liquids

journal homepage: www.elsevier.com/locate/molliq

Micellization of alkylated 1,4-diazabicyclo[2.2.2]octane by nuclear magnetic resonance technique using pulsed gradient of static magnetic field

Lucia Ya. Zakharova^{a,b,*}, Nail K. Gaysin^a, Oleg I. Gnezdilov^{c,d}, Farid I. Bashirov^a, Ruslan R. Kashapov^{a,b}, Elena P. Zhiltsova^b, Tatiana N. Pashirova^b, Svetlana S. Lukashenko^b

^a Kazan State Technological University, Marks 68, 420015 Kazan, Russia

^b A.E. Arbuzov Institute of Organic and Physical Chemistry, Kazan' Scientific Center of the Russian Academy of Sciences, Arbuzova 8, 420088 Kazan, Russia

^c Zavoisky Kazan Physical-Technical Institute, Kazan' Scientific Center of the Russian Academy of Sciences, Sibirsky tract 10/7, 420029 Kazan, Russia

^d Kazan (Volga region) Federal University, Kremlyovskaya 18, 420008 Kazan, Russia

ARTICLE INFO

Article history:

Received 11 March 2011

Received in revised form 26 November 2011

Accepted 30 December 2011

Available online 14 January 2012

Keywords:

Alkylated 1,4-diazabicyclo[2.2.2]octane

Micellization

Critical micelle concentration

Aggregation number

Translational diffusion

ABSTRACT

A phenomenon of micellization of 4-aza-1-tetradecyl-1-azoniabicyclo[2.2.2]octane bromide (AB-14) in aqueous solution has been studied by the methods of nuclear magnetic resonance using magnetic field pulse gradient and fluorimetry. The critical micelle concentration is determined; concentration dependences of percentage of free AB-14 molecules and those associated with micelles as well as radii of micelles and numbers of their aggregation are established. Effect of external and internal translational diffusion of molecules on lifetime of AB-14 molecules in micelles is investigated.

© 2012 Elsevier B.V. All rights reserved.

1. Introduction

Studying the nature of interaction of the surfactant (SA) molecules with a dissolvent is one of the main purposes of the micellization theory being reduced to revealing a role of these interaction in both a micellization and an exchange of free and aggregated molecules of SA and an arrival to an equilibrium state between the processes of aggregation and disaggregation of micelles. A method of nuclear magnetic resonance with magnetic field pulse gradient (NMR-MFPG) is largely used in solving these problems. By means of this method, it is possible to determine self-diffusion coefficients (SDC) of the free SA molecules in a solution and those associated in micelles, a critical micelle concentration (CMC), a percentage of free SA molecules and the micelle-forming molecules (MFM), sizes of micelles and a number of their aggregation.

A fast molecular exchange takes place between the free SA molecules and the SA molecules of micelles. The lifetime of a SA molecule makes 10^{-5} – 10^{-7} s in a micelle [1–3]. At the same time, the micelles themselves are not long-lived formations: they can both disintegrate and arise again. A time of half-decay of micelles takes the value within the limits of several milliseconds up to several seconds.

The “drop-wise” model of micelles assumes that their geometrical shape is spherical at small concentrations of the SA [4]. It has been shown by thermodynamic and experimental studies that the micelles seem to be liquids: molecules of SA are movable in micelles as in a usual liquid [2]. It is confirmed by the line width of fine NMR spectra of surfactant solutions, which does not change by transferring a solution from a nonmicellar to a micellar state.

The present work has as its objects to examine by using the proton NMR-MFPG method and fluorimetry the concentration dependence of the self-diffusion coefficient of 4-aza-1-tetradecyl-1-azoniabicyclo[2.2.2]octane bromide (AB-14) in deuterioxide solution for both kinds of the SA – outside the micellar formations and inside them, the distribution of surfactant molecules on these kinds, the dependence of the size and the number of aggregation of micelles, the role of external and internal diffusion of surfactant molecules in the process of exchange between their different kinds.

2. Experimental and theoretical procedures

2.1. Chemicals

Bicyclic surfactant AB-14 (Scheme 1) has been synthesized from a solution of 1,4-diazabicyclo[2.2.2]octane and tetradecyl bromide upon a procedure offered in refs. [5,6]. The deuterated water with ²H isotope content of 99.5% has been taken in order to explore the desired solutions. Cetylpyridinium bromide (CPB) (AppliChem, BioChemica, Germany),

* Corresponding author.

E-mail address: lucia@iopc.ru (L.Y. Zakharova).