

Zeitschrift fur Anorganische und Allgemeine Chemie 2009 vol.635 N11, pages 1613-1619

Complexes of crown ether-containing N-phosphorylated thioamides and thioureas with Coll, ZnII and PdII cations

Safin D., Babashkina M., Sokolov F., Baranov S., Ekkehardt Hahn F., Pape T.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

The reaction of the potassium salts of N-phosphorylated thioureas [4'-benzo-15-crown-5]NHC(S)NHP(Y)(O*i*Pr)₂ (Y = S, HLI; Y = O, HLII) with ZnII and Coll cations in aqueous EtOH leads to complexes of formulae Zn(LI,IIS,Y)₂ (Y = S, 1; Y = O, 2) and Co(LIS,S')₂ (3), while interaction of the potassium salt of N-phosphorylated thioamide [4'-benzo-15-crown-5]C(S)NHP(O) (O*i*Pr)₂ (HLIII) with ZnII in the same conditions leads to the complex Zn(HLIII)(LIII-S,O)₂ (4). The reaction of the potassium salt of crown ether-containing N-phosphorylated bis-thiourea N,N'-[C(S)NHP(O)(O*i*PT)₂] 2-1,10-diaza-18crown-6 (H₂L) with Coll, Zn.II and PdII cations in anhydrous CH₃OH leads to complexes M₂(L-O,A)₂ (M = Co, 5; Zn, 6; M = Pd, 7). Thioamide HLIII was investigated by single-crystal X-ray diffraction. © 2009 Wiley-VCH Verlag GmbH & Co. KGaA.

<http://dx.doi.org/10.1002/zaac.200801418>

Keywords

Chelates, Cobalt, Crown compounds, Palladium, Zinc