Molekuliarnaia biologiia 2016 vol.50 N6, pages 992-998

Activation of biosynthesis of guanyl-specific ribonuclease secreted by Bacillus circulans under salt stress

Kharitonova M., Kipenskaya L., Ilinskaya O. Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

The gene transcription of guanyl-specific ribonucleases (RNases), which provide available phosphate to cells of Bacillus, is controlled by the signal transduction system PhoP-PhoR. However, the biosynthesis of B. circulans RNase does not depend on the signal-transduction regulatory proteins of Pho regulon. It has been found that raising the salt molar concentration in culture medium increases the level of extracellular guanyl-specific ribonuclease Bci synthesized by B. circulans. Sequences homologous to the binding sites of the regulatory protein DegU were found in RNase Bci promoter. The functioning of the DegS-DegU signal transduction system is stimulated by a high salt concentration. Using a strain of B. subtilis that is defective in the DegU regulatory protein, we have shown that the DegS-DegU system participates in the regulation of RNase Bci expression under salt stress.

http://dx.doi.org/10.7868/S0026898416050086

Keywords

Bacillus circulans, biosynthesis activation, guanyl-specific ribonuclease, salt stress