

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

# Metabolism

[www.metabolismjournal.com](http://www.metabolismjournal.com)

## A new hope for obesity management: Boron inhibits adipogenesis in progenitor cells through the Wnt/ $\beta$ -catenin pathway



Ayşegül Doğan<sup>a,b,1</sup>, Selami Demirci<sup>a,c,\*</sup>, Hüseyin Apdik<sup>a</sup>, Omer Faruk Bayrak<sup>d</sup>,  
Sukru Gulluoglu<sup>d</sup>, Emre Can Tuysuz<sup>d</sup>, Oleg Gusev<sup>e,f</sup>, Albert A. Rizvanov<sup>e</sup>,  
Emrah Nikerel<sup>a</sup>, Fikretin Şahin<sup>a</sup>

<sup>a</sup> Department of Genetics and Bioengineering, Faculty of Engineering, Yeditepe University, Kayisdagi Cad. 26 Agustos Yerlesimi, 34755 Atasehir, Istanbul, Turkey

<sup>b</sup> National Cancer Institute, CDBL, NIH, Frederick, MD

<sup>c</sup> National Heart, Lung, and Blood Institute (NHLBI), NIH, Bethesda, MD

<sup>d</sup> Department of Medical Genetics, Yeditepe University Medical School Inonu Mah, Kayisdagi Cad. 26 Agustos Yerlesimi, 34755 Atasehir, Istanbul, Turkey

<sup>e</sup> Institute of Fundamental Medicine and Biology, Kazan Federal University, Kazan, Russia

<sup>f</sup> Riken Innovation Center, Riken, Yokohama, Japan

### ARTICLE INFO

#### Article history:

Received 18 October 2016

Accepted 12 January 2017

#### Keywords:

Boron  
Lipogenesis  
 $\beta$ -catenin  
PPAR $\gamma$   
Adipogenesis

### ABSTRACT

Obesity is a worldwide medical problem resulting in serious morbidity and mortality involving differentiation of pre-adipocytes into mature adipocytes (adipogenesis). Boron treatment has been reported to be associated with weight reduction in experimental animals; however, its effects on pre-adipocyte differentiation and anti-adipogenic molecular mechanisms are unknown.

In this study, we demonstrate the inhibitory activities of boric acid (BA) and sodium pentaborate pentahydrate (NaB) on adipogenesis using common cellular models. Boron treatment repressed the expression of adipogenesis-related genes and proteins, including CCAAT-enhancer-binding protein  $\alpha$  and peroxisome proliferator-activated receptor  $\gamma$ , by regulating critical growth factors and the  $\beta$ -catenin, AKT, and extracellular signal-regulated kinase signaling pathways. In addition, although boron treatment did not induce apoptosis in pre-adipocytes, it depressed mitotic clonal expansion by regulation of cell cycle genes.

Overall, these data offer promising insights into the prevention/treatment of obesity and associated diseases.

Published by Elsevier Inc.

## 1. Introduction

Obesity and associated comorbidities, such as type 2 diabetes, hypertension, cardiovascular disease, cancer, and metabolic syndrome or posttraumatic stress disorder (PTSD), are worldwide health concerns, particularly in developed countries, leading to

increased healthcare costs, morbidity, and mortality [1,2]. An estimated 2.1 billion people worldwide are thought to be overweight or obese, and 2.8 million deaths are associated with obesity annually [3]. Although the exact molecular mechanisms of obesity are not well understood, uncontrolled hyperplasia and hypertrophy of adipocytes, the main units in fat tissue, are

\* Corresponding author at: Genetics and Bioengineering Department, Faculty of Engineering and Architecture, Yeditepe University Kayisdagi, Istanbul, Turkey. Tel.: +90 216 578 0619; fax: +90 216 578 2182.

E-mail addresses: [selami.demirci@nih.gov](mailto:selami.demirci@nih.gov), [selamidemirci@hotmail.com](mailto:selamidemirci@hotmail.com) (S. Demirci).

<sup>1</sup> These authors contributed equally to this work.