

Novel type of isoprenoid membrane anchors: an investigation of binding properties with dipalmitoylphosphatidylcholine vesicles

Faizullin D., Dzyurkevich M., Valiullina Y., Islamov D., Kataeva O., Zuev Y., Plemenkov V., Stoikov I.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

Copyright © 2016 John Wiley & Sons, Ltd. In this work, we present a new type of amphiphilic membrane-anchoring agents that can be easily obtained by the Diels-Alder reaction between terpene myrcene and N-substituted maleimides. The interaction between the compounds and small unilamellar dipalmitoylphosphatidylcholine vesicles was investigated using infrared spectroscopy, microgravimetry, and turbidimetry. The ability of the compounds to embed in the phospholipid membrane was shown to be strongly dependent on the charge of their polar group. The insertion of the compounds studied into the lipid bilayer did not lead to disruption of the dipalmitoylphosphatidylcholine vesicles up to the highest tested drug to lipid molar ratio of 0.5 to 0.6. Low lipid solubilization ability of the compounds as well as their rigid nonplanar structure makes them an interesting alternative to the common membrane-anchoring structural motifs.

<http://dx.doi.org/10.1002/poc.3618>

Keywords

dipalmitoylphosphatidylcholine vesicles, membrane anchors, N-substituted maleimides, terpenoids

References

- [1] C. A. Lipinski, F. Lombardo, B. W. Dominy, P. J. Feeney, *Adv. Drug Delivery Rev.* 2001, 46, 3.
- [2] L. Rajendran, V. Udayar, Z. V. Goodger, *Trends Pharmacol. Sci.* 2012, 33, 215.
- [3] L. Rajendran, A. Schneider, G. Schlechtingen, S. Weidlich, J. Ries, T. Braxmeier, P. Schwille, J. B. Schulz, C. Schroeder, M. Simons, G. Jennings, H. J. Knölker, K. Simons, *Science* 2008, 320, 520.
- [4] P. Ingallinella, E. Bianchi, N. A. Ladwa, Y.-J. Wang, R. Hrin, M. Veneziano, F. Bonelli, T. J. Ketas, J. P. Moore, M. D. Miller, A. Pessia, *Proc. Natl. Acad. Sci. U. S. A.* 2009, 106, 5801.
- [5] E. C. Naumann, S. Göring, I. Ogorek, S. Weggen, B. Schmidt, *Bioorg. Med. Chem. Lett.* 2013, 23, 3852.
- [6] L. Miao, M. Nielsen, J. Thewalt, J. H. Ipsen, M. Bloom, M. J. Zuckermann, O. G. Mouritsen, *Biophys. J.* 2002, 82, 1429.
- [7] A. C. Simonsen, *Biophys. J.* 2008, 94, 3966.
- [8] J. Atkinson, R. F. Epand, R. M. Epand, *Free Radical Bio. Med.* 2008, 44, 739.
- [9] S. Witzke, L. Duelund, J. Kongsted, M. Petersen, O. G. Mouritsen, K. Himanshu, *J. Phys. Chem. B* 2010, 114, 15825.
- [10] T. Rog, M. Pasenkiewicz-Gierula, *Biophys. J.* 2003, 84, 1818.

- [11] B. B. Lorenzetti, G. E. P. Souza, S. J. Sarti, D. S. Filho, S. H. Ferreira, J. Ethnopharmacol 1991, 34(1), 43.
- [12] M. S. Dzyurkevich, K. N. Timofeeva, D. A. Faizullin, Yu. F. Zuev, I. I. Stoikov, V. V. Plemenkov, Mend. Comm. 2014, 24, 224.
- [13] F. Eker, H. O. Durmus, B. G. Akinoglu, F. Severcan, J. Mol. Struct. 1999, 693, 482.
- [14] G. Kateman, H. C. Smit, L. Meites, Anal. Chim. Acta 1983, 152 (C), 61.
- [15] D. R. Gabdrakhmanov, M. A. Voronin, L. Y. Zakharova, A. I. Konovalov, R. N. Khaybullin, I. Y. Strobykina, V. E. Kataev, D. A. Faizullin, N. E. Gogoleva, T. A. Konnova, V. V. Salnikov, Y. F. Zuev, Phys. Chem. Chem. Phys. 2013, 15, 16725.
- [16] Y. Okahata, H. Ebato, Anal. Chem. 1991, 63, 203.
- [17] Bruker. APEX2 Software Suite for Crystallographic Programs, Bruker AXS, Inc., Madison, WI, USA, 2009.
- [18] Bruker. Area detector control and integration software. Version 5.x. In: SMART and SAINT. Madison, Wisconsin (USA): Bruker Analytical X-ray Instruments Inc.; 1996.
- [19] G. M. Sheldrick, SADABS, University of Göttingen, Institut für Anorganische Chemie der Universität, Tammanstrasse 4, D-3400 Göttingen, Germany, 1997.
- [20] G. M. Sheldrick, Acta Crystallogr 2008, 64, 112.
- [21] C. F. Macrae, P. R. Edgington, P. McCabe, E. Pidcock, G. P. Shields, R. Taylor, M. Towler, J. van de Streek, J. Appl. Crystallogr. 2006, 39, 453.
- [22] M. S. Dzyurkevich, D. B. Krivolapov, O. A. Lodochnikova, I. A. Litvinov, V. V. Plemenkov, Rus. J. Gen. Chem. 2011, 81, 1181.
- [23] E. V. Mironova, M. S. Dzyurkevich, O. A. Lodochnikova, D. B. Krivolapov, I. A. Litvinov, V. V. Plemenkov, J. Struct. Chem. 2012, 53, 361.
- [24] S. F. Burlatsky, V. V. Atrazhev, D. V. Dmitriev, V. I. Sultanov, E. N. Timokhina, E. A. Ugolkova, S. Tulyani, A. Vincitore, J. Colloid Interface Sci. 2013, 393, 151.
- [25] T. Inoue, K. Miyakawa, R. Shimosawa, Chem. Phys. Lipids 1986, 42(4), 261.
- [26] R. Koynova, M. Caffrey, Biochim. Biophys. Acta 1998, 1376, 91.
- [27] M. A. Morini, M. B. Sierra, V. I. Pedroni, L. M. Alarcon, G. A. Appignanesi, E. A. Disalvo, Colloids Surf., B 2015, 131, 54.
- [28] Y.-P. Zhang, R. N. A. H. Lewis, R. N. McElhaney, Biophys. J. 1997, 72, 779.
- [29] R. N. A. H. Lewis, R. N. McElhaney, Methods Mol. Biol. 2007, 400, 207.
- [30] R. N. A. H. Lewis, R. N. McElhaney, W. Pohle, H. H. Mantsch, Biophys. J. 1994, 67, 2367.
- [31] T.-X. Xiang, B. D. Anderson, Adv. Drug Delivery Rev. 2006, 58, 1357.
- [32] S. A. Mendanha, A. Alonso, Biophys. Chem. 2015, 198, 45.
- [33] M. Keller, A. Kerth, A. Blume, BBA-Biomembranes 1997, 1326, 178.