

Structural and dynamic features of water and amorphous ice

Khusnutdinoff R.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2017, Pleiades Publishing, Ltd. Structural properties and microscopic dynamics of water and amorphous ice have been studied by the molecular dynamics method. It has been found that the distribution function of the tetrahedrality parameter exhibits two ranges, which correspond to local molecular formations with low and high degrees of tetrahedrality. The number of molecular clusters with a high degree of tetrahedrality grows as temperature decreases. It has been shown that the vibrational density of states comprises two vibrational modes. A low-frequency vibrational mode strongly depends on pressure and is almost independent of temperature, while a high-frequency mode is relevant to the pressure-independent heat motion of molecules. The geometric criterion of hydrogen bonds has been used to evaluate their continuous lifetime as depending on temperature for molecules with different coordination values. The average lifetime of a hydrogen bond substantially depends on the coordination of molecules, with the temperature dependence of the coordination obeying the activation dynamics.

<http://dx.doi.org/10.1134/S1061933X17010070>

References

- [1] Zheligovskaya, E.A. and Malenkov, G.G., *Usp. Khim.*, 2006, vol. 75, p. 64.
- [2] Salzmann, Ch.G., Radaelli, P.G., Slater, B., and Finney, J.L., *Phys. Chem. Chem. Phys.*, 2011, vol. 13, p. 18468.
- [3] Angell, C.A., *Annu. Rev. Phys. Chem.*, 2004, vol. 55, p. 559.
- [4] Khusnutdinoff, R.M. and Mokshin, A.V., *J. Non-Cryst. Solids*, 2011, vol. 357, p. 1677.
- [5] Khusnutdinoff, R.M. and Mokshin, A.V., *Physica A (Amsterdam)*, 2012, vol. 391, p. 2842.
- [6] Kumar, P., Wikfeldt, K.Th., Schlesinger, D., Pettersson, L.G.M., and Stanley, H.E., *Sci. Rep.*, 2013, vol. 3, p. 1980.
- [7] Paciaroni, A., Bizzarri, A.R., and Cannistraro, S., *Phys. Rev. E: Stat. Phys., Plasmas, Fluids, Relat. Interdiscip. Top.*, 1999, vol. 60, p. R2476.
- [8] Li, J.C. and Kolesnikov, A.I., *Physica B (Amsterdam)*, 2002, vols. 316-317, p. 493.
- [9] Mishima, O., Calvert, L.D., and Whalley, E., *Nature (London)*, 1985, vol. 314, p. 76.
- [10] Malenkov, G.G., *Zh. Strukt. Khim.*, 2006, vol. 47, p. S5.
- [11] Falenty, A., Hansen, Th.C., and Kuhs, W.F., *Nature (London)*, 2014, vol. 516, p. 231.
- [12] Huang, Y., Zhu, Ch., Wang, L., Cao, X., Su, Y., Jiang, X., Meng, Sh., Zhao, J., and Cheng Zeng, X., *Sci. Adv.*, 2016, vol. 2, p. 1.
- [13] Loerting, Th. and Giovambattista, N., *J. Phys.: Condens. Matter*, 2006, vol. 18, p. R919.
- [14] Mishima, O., *Jpn. Acad. B-Phys.*, 2010, vol. 86, p. 165.
- [15] Mishima, O. and Stanley, H.E., *Nature (London)*, 1998, vol. 396, p. 329.

- [16] Abascal, J.L.F. and Vega, C., *J. Chem. Phys.*, 2005, vol. 123, p. 234505.
- [17] Pi, H.L., Aragonés, J.L., Vega, C., Noya, E.G., Abascal, J.L.F., González, M.A., and McBride, C., *Mol. Phys.*, 2009, vol. 107, p. 365.
- [18] Khusnutdinov, R.M. and Mokshin, A.V., *Bull. Russ. Acad. Sci.: Phys.*, 2010, vol. 74, p. 640.
- [19] Allen, M.P. and Tildesley, D.J., *Computer Simulation of Liquids*, Oxford: Clarendon, 1987.
- [20] Mallamace, F., Corsaro, C., and Stanley, H.E., *PNAS*, 2013, vol. 110, p. 4899.
- [21] Khusnutdinoff, R.M., *Colloid J.*, 2013, vol. 75, p. 726.
- [22] Khusnutdinoff, R.M., *Colloid J.*, 2016, vol. 78, p. 225.
- [23] Mokshin, A.V., Yulmetyev, R.M., Khusnutdinov, R.M., and Hänggi, P., *Phys. Solid State*, 2006, vol. 48, p. 1760.
- [24] Sarkisov, G.N., *Usp. Fiz. Nauk*, 2002, vol. 172, p. 647.
- [25] Mokshin, A.V., Zabegaev, S.O., and Khusnutdinov, R.M., *Phys. Solid State*, 2011, vol. 53, p. 570.
- [26] Wendt, H.R. and Abraham, F.F., *Phys. Rev. Lett.*, 1978, vol. 41, p. 1244.
- [27] Debenedetti, P.G. and Stanley, H.E., *Phys. Today*, 2003, vol. 56, p. 40.
- [28] Mokshin, A.V., Yulmetyev, R.M., Khusnutdinoff, R.M., and Hänggi, P., *J. Phys.: Condens. Matter*, 2007, vol. 19, p. 046209.
- [29] Medvedev, N.N., Voloshin, V.P., and Naberukhin, Yu.I., *Zh. Strukt. Khim.*, 1989, vol. 30, p. 98.
- [30] Naberukhin, Yu.I. and Voloshin, V.P., *Zh. Strukt. Khim.*, 2006, vol. 47, p. S129.
- [31] Pereyra, R.G., Bermudez di Lorenzo, A.J., Malaspina, D.C., and Carignano, M.A., *Chem. Phys. Lett.*, 2012, vol. 538, p. 35.
- [32] Stillinger, F.H. and Rahman, A., *J. Chem. Phys.*, 1972, vol. 57, p. 1281.
- [33] Swiatla-Wojcik, D., *Chem. Phys.*, 2007, vol. 342, p. 260.
- [34] Kumar, R., Schmidt, J.R., and Skinner, J.L., *J. Chem. Phys.*, 2007, vol. 126, p. 204107.
- [35] Rapaport, D.C., *Mol. Phys.*, 1983, vol. 50, p. 1151.
- [36] Kalinichev, A.G. and Bass, J.D., *Chem. Phys. Lett.*, 1994, vol. 231, p. 301.
- [37] Malenkov, G.G. and Tytik, D.L., *Izv. Akad. Nauk, Ser. Fiz.*, 2000, vol. 64, p. 1469.
- [38] Starr, F.W., Nielsen, J.K., and Stanley, H.E., *Phys. Rev. E: Stat. Phys., Plasmas, Fluids, Relat. Interdiscip. Top.*, 2000, vol. 62, p. 579.
- [39] Luzar, A. and Chandler, D., *Phys. Rev. Lett.*, 1996, vol. 76, p. 928.
- [40] Luzar, A. and Chandler, D., *Nature (London)*, 1996, vol. 379, p. 55.
- [41] Luzar, A., *J. Chem. Phys.*, 2000, vol. 113, p. 10663.
- [42] Naberukhin, Yu.I. and Voloshin, V.P., *Z. Phys. Chem.*, 2009, vol. 223, p. 1119.
- [43] Conde, O. and Teixeira, J., *J. Phys.*, 1983, vol. 44, p. 525.
- [44] Khusnutdinov, R.M. and Mokshin, A.V., *JETP Lett.*, 2014, vol. 100, p. 39.
- [45] Hansen, J.P. and McDonald, I.R., *Theory of Simple Liquids*, New York: Academic, 2006.
- [46] Mokshin, A.V., Chvanova, A.V., and Khusnutdinov, R.M., *Teor. Mat. Fiz.*, 2012, vol. 171, p. 135.
- [47] Gallo, P. and Rovere, M., *J. Chem. Phys.*, 2012, vol. 37, p. 164503.
- [48] Götze, W., *Complex Dynamics of Glass-Forming Liquids*, Oxford: Oxford Univ. Press, 2009.
- [49] Khusnutdinov R.M., Mokshin A.V., and Khadeev I.I., *J. Surf. Invest.: X-ray, Synchrotron Neutron Tech.*, 2014, vol. 8, p. 84.
- [50] Cunsolo, A., Kodituwakku, C.N., Bencivenga, F., Frontzek, M., Leu, B.M., and Said, A.H., *Phys. Rev. B*, 2012, vol. 85, p. 174305.