Russian Meteorology and Hydrology 2017 vol.42 N3, pages 168-180

Comparison of the results of ⁸⁵Kr transport modeling with the ACURATE field experiment data

Rubinshtein K., Safronov A., Pripachkin D., Ignatov R., Emelina S., Nabokova E., Kurbatova M., Blagodatskikh D., Arutyunyan R., Sorokovikova O., Semenov V. *Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia*

Abstract

© 2017, Allerton Press, Inc.Currently the Nuclear Safety Institute of the Russian Academy of Sciences (NSI RAS) jointly with the Hydrometcenter of Russia is developing the system for forecasting the transfer of radio-active substances in the atmosphere in case of radiation accidents at Russian nuclear power plants. The operation of the system is based on the numerical hydrodynamic model which allows forecasting meteorological parameters and is coupled with the mesoscale dispersion model of the transfer ofradioactive substances in the atmosphere. The results are presented of 85Kr transport modeling under the conditions of the ACURATE experiment with three transport models: FLEXPART, HYSPLIT, and the model from the NOSTRADAMUS software package. It is demonstrated that all three Lagrangian models can give a qualitative description of concentration fields from the ACURATE experiment with the best value of the RANK metric (2.5) based on three statistics.

http://dx.doi.org/10.3103/S1068373917030049

Keywords

contaminant transfer, NPP, radionuclide, transport-dispersion models, WRF-ARW model

References

- R. V. Arutyunyan, L. A. Bol'shov, D. A. Pripachkin, et al., "Actionable Analysis of the Accident at the Fukushima-1 Nuclear Power Plant (Japan) and Prediction of Its Consequences," Atomnaya Energiya, No. 3, 112 (2012) [Atomic Energy, No. 3, 112 (2012)].
- [2] R. V. Arutyunyan, D. A. Pripachkin, O. S. Sorokovikova, et al., "Cross-border Transport during Small Discharges of Radioactive Substances into the Atmosphere," Atomnaya Energiya, No. 2, 117 (2014) [Atomic Energy, No. 2, 117 (2014)].
- [3] V. V. Belikov, V. M. Goloviznin, Yu. V. Katyshkov, et al., "NOSTRADAMUS: A Computer System for Predicting Radiation Conditions. Verification of the Model of Atmospheric Transport of Contaminants," in Modeling of the Transport of Radionuclides in the Environment: Proceedings of Nuclear Safety Institute (IBRAE) (Nauka, Moscow, 2008) [in Russian].
- [4] K. G. Rubinshtein, R. Yu. Ignatov, E. V. Nabokova, et al., "Use of the Regional Non-hydrostatic Model for Determining Meteorological Variables in Forecasting Radionuclides Propagation and Fallout," Meteorol. Gidrol., No. 8 (2009) [Russ. Meteorol. Hydrol., No. 8, 34 (2009)].
- [5] D. M. Barker, W. Huang, Y. R. Guo, and Q. N. Xiao, "A Three-dimensional (3DVAR) Data Assimilation Systems for Use with MM5: Implementation and Initial Results," Mon. Wea. Rev., 132 (2004).
- [6] H. W. Barker, J. N. S. Cole, et al., "The Monte Carlo Independent Column Approximation: An Assessment Using Several Global Atmospheric Models," Quart. J. Roy. Meteorol. Soc., 134 (2008).

- [7] H. W. Barker, G. L. Stephens, P. T. Partain, et al., "Assessing 1d Atmospheric Solar Radiative Transfer Models: Interpretation and Handling of Unresolved Clouds," J. Climate, 16 (2003).
- [8] A. K. Betts, "A New Convective Adjustment Scheme. Observational and Theoretical Basis," Quart. J. Roy. Meteorol. Soc., 112 (1986).
- [9] L. Bolshov and V. Strizhov, "SOCRAT—The System of Codes for Realistic Analysis of Severe Accidents," in Proceedings of International Congress on Advances in Nuclear Power Plants (ICAPP'06), Reno, NV, USA, June 4-8, 2006.
- [10] J. Brioude, D. Arnold, A. Stohl, et al., "The Lagrangian Particle Dispersion Model FLEXPART-WRF Version 3.1," Geosci. Model Dev., 6 (2013).
- [11] F. Chen and J. Dudhia, "Coupling an Advanced Land Surface—Hydrology Model with the Penn State-NCAR MM5 Modeling System. Part I: Model Implementation and Sensitivity," Mon. Wea. Rev., 129 (2001).
- [12] Dataset DATEM (Data Archive of Tracer Experiments and Meteorology), http://www.arl.noaa.gov/DATEM.php (Accessed March 2015).
- [13] R. R. Draxler, "Demonstration of a Global Modeling Methodology to Determine the Relative Importance of Local and Long-distance Sources," Atmos. Environ., 41 (2007).
- [14] R. R. Draxler, HYSPLIT4 User's Guide, NOAA Tech. Memo. ERL ARL-230 (Air Resources Laboratory, Silver Spring, MD, 1999), www.arl.noaa.gov/documents/reports/arl-230.pdf.
- [15] R. R. Draxler and G. D. Hess, "An Overview of the HYSPLIT_4 Modelling System for Trajectories, Dispersion, and Deposition," Austral. Meteorol. Mag., 47 (1998).
- [16] R. R. Draxler and G. D. Hess, Description of the HYSPLIT 4 Modeling System, NOAA Tech. Memo. ERL ARL-224 (Air Resources Laboratory, Silver Spring, MD, 1997), www.arl.noaa.gov/documents/reports/arl-224.pdf.
- [17] R. R. Draxler and G. D. Rolph, "Evaluation of the Transfer Coefficient Matrix (TCM) Approach to Model the Atmospheric Radionuclide Air Concentrations from Fukushima," J. Geophys. Res. Atmos., 117 (2012).
- [18] M. B. Ek, K. E. Mitchell, Y. Lin, et al., "Implementation of NOAH Land Surface Model Advances in the NCEP Operational Mesoscale Eta Model," J. Geophys. Res., No. D22, 108 (2003).
- [19] ERA-Interim Data of ECMWF (The European Centre for Medium-range Weather Forecasts), http://apps.ecmwf. int/datasets/data/interim-full-daily/?levtype=pl (Accessed March 2015).
- [20] Q. Fu, "An Accurate Parameterization of the Solar Radiative Properties of Cirrus Clouds for Climate Models," J. Climate, 9 (1996).
- [21] S. R. Hanna, "Applications in Air Pollution Modeling," in Atmospheric Turbulence and Air Pollution Modelling (D. Reidel Publishing Company, Dordrecht, Holland, 1982).
- [22] Y. X. Hu and K. Stamnes, "An Accurate Parameterization of the Radiative Properties of Water Clouds Suitable for Use in Climate Models," J. Climate, 6 (1993).
- [23] M. J. Iacono, J. S. Delamere, E. J. Mlawer, et al., "Radiative Forcing by Long-lived Greenhouse Gases: Calculations with the AER Radiative Transfer Models," J. Geophys. Res., 113 (2008).
- [24] Integrated Global Radiosonde Archive (IGRA), http://www.ncdc.noaa.gov/oa/cliamte/igra/index.php (Accessed March 2015).
- [25] Integrated Surface Database (ISD), www.ncdc.noaa.gov (Accessed March 2015).
- [26] Z. I. Janjic, "Comments on "Development and Evaluation of a Convection Scheme for Use in Climate Models," J. Atmos. Sci., No. 21, 57 (2000).
- [27] Z. I. Janjic, "The Step-Mountain Eta Coordinate Model: Further Developments of the Convection, Viscous Sublayer, and Turbulence Closure Schemes," Mon. Wea. Rev., 122 (1994).
- [28] L. Mahrt and M. Ek, "The Influence of Atmospheric Stability on Potential Evaporation," J. Climate Appl. Meteorol., No. 2, 23 (1984).
- [29] E. J. Mlawer, S. J. Taubman, P. D. Brown, et al., "Radiative Transfer for Inhomogeneous Atmospheres: RRTM, a Validated Correlated-k Model for the Long-wave," J. Geophys. Res., No. D14, 102 (1997).
- [30] Model MM5 Description (Fifth-generation NCAR/Penn State Mesoscale Model), www2.mmm.ucar.edu/mm5/ mm5 -home.html.
- [31] B. E. Moroz, H. L. Beck, A. Bouville, and S. L. Simon, "Predictions of Dispersion and Deposition of Fallout from Nuclear Testing Using the NOAA-HYSPLIT Meteorological Model," Health Phys., No. 2, 99 (2010).
- [32] M. Nakanishi, "Improvement of Mellor-Yamada Turbulence Closure Model Based on Large-eddy Simulation Data," Boundary-Layer Meteorol., 99 (2001).
- [33] M. Nakanishi and H. Niino, "An Improved Mellor-Yamada Level-3 Model with Condensation Physics: Its Design and Verification," Boundary-Layer Meteorol., 112 (2004).
- [34] M. Nakanishi and H. Niino, "An Improved Mellor-Yamada Level-3 Model: Its Numerical Stability and Application to a Regional Prediction of Advection Fog," Boundary-Layer Meteorol., 119 (2006).
- [35] M. Nakanishi and H. Niino, "Development of an Improved Turbulence Closure Model for the Atmospheric Boundary Layer," J. Meteorol. Soc. Japan, 87 (2009).

- [36] NARR Dataset, National Centers for Environmental Prediction/National Weather Service/NOAA/U.S. Department of Commerce. 2005, Updated Monthly. NCEP North American Regional Reanalysis (NARR). Research Data Archive at the National Center for Atmospheric Research, Computational and Information Systems Laboratory, http://rda.ucar.edu/datasets/ds608.0Z (Accessed March 2015).
- [37] J. P. Olson and J. M. Brown, "A Comparison of Two Mellor-Yamada-based PBL Schemes in Simulating a Hybrid Barrier Jet," in Proceedings of the 23rd Conference on Weather Analysis and Forecasting/19th Conference on Numerical Weather Prediction, Omaha, NE, 2009.
- [38] H.-L. Pan and L. Mahrt, "Interaction between Soil Hydrology and Boundary-layer Development," Boundary-Layer Meteorol., No. 1/2, 38 (1987).
- [39] R. Pincus, H. W. Barker, and J.-J. Morcrette, "A Fast, Flexible, Approximate Technique for Computing Radiative Transfer in Inhomogeneous Cloud Fields," J. Geophys. Res. (2003).
- [40] J. Reisner, R. M. Rasmussen, and R. T. Bruintjes, "Explicit Forecasting of Supercooled Liquid Water in Winter Storms Using the MM5 Mesoscale Model," Quart. J. Roy. Meteorol. Soc., No. 548, 124 (1998).
- [41] W. C. Skamarock, "Positive-definite and Monotonie Limiters for Unrestricted-timestep Transport Schemes," Mon. We a. Rev., 134 (2006).
- [42] W. C. Skamarock, J. B. Klemp, J. Dudhia, et al., A Description of the Advanced Research WRF Version 3, NCAR Tech. Note NCAR/TN-475+STR (2008).
- [43] J. Smagorinsky, "General Circulation Experiments with the Primitive Equations: 1. The Basic Experiment," Mon. Wea. Rev., 91 (1963).
- [44] A. Stohl, "Computation, Accuracy and Applications of Trajectories—a Review and Bibliography," Atmos. Environ., 32 (1998).
- [45] A. Stohl, C. Forster, A. Frank, et al., "Technical Note: The Lagrangian Particle Dispersion Model FLEXPART Version 6.2," Atmos. Chem. Phys., 5 (2005).
- [46] A. Stohl, M. Hittenberger, and G. Wotawa, "Validation of the Lagrangian Particle Dispersion Model FLEXPART against Large Scale Tracer Experiment Data," Atmos. Environ., 32 (1998).
- [47] A. Stohl, P. Seibert, G. Wotawa, et al., "Xenon-133 and Caesium-137 Releases into the Atmosphere from the Fukushima Dai-ichi Nuclear Power Plant: Determination of the Source Term, Atmospheric Dispersion, and Deposition," Atmos. Chem. Phys., 12 (2012).
- [48] G. Thompson and T. Eidhammer, "A Study of Aerosol Impacts on Clouds and Precipitation Development in a Large Winter Cyclone," J. Atmos. Sci., 71 (2014).
- [49] G. Thompson, P. R. Field, R. M. Rasmussen, and W. D. Hall, "Explicit Forecasts of Winter Precipitation Using an Improved Bulk Microphysics Scheme. Part II: Implementation of a New Snow Parameterization," Mon. Wea. Rev., 136 (2008).
- [50] G. Thompson, R. M. Rasmussen, and K. Manning, "Explicit Forecasts of Winter Precipitation Using an Improved Bulk Microphysics Scheme. Part I: Description and Sensitivity Analysis," Mon. Wea. Rev., 132 (2004).
- [51] L. J. Wicker and W. C. Skamarock, "Time-splitting Methods for Elastic Models Using Forward Time Schemes," Mon. Wea. Rev., 130 (2002).