



ELSEVIER

Available online at www.sciencedirect.com**ScienceDirect**

Russian Geology and Geophysics 57 (2016) 653–671

RUSSIAN GEOLOGY
AND GEOPHYSICSwww.elsevier.com/locate/rgg

The 1501 Ma Kuonamka Large Igneous Province of northern Siberia: U–Pb geochronology, geochemistry, and links with coeval magmatism on other crustal blocks

R.E. Ernst^{a,b,*}, A.V. Okrugin^c, R.V. Veselovskiy^{d,e}, S.L. Kamo^f, M.A. Hamilton^f,
V. Pavlov^{d,g}, U. Söderlund^h, K.R. Chamberlainⁱ, C. Rogers^a^a Department of Earth Sciences, Carleton University, Ottawa, ON, K1S 5B6, Canada^b Faculty of Geology and Geography, Tomsk State University, pr. Lenina 36, Tomsk, 634050, Russia^c Diamond & Precious Metal Geology Institute, Russian Academy of Sciences, pr. Lenina 39, Yakutsk, 677000, Russia^d Institute of Physics of the Earth, Russian Academy of Sciences, ul. Bol'shaya Gruzinskaya 10, build. 1, 123995, Moscow, Russia^e Geological Department, Lomonosov Moscow State University, ul. Leninskie Gory 1, 119991, Moscow, Russia^f Jack Satterly Geochronology Laboratory, University of Toronto, Toronto, ON, M5S 3B1, Canada^g Kazan Federal University, ul. Kremlevskaya 18, Kazan, 420000, Russia^h Department of Geology, Lund University, Lund, 223 62, Swedenⁱ Department of Geology and Geophysics, University of Wyoming, Laramie, Wyoming, 82071, USA

Received 29 December 2015; accepted 29 January 2016

Abstract

A new large igneous province (LIP), the 1501 ± 3 Ma Kuonamka LIP, extends across 700 km of northern Siberia and is linked with coeval dikes and sills in the formerly attached São Francisco craton (SFC)–Congo craton to yield a short-duration event 2000 km across. The age of the Kuonamka LIP can be summarized as 1501 ± 3 Ma (95% confidence), based on 7 U–Pb ID-TIMS ages (6 new herein) from dolerite dikes and sills across the Anabar shield and within western Riphean cover rocks for a distance of 270 km. An additional sill yielded a SIMS (CAMECA) age of 1483 ± 17 Ma and sill in the Olenek uplift several hundred kilometers farther east, a previous SIMS (SHRIMP) age of ca. 1473 Ma was obtained on a sill; both SIMS ages are within the age uncertainty of the ID-TIMS ages. Geochemical data indicate a tholeiitic basalt composition with low MgO (4–7 wt%) within-plate character based on trace element classification diagrams and source between E-MORB and OIB with only minor contamination from crust or metasomatized lithospheric mantle. Two subgroups are distinguished: Group 1 has gently sloping LREE ((La/Sm)_{PM} = 1.9) and HREE ((Gd/Yb)_{PM} = 1.8) patterns, slightly negative Sr and moderate TiO₂ (2.2 wt%), and Group 2 has steeper LREE ((La/Sm)_{PM} = 2.3) and HREE ((Gd/Yb)_{PM} = 2.3), strong negative Sr anomaly, is higher in TiO₂ (2.7 wt%), and is transitional from tholeiitic to weakly alkaline in composition. The slight differences in REE slopes are consistent with Group 2 on average melting at deeper levels. Proposed reconstructions of the Kuonamka LIP with 1500 Ma magmatism of the SFC–Congo craton are supported by a geochemical comparison. Specifically, the chemistry of the Chapada Diamantina and Curaça dikes of the SFC can be linked to that of Groups 1 and 2, respectively, of the Kuonamka LIP and are consistent with a common mantle source between EMORB and OIB and subsequent differentiation history. However, the coeval Humpata sills and dikes of the Angola block of the Congo craton represent a different magma batch.

© 2016, V.S. Sobolev IGM, Siberian Branch of the RAS. Published by Elsevier B.V. All rights reserved.

Keywords: magmatism; dikes; sills; igneous province; northern Siberia

Introduction

Large Igneous Provinces (LIPs) represent large volume (>0.1 Mkm³; frequently above >1 Mkm³), mainly mafic (ultramafic) magmatic events of intraplate affinity, that occur

in both continental and oceanic settings, and are typically of short duration (<5 m.y.) or consist of multiple short pulses over a maximum of a few 10s of m.y. (Ernst, 2014 and references therein). They comprise volcanic packages (flood basalts), and a plumbing system of dikes, sills and layered intrusions, and can be associated with silicic magmatism, carbonatites and kimberlites. LIPs are linked with continental

* Corresponding author.

E-mail address: Richard.Ernst@ErnstGeosciences.com (R.E. Ernst)