

Test determinations of paleointensity in historical lavas of Kamchatka

Zhidkov G., Shcherbakov V., Dolotov A., Smirnov M., Ovsyannikov A., Plechov P.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2017, Pleiades Publishing, Ltd. The reliability of the Thellier method for determining the paleointensity of a geomagnetic field is explored on recent igneous rocks of Kamchatka. The main magnetic mineral in the studied rocks is titanomagnetite with different degree of oxidation. It is obtained that the reliability of the results can be assessed based on the deviations of the check points of the partial thermoremanent magnetization (pTRM) during the Thellier experiment. Besides, for different rocks, it is found that the stability of titanomagnetites to heating during the experiments can be insufficient for validating the reliability of the results of paleointensity determination; however, at the same time, the reliability may depend on the initial (oxidation) state of the magnetic minerals of the studied rocks.

<http://dx.doi.org/10.1134/S1069351316060136>

References

- [1] Abokhodair, A.A., The accuracy of the Thellier's technique for the determination of paleointensities of the Earth's magnetic field, Ph.D. Thesis. Santa Cruz: University of California, 1977.
- [2] Burakov, K.S., The method for determining the geomagnetic field strength from thermal demagnetization curves In and Irt, Materialy IX konferentsii po voprosam postoyannogo geomagnitnogo polya, magnetizmu gornykh porod i paleomagnetizmu. Ch. 2 (Proc. 9th Conf. on the Problems of the Main Geomagnetic Field, Rock Magnetism, and Palaeomagnetism), Baku, 1973, part 2, pp. 56–57.
- [3] Chauvin, A., Roperch, P., and Levi, S., Reliability of geomagnetic paleointensity data: the effects of the NRM fraction and concave-up behavior on paleointensity determinations by the Thellier method, Phys. Earth Planet. Inter., 2005, vol. 150, no. 4, pp. 265–286. <http://dx.doi.org/doi 10.1016/j.pepi.2004.11.008>
- [4] Coe, R.S., The determination of paleointensities of the Earth's magnetic field with special emphasize on mechanisms which could cause nonideal behavior in Thellier's method, J. Geomagn. Geoelectr., 1967, vol. 19, no. 3, pp. 157–178.
- [5] Coe, R.S., Grommé, S., and Mankinen, E.A., Geomagnetic paleointensities from radiocarbon-dated lava flows on Hawaii and the question of the Pacific nondipole low, J. Geophys. Res., 1978, vol. 83, no. B4, pp. 1740–1756. doi [10.1029/JB083iB04p01740](https://doi.org/10.1029/JB083iB04p01740)
- [6] Cottrell, R.D. and Tarduno, J.A., Geomagnetic paleointensity derived from single plagioclase crystals, Earth Planet. Sci. Lett., 1999, vol. 169, nos. 1–2, pp. 1–5. doi [10.1016/S0012-821X\(99\)00068-0](https://doi.org/10.1016/S0012-821X(99)00068-0)
- [7] Day, R., Fuller, M., and Schmidt, V.A., Hysteresis properties of titanomagnetites: grain-size and compositional dependence, Phys. Earth Planet. Inter., 1977, vol. 13, no. 4, pp. 260–267. doi [10.1016/0031-9201\(77\)90108-x](https://doi.org/10.1016/0031-9201(77)90108-x)
- [8] Dekkers, M.J. and Bohnel, H.N., Reliable absolute palaeointensities independent of magnetic domain state, Earth Planet. Sci. Lett., 2006, vol. 248, nos. 1–2, pp. 508–517. doi [10.1016/j.epsl.2006.05.040](https://doi.org/10.1016/j.epsl.2006.05.040)
- [9] Dunlop, D.J. and Ozdemir, O., Effect of grain size and domain state on thermal demagnetization tails, Geophys. Rev. Lett., 2000, vol. 27, pp. 1311–1314. doi [10.1029/1999GL008461](https://doi.org/10.1029/1999GL008461)
- [10] Fabian, K., Shcherbakov, V.P., and McEnroe, S.A., Measuring the Curie temperature, Geochem. Geophys. Geosyst., 2013, vol. 14, no. 4, pp. 947–961.

- [11] Herrero-Bervera, E. and Valet, J.-P., Testing determinations of absolute paleointensity from the 1955 and 1960 Hawaiian flows, *Earth Planet. Sci. Lett.*, 2009, vol. 287, nos. 3–4, pp. 420–433. <http://dx.doi.org/doi/10.1016/j.epsl.2009.08.035>
- [12] Hill, M.J. and Shaw, J., Magnetic field intensity study of the 1960 Kilauea lava flow, Hawaii, using the microwave palaeointensity technique, *Geophys. J. Int.*, 2000, vol. 142, no. 2, pp. 487–504. doi 10.1046/j.1365-246x.2000.00164.x
- [13] Hoffman, K.A., Constantine, V.L., and Morse, D.L., Determination of absolute palaeointensity using a multi-specimen procedure, *Nature*, 1989, vol. 339, no. 6222, pp. 295–297. doi 10.1038/339295a0
- [14] Kono, M., Reliability of palaeointensity methods using alternating field demagnetization and anhysteretic remanence, *Geophys. J. Int.*, 1978, vol. 54, pp. 241–261. doi 10.1111/j.1365-246X.1978.tb04258.x
- [15] McClelland, E. and Briden, J.C., An improved methodology for thellier-type paleointensity determination in igneous rocks and its usefulness for verifying primary thermoremanence, *J. Geophys. Res.: Solid Earth*, 1996, vol. 101, no. B10, pp. 21995–22013. doi 10.1029/96jb02113
- [16] Merrill, R.T., McElhinny, M.W., and McFadden, P.L., *The Magnetic Field of the Earth: Paleomagnetism, the Core, and the Deep Mantle*, Int. Geophys. Ser., San Diego: Academic, 1996.
- [17] Nagata, T., *Rock Magnetism*, Tokyo: Maruzen, 1961.
- [18] Nagata, T., Arai, Y., and Momose, K., Secular variation of the geomagnetic total force during the last 5000 years, *J. Geophys. Res.*, 1963, vol. 68, pp. 5277–5282.
- [19] Nishitani, T. and Kono, M., Curie temperature and lattice constant of oxidized titanomagnetite, *Geophys. J. Int.*, 1983, vol. 74, no. 2, pp. 585–600. doi 10.1111/j.1365-246X.1983.tb01890.x
- [20] Pick, T. and Tauxe, L., Geomagnetic palaeointensities during the Cretaceous normal superchron measured using submarine basaltic glass, *Nature*, 1993, vol. 366, no. 6452, pp. 238–242. doi 10.1038/366238a0
- [21] Prevot, M., Mankinen, E.A., Coe, R.S., and Gromme, S.C., The Steens Mountain (Oregon) geomagnetic polarity transition: 2. Field intensity variations and discussion of reversal models, *J. Geophys. Res.*, 1985, vol. 90, no. B12, pp. 10417–10448.
- [22] Riisager, P. and Riisager, J., Detecting multidomain magnetic grains in Thellier palaeointensity experiments, *Phys. Earth Planet. Inter.*, 2001, vol. 125, nos 1–4, pp. 111–117. doi 10.1016/s0031-9201(01)00236-9
- [23] Rolph, T.C. and Shaw, J., A new method of paleofield magnitude correction for thermally altered samples and its application to Lower Carboniferous lavas, *Geophys. J. Int.*, 1985, vol. 80, pp. 773–781. doi 10.1111/j.1365-246X.1985.tb05124.x
- [24] Selkin, P.A. and Tauxe, L., Long-term variations in palaeointensity, *Philos. Trans. R. Soc.*
- [25] London A, 2000, vol. 358, no. 1768, pp. 1065–1088. doi 10.1098/rsta.2000.0574
- [26] Shaw, J., A new method of determining the magnitude of the palaeomagnetic field: application to five historic lavas and five archaeological samples, *Geophys. J. R. Astron. Soc.*, 1974, vol. 39, pp. 133–141. doi 10.1111/j.1365-246X.1974.tb05443.x
- [27] Shaw, J., Walton, D., Yang, S., Rolph, T.C., and Share, J.A., Microwave archeointensities from Peruvian ceramics, *Geophys. J. Int.*, 1996, vol. 124, pp. 241–244.
- [28] Shcherbakova, V.V., Shcherbakov, V.P., and Heider, F., Properties of partial thermoremanent magnetization in pseudosingle domain and multidomain magnetite grains, *J. Geophys. Res.*, 2000, vol. 105, no. B1, pp. 767–781. doi 10.1029/1999jb900235
- [29] Shcherbakova, V.V. and Shcherbakov, V.P., On the criteria of reliability of the results in determining the intensity of the ancient geomagnetic field, *Paleomagnetizm i magnetizm gornykh porod: teoriya, praktika, eksperiment. Materialy mezdunarodnogo seminara (Proc. Int. Symp.: Palaeomagnetism and Rock Magnetism: Theory, Practice, Experiment)*, Kazan: Kazan. Univ., 2004, pp. 117–121.
- [30] Sherwood, G.J., Evaluation of a multi-specimen approach to paleointensity determination, *J. Geomagn. Geoelectr.*, 1991, vol. 43, pp. 341–349.
- [31] Tanaka, H. and Kono, M., Preliminary results and reliability of palaeointensity studies on historical and C dated Hawaiian lavas, *J. Geomagn. Geoelectr.*, 1991, vol. 43, no. 5, pp. 375–388.
- [32] Tanaka, H., Athanassopoulos, J.D.E., Dunn, J.R., and Fuller, M., Paleointensity determinations with measurements at high temperature, *J. Geomagn. Geoelectr.*, 1995, vol. 47, pp. 103–113. doi 10.5636/jgg.47.103
- [33] Tarduno, J.A., Cottrell, R.D., and Smirnov, A.V., High geomagnetic intensity during the mid-Cretaceous from Thellier analyses of single plagioclase crystals, *Science*, 2001, vol. 291, no. 5509, pp. 1779–1783. doi 10.1126/science.1057519
- [34] Thellier, E. and Thellier, O., Sur l'intensité du champ magnétique terrestre dans le passé historique et géologique, *Ann. Geophys.*, 1959, vol. 15, pp. 285–376.
- [35] Tsunakawa, H. and Shaw, J., The Shaw method of paleointensity determinations and its application to recent volcanic rocks, *Geophys. J. Int.*, 1994, vol. 118, pp. 781–787. doi 10.1111/j.1365-246X.1994.tb03999.x

- [36] Valet, J.-P. and Herrero-Bervera, E., Paleointensity experiments using alternating field demagnetization, *Earth Planet. Sci. Lett.*, 2000, vol. 177, nos. 1-2, pp. 43-58. <http://dx.doi.org/>. doi 10.1016/S0012-821X(00)00036-4
- [37] Wilson, R.L., The thermal demagnetization of natural magnetic moments in rocks, *Geophys. J. R. Astron. Soc.*, 1961, vol. 5, no. 1, pp. 45-58. doi 10.1111/j.1365-246X.1961.tb02928.x
- [38] Yamamoto, Y., Tsunakawa, H., and Shibuya, H., Palaeointensity study of the Hawaiian 1960 lava: implications for possible causes of erroneously high intensities, *Geophys. J. Int.*, 2003, vol. 153, no. 1, pp. 263-276. doi 10.1046/j.1365-246X.2003.01909.x