

Mineralization of “non-metabolizable” glucose analogues in soil: potential chemosensory mimics of glucose

Mason-Jones K., Gilmullina A., Kuzyakov Y.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2017 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim Glucose is widely used to study the dynamics of easily available organics in soil. Pure culture studies have revealed that many microorganisms can sense and respond to glucose through chemosensory mechanisms that are not directly reliant on energy catabolism. However, the rapid mineralization of glucose by microorganisms makes it difficult to disentangle its energy effects from such non-catabolic interactions. “Non-metabolizable” glucose analogues have proven useful in mechanistic studies of glucose in pure culture, but have never been applied to complex microbial communities in soil. We sought to determine how their mineralization in soil differs from that of glucose, and whether they have potential as a new approach for investigating chemosensory mechanisms in soil microbiology. We incubated soil from an agricultural Haplic Luvisol under controlled conditions for 24 d and monitored CO₂ efflux after addition of (1) glucose, and three “non-metabolizable” glucose analogues: (2) 2-deoxyglucose (DG), (3) α -methylglucoside (α MG), and (4) 3-O-methyl-glucose (OMG), at three concentration levels, along with a control. All three analogues did in fact produce a large increase in soil CO₂ efflux, but the dynamics of their mineralization differed from the rapid degradation seen for glucose. At medium and high concentrations, CO₂ efflux peaked between 2.5 and 4 d after amendment with DG and α MG, and was delayed by about one week for OMG. The markedly different patterns of mineralization between glucose and OMG offer a new tool for investigating the behavior of glucose in soil. By using OMG as a glucose model, chemosensory mechanisms could be studied with limited interference from energy catabolism.

<http://dx.doi.org/10.1002/jpln.201600204>

Keywords

carbohydrate, LMWOS, microbial metabolism, soil respiration, soil sugars

References

- [1] Adler, J. (1969): Chemoreceptors in bacteria. *Science* 166, 1588–1597.
- [2] Adler, J., Epstein, W. (1974): Phosphotransferase-system enzymes as chemoreceptors for certain sugars in *Escherichia coli* chemotaxis. *Proc. Natl. Acad. Sci. USA*. 71, 2895–2899.
- [3] Anderson, J. P. E., Domsch, K. H. (1978): A physiological method for the quantitative measurement of microbial biomass in soils. *Soil Biol. Biochem.* 10, 215–221.

- [4] Anderson, T. H, Joergensen, R. G. (1997): Relationship between SIR and FE estimates of microbial biomass C in deciduous forest soils at different pH. *Soil Biol. Biochem.* 29, 1033-1042.
- [5] Blagodatskaya, E, Kuzyakov, Y. (2008): Mechanisms of real and apparent priming effects and their dependence on soil microbial biomass and community structure: Critical review. *Biol. Fert. Soils* 45, 115-131.
- [6] De Nobili, M, Contin, M, Mondini, C, Brookes, P. C. (2001): Soil microbial biomass is triggered into activity by trace amounts of substrate. *Soil Biol. Biochem.* 33, 1163-1170.
- [7] Derrien, D, Marol, C, Balesdent, J. (2004): The dynamics of neutral sugars in the rhizosphere of wheat. An approach by C pulse-labelling and GC/C/IRMS. *Plant Soil* 267, 243-253.
- [8] Fischer, H, Ingwersen, J, Kuzyakov, Y. (2010): Microbial uptake of low-molecular-weight organic substances out-competes sorption in soil. *Eur. J. Soil Sci.* 61, 504-513.
- [9] Gunina, A, Kuzyakov, Y. (2015): Sugars in soil and sweets for microorganisms: Review of origin, content, composition and fate. *Soil Biol. Biochem.* 90, 87-100.
- [10] Halpern, Y. S, Lupo, M. (1966): Effect of glucose and other carbon compounds on the transport of α -methylglucoside in *Escherichia coli* K12. *Biochim. Biophys. Acta* 126, 163-167.
- [11] Kirby, J. R. (2009): Chemotaxis-like regulatory systems: unique roles in diverse bacteria. *Annu. Rev. Microbiol.* 63, 45-59.
- [12] Knaff, D. B, Whetstone, R. (1980): Active transport of α -methylglucoside by the photosynthetic bacterium *Chromatium vinosum*. *Arch. Biochem. Biophys.* 203, 697-701.
- [13] Kögel-Knabner, I. (2002): The macromolecular organic composition of plant and microbial residues as inputs to soil organic matter. *Soil Biol. Biochem.* 34, 139-162.
- [14] Komor, E, Tanner, W. (1971): Characterization of the active hexose transport system of *Chlorella vulgaris*. *Biochim. Biophys. Acta* 241, 170-179.
- [15] Koser, S. A, Saunders, F. (1933): The utilization of certain sugars and their derivatives by bacteria. *J. Bacteriol.* 26, 475-488.
- [16] Kramer, S, Marhan, S, Ruess, L, Armbruster, W, Butenschoen, O, Haslwimmer, H, Kuzyakov, Y, Pausch, J, Scheunemann, N, Schoene, J, Schmalwasser, A, Totsche, K. U, Walker, F, Scheu, S, Kandeler, E. (2012): Carbon flow into microbial and fungal biomass as a basis for the belowground food web of agroecosystems. *Pedobiologia* 55, 111-119.
- [17] Lengeler, J. W, Jahreis, K. (2009): Bacterial PEP-dependent carbohydrate: phosphotransferase systems couple sensing and global control mechanisms, in Collin, M, Schuch, R. (eds.): *Bacterial Sensing and Signalling. Contributions to Microbiology*, Vol. 16. Karger, Basel, Switzerland, pp. 65-87.
- [18] Madigan, M. T, Martinko, J. M, Parker, J. (1997): *Brock Biology of Microorganisms*, 8th ed. Prentice Hall, Upper Saddle River, NJ, USA.
- [19] McDermott, J. C, Jennings, D. H. (1976): The relationship between the uptake of glucose and 3-O-methylglucose and soluble carbohydrate and polysaccharide in the fungus *Dendryphiella salina*. *Microbiology* 97, 193-209.
- [20] Pausch, J, Kuzyakov, Y. (2012): Soil organic carbon decomposition from recently added and older sources estimated by C values of CO and organic matter. *Soil Biol. Biochem.* 55, 40-47.
- [21] Persson, C, Olsson, S, Jansson, H. B. (2000): Growth of *Arthrotrix superba* from a birch wood resource base into soil determined by radioactive tracing. *FEMS Microbiol. Ecol.* 31, 47-51.
- [22] R Core Team (2015): *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. Available at: www.R-project.org.
- [23] Scarborough, G. A. (1970): Sugar transport in *Neurospora crassa*. *J. Biol. Chem.* 245, 1694-1698.
- [24] Schneckenberger, K, Demin, D, Stahr, K, Kuzyakov, Y. (2008): Microbial utilization and mineralization of [C]glucose added in six orders of concentration to soil. *Soil Biol. Biochem.* 40, 1981-1988.
- [25] Tarshis, M. A, Bekkouzjin, A. G, Ladygina, V. G, Panchenko, L. F. (1976): Properties of the 3-O-methyl-D-glucose transport system in *Acholeplasma laidlawii*. *J. Bacteriol.* 125, 1-7.
- [26] Tyler, B, Loomis, W. F, Magasanik, B. (1967): Transient repression of the lac operon. *J. Bacteriol.* 94, 2001-2011.
- [27] Zyakun, A. M, Dilly, O. (2005): Use of carbon isotope composition for characterization of microbial activity in arable soils. *Appl. Biochem. Microbiol.* 41, 582-591.