

# Turnover of microbial groups and cell components in soil: $^{13}\text{C}$ analysis of cellular biomarkers

Gunina A., Dippold M., Glaser B., Kuzyakov Y.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

---

## Abstract

© 2017 The Author(s). Microorganisms regulate the carbon (C) cycle in soil, controlling the utilization and recycling of organic substances. To reveal the contribution of particular microbial groups to C utilization and turnover within the microbial cells, the fate of  $^{13}\text{C}$ -labelled glucose was studied under field conditions. Glucose-derived  $^{13}\text{C}$  was traced in cytosol, amino sugars and phospholipid fatty acid (PLFA) pools at intervals of 3, 10 and 50 days after glucose addition into the soil.  $^{13}\text{C}$  enrichment in PLFAs ( $\sim 1.5\%$  of PLFA C at day 3) was an order of magnitude greater than in cytosol, showing the importance of cell membranes for initial C utilization. The  $^{13}\text{C}$  enrichment in amino sugars of living microorganisms at day 3 accounted for 0.57% of total C pool; as a result, we infer that the replacement of C in cell wall components is 3 times slower than that of cell membranes. The C turnover time in the cytosol (150 days) was 3 times longer than in PLFAs (47 days). Consequently, even though the cytosol pool has the fastest processing rates compared to other cellular compartments, intensive recycling of components here leads to a long C turnover time. Both PLFA and amino-sugar profiles indicated that bacteria dominated in glucose utilization.  $^{13}\text{C}$  enrichment decreased with time for bacterial cell membrane components, but it remained constant or even increased for filamentous microorganisms.  $^{13}\text{C}$  enrichment of muramic acid was the 3.5 times greater than for galactosamine, showing a more rapid turnover of bacterial cell wall components compared to fungal. Thus, bacteria utilize a greater proportion of low-molecular-weight organic substances, whereas filamentous microorganisms are responsible for further C transformations. Thus, tracing  $^{13}\text{C}$  in cellular compounds with contrasting turnover rates elucidated the role of microbial groups and their cellular compartments in C utilization and recycling in soil. The results also reflect that microbial C turnover is not restricted to the death or growth of new cells. Indeed, even within living cells, highly polymeric cell compounds are constantly replaced and renewed. This is especially important for assessing C fluxes in soil and the contribution of C from microbial residues to soil organic matter.

<http://dx.doi.org/10.5194/bg-14-271-2017>

---

## References

- [1] Amelung, W., Brodowski, S., Sandhage-Hofmann, A., Bol, R.: Combining biomarker with stable isotope analyses for assessing the transformation and turnover of soil organic matter, in: Advances in Agronomy, edited by: Sparks, D. L., Advances in Agronomy, 100, 155-250, 2008.
- [2] Amelung, W., Kimble, J. M., Samson-Liebig, S., Follett, R. F.: Restoration of microbial residues in soils of the Conservation Reserve Program, Soil Sci. Soc. Am. J., 65, 1704-1709, 2001a.

- [3] Amelung, W., Miltner, A., Zhang, X., Zech, W.: Fate of microbial residues during litter decomposition as affected by minerals, *Soil Sci.*, 166, 598-606, doi:10.1097/00010694-200109000-00003, 2001b.
- [4] Apostel, C., Dippold, M. A., Glaser, B., Kuzyakov, Y.: Biochemical pathways of amino acids in soil: Assessment by position-specific labeling and <sup>13</sup>C-PLFA analysis, *Soil Biol. Biochem.*, 67, 31-40, 2013.
- [5] Baath, E.: Growth rates of bacterial communities in soils at varying pH: A comparison of the thymidine and leucine incorporation techniques, *Microb. Ecol.*, 36, 316-327, doi:10.1007/s002489900118, 1998.
- [6] Bai, Z., Bode, S., Huygens, D., Zhang, X., Boeckx, P.: Kinetics of amino sugar formation from organic residues of different quality, *Soil Biol. Biochem.*, 57, 814-821, doi:10.1016/j.soilbio.2012.08.006, 2013.
- [7] Blagodatskaya, E. and Kuzyakov, Y.: Active microorganisms in soil: critical review of estimation criteria and approaches, *Soil Biol. Biochem.*, 67, 192-211, 2013.
- [8] Bossio, D. A., Scow, K. M., Gunapala, N., Graham, K. J.: Determinants of soil microbial communities, Effects of agricultural management, season, and soil type on phospholipid fatty acid profiles, *Microb. Ecol.*, 36, 1-12, doi:10.1007/s002489900087, 1998.
- [9] Brant, J. B., Sulzman, E.W., Myrold, D. D.: Microbial community utilization of added carbon substrates in response to longterm carbon input manipulation, *Soil Biol. Biochem.*, 38, 2219-2232, doi:10.1016/j.soilbio.2006.01.022, 2006.
- [10] Bremer, E. and Kuikman, P.: Microbial utilization of <sup>14</sup>C-U-glucose in soil is affected by the amount and timing of glucose additions, *Soil Biol. Biochem.*, 26, 511-517, doi:10.1016/0038-0717(94)90184-8, 1994.
- [11] Derrien, D., Marol, C., Balesdent, J.: The dynamics of neutral sugars in the rhizosphere of wheat. An approach by C-13 pulse-labelling and GC/C/IRMS, *Plant Soil*, 267, 243-253, doi:10.1007/s11104-005-5348-8, 2004.
- [12] Dippold, M. A., Boesel, S., Gunina, A., Kuzyakov, Y., Glaser, B.: Improved delta C-13 analysis of amino sugars in soil by ion chromatography-oxidation-isotope ratio mass spectrometry, *Rapid Commun. Mass Sp.*, 28, 569-576, doi:10.1002/rcm.6814, 2014.
- [13] Dungait, J. A. J., Kemmitt, S. J., Michallon, L., Guo, S., Wen, Q., Brookes, P. C., Evershed, R. P.: Variable responses of the soil microbial biomass to trace concentrations of <sup>13</sup>C-labelled glucose, using <sup>13</sup>C-PLFA analysis, *Eur. J. Soil Sci.*, 62, 117-126, doi:10.1111/j.1365-2389.2010.01321.x, 2011.
- [14] Engelking, B., Flessa, H., Joergensen, R. G.: Shifts in amino sugar and ergosterol contents after addition of sucrose and cellulose to soil, *Soil Biol. Biochem.*, 39, 2111-2118, doi:10.1016/j.soilbio.2007.03.020, 2007.
- [15] Fischer, H., Meyer, A., Fischer, K., Kuzyakov, Y.: Carbohydrate and amino acid composition of dissolved organic matter leached from soil, *Soil Biol. Biochem.*, 39, 2926-2935, doi:10.1016/j.soilbio.2007.06.014, 2007.
- [16] Fischer H., Ingwersen J., Kuzyakov Y.: Microbial uptake of low-molecular-weight organic substances outcompetes sorption in soil, *Eur. J. Soil Sci.*, 61, 504-513, 2010.
- [17] Frostegard, A., Tunlid, A., Baath, E.: Microbial biomass measured as total lipid phosphate in soils of different organic content, *J. Microbiol. Meth.*, 14, 151-163, doi:10.1016/0167-7012(91)90018-I, 1991.
- [18] Frostegard, A., Tunlid, A., Baath, E.: Use and misuse of PLFA measurements in soils, *Soil Biol. Biochem.*, 43, 1621-1625, doi:10.1016/j.soilbio.2010.11.021, 2011.
- [19] Gearing, P. J., Gearing, J. N., Maughan, J. T., Oviatt, C. A.: Isotopic Distribution of Carbon from Sewage Sludge and Eutrophication in the Sediments and Food Web of Estuarine Ecosystems, *Environ. Sci. Technol.*, 25, 295-301, 1991.
- [20] Glanville, H., Rousk, J., Golyshin, P., Jones, D. L.: Mineralization of low molecular weight carbon substrates in soil solution under laboratory and field conditions, *Soil Biol. Biochem.*, 48, 88-95, doi:10.1016/j.soilbio.2012.01.015, 2012.
- [21] Glaser: Compound-specific stable-isotope (<sup>d</sup><sup>13</sup>C) analysis in soil science, 2005.
- [22] Glaser, B. and Amelung, W.: Determination of <sup>13</sup>C natural abundance of amino acid enantiomers in soil: methodological considerations and first results, *Rapid Commun. Mass Sp.*, 16, 891-898, doi:10.1002/rcm.650, 2002.
- [23] Glaser, B. and Gross, S.: Compound-specific-<sup>13</sup>C analysis of individual amino sugars-a tool to quantify timing and amount of soil microbial residue stabilization, *Rapid Commun. Mass Sp.*, 19, 1409-1416, doi:10.1002/rcm.1913, 2005.
- [24] Glaser, B., Turrión, M. B., Alef, K.: Amino sugars and muramic acid-biomarkers for soil microbial community structure analysis, *Soil Biol. Biochem.*, 36, 399-407, doi:10.1016/j.soilbio.2003.10.013, 2004.
- [25] Glaser, B., Millar, N., Blum, H.: Sequestration and turnover of bacterial-and fungal-derived carbon in a temperate grassland soil under long-term elevated atmospheric pCO<sub>2</sub>, *Glob. Change Biol.*, 12, 1521-1531, doi:10.1111/j.1365-2486.2006.01186.x, 2006.
- [26] Gottschalk, G.: *Bacterial Metabolism*, Springer-Verlag New York, New York, 1979.
- [27] Gregorich, E. G., Voroney, R. P., Kachanoski, R. G.: Turnover of carbon through the microbial biomass in soil with different textures, *Soil Biol. Biochem.*, 23, 799-805, doi:10.1016/0038-0717(91)90152-a, 1991.
- [28] Gunina, A. and Kuzyakov, Y.: Sugars in soil and sweets for microorganisms: Review of origin, content, composition and fate, *Soil Biol. Biochem.*, 90, 87-100, doi:10.1016/j.soilbio.2015.07.021, 2015.

- [29] Gunina, A., Dippold, M., Glaser, B., Kuzyakov, Y.: Fate of low molecular weight organic substances in an arable soil: From microbial uptake to utilisation and stabilisation, *Soil Biol. Biochem.*, 77, 304-313, 2014.
- [30] Indorf, C., Stamm, F., Dyckmans, J., Joergensen, R. G.: Determination of saprotrophic fungi turnover in different substrates by glucosamine-specific delta C-13 liquid chromatography/isotope ratio mass spectrometry, *Fungal Ecol.*, 5, 694-701, doi:10.1016/j.funeco.2012.07.001, 2012.
- [31] IUSS Working Group WRB: World Reference Base for Soil Resources 2014, update 2015 International soil classification system for naming soils and creating legends for soil maps, *World Soil Resources Reports No. 106*, FAO, Rome, 2015.
- [32] Jenkinson, D. S. and Ladd, J. N.: Microbial biomass in soil: measurement and turnover, *Soil Biochemistry*, Marcel Dekker, New York, 415-471, 1981.
- [33] Kieft, T. L., Wilch, E., O'Connor, K., Ringelberg, D. B., and White, D. C.: Survival and phospholipid fatty acid profiles of surface and subsurface bacteria in natural sediment microcosms, *Appl. Environ. Microbiol.*, 63, 1531-1542, 1997.
- [34] Kindler, R., Miltner, A., Richnow, H.-H., Kaestner, M.: Fate of gram-negative bacterial biomass in soil-mineralization and contribution to SOM, *Soil Biol. Biochem.*, 38, 2860-2870, doi:10.1016/j.soilbio.2006.04.047, 2006.
- [35] Kindler, R., Miltner, A., Thullner, M., Richnow, H.-H., Kaestner, M.: Fate of bacterial biomass derived fatty acids in soil and their contribution to soil organic matter, *Org. Geochem.*, 40, 29-37, doi:10.1016/j.orggeochem.2008.09.005, 2009.
- [36] Knapp, D.: *Handbook of Analytical Derivatization Reagents*, Wiley, New York, 1979.
- [37] Kouno, K., Wu, J., Brookes, P. C.: Turnover of biomass C and P in soil following incorporation of glucose or ryegrass, *Soil Biol. Biochem.*, 34, 617-622, doi:10.1016/s0038-0717(01)00218-8, 2002.
- [38] Kuzyakov, Y.: How to link soil C pools with CO<sub>2</sub> fluxes, *Biogeosciences*, 8, 1523-1537, doi:10.5194/bg-8-15-3-2011, 2011.
- [39] Lengeler, J. W., Drews, G., Schlegel, H. G.: *Biology of the prokaryotes*, Georg Thieme Verlag, 955, 110-163, 1999.
- [40] Liang, C., Fujinuma, R., Balser, T. C.: Comparing PLFA and amino sugars for microbial analysis in an Upper Michigan old growth forest, *Soil Biol. Biochem.*, 40, 2063-2065, doi:10.1016/j.soilbio.2008.01.022, 2008.
- [41] Lutzow, M. v., Koegel-Knabner, I., Ekschmitt, K., Matzner, E., Guggenberger, G., Marschner, B., Flessa, H.: Stabilization of organic matter in temperate soils: mechanisms and their relevance under different soil conditions-a review, *Eur. J. Soil Sci.*, 57, 426-445, doi:10.1111/j.1365-2389.2006.00809.x, 2006.
- [42] Malik, A., Blagodatskaya, E., Gleixner, G.: Soil microbial carbon turnover decreases with increasing molecular size, *Soil Biol. Biochem.*, 62, 115-118, doi:10.1016/j.soilbio.2013.02.022, 2013.
- [43] Malik, A., Dannert, H., Griffiths, R. I., Thomson, B. C., Gleixner, G.: Rhizosphere bacterial carbon turnover is higher in nucleic acids than membrane lipids: implications for understanding soil carbon cycling, *Front. Microbiol.*, 6, 268, doi:10.3389/fmicb.2015.00268, 2015.
- [44] Moore, J. C., McCann, K., Ruiter, P. C. de: Modeling trophic pathways, nutrient cycling, and dynamic stability in soils, *Pedobiologia*, 49, 499-510, doi:10.1016/j.pedobi.2005.05.008, 2005.
- [45] Park, J. T. and Uehara, T.: How bacteria consume their own exoskeletons (Turnover and recycling of cell wall peptidoglycan), *Microbiol. Mol. Biol. R.*, 72, 211-227, doi:10.1128/mmbr.00027-07, 2008.
- [46] Parton, W. J., Schimel, D. S., Cole, C. V., Ojima, D. S.: Analysis of factors controlling soil organic matter levels in Great Plains grasslands, *Soil Sci. Soc. Am. J.*, 51, 1173-1179, 1987.
- [47] Perelo, L. W. and Munch, J. C.: Microbial immobilisation and turnover of <sup>13</sup>C labelled substrates in two arable soils under field and laboratory conditions, *Soil Biol. Biochem.*, 37, 2263-2272, doi:10.1016/j.soilbio.2005.02.039, 2005.
- [48] Reischke, S., Rousk, J., Baåth, E.: The effects of glucose loading rates on bacterial and fungal growth in soil, *Soil Biol. Biochem.*, 70, 88-95, 2014.
- [49] Rethemeyer, J.: Age heterogeneity of soil organic matter, 2004.
- [50] Rethemeyer, J., Kramer, C., Gleixner, G., John, B., Yamashita, T., Flessa, H., Andersen, N., Nadeau, M. J., Grootes, P. M.: Transformation of organic matter in agricultural soils: radiocarbon concentration versus soil depth, *Geoderma*, 128, 94-105, doi:10.1016/j.geoderma.2004.12.017, 2005.
- [51] Rousk, J. and Baath, E.: Fungal biomass production and turnover in soil estimated using the acetate-in ergosterol technique, *Soil Biol. Biochem.*, 39, 2173-2177, doi:10.1016/j.soilbio.2007.03.023, 2007.
- [52] Ruess, L., Schutz, K., Haubert, D., Haggblom, M. M., Kandeler, E., Scheu, S.: Application of lipid analysis to understand trophic interactions in soil, *Ecology*, 86, 2075-2082, doi:10.1890/04-1399, 2005.
- [53] Schimel, J. P. and Weintraub, M. N.: The implications of exoenzyme activity on microbial carbon and nitrogen limitation in soil: a theoretical model, *Soil Biol. Biochem.*, 35, 549-563, 2003.

- [54] Schmidt, S. K., Costello, E. K., Nemergut, D. R., Cleveland, C. C., Reed, S. C., Weintraub, M. N., Meyer, A. F., Martin, A. M.: Biogeochemical consequences of rapid microbial turnover and seasonal succession in soil, *Ecology*, 88, 1379-1385, doi:10.1890/06-0164, 2007.
- [55] Schneckenberger, K., Demin, D., Stahr, K., Kuzyakov, Y.: Microbial utilization and mineralization of <sup>14</sup>C glucose added in six orders of concentration to soil, *Soil Biol. Biochem.*, 40, 1981-1988, doi:10.1016/j.soilbio.2008.02.020, 2008.
- [56] Wagner, G. H.: Significance of microbial tissue to soil organic matter. In: *Isotopes and radiation in soil organic matter studies*, FAO/IAEA, Vienna, 197-205, 1968.
- [57] Wallander, H., Ekblad, A., Godbold, D. L., Johnson, D., Bahr, A., Baldrian, P., Bjork, R. G., Kieliszewska-Rokicka, B., Kjoller, R., Kraigher, H., Plassard, C., Rudawska, M.: Evaluation of methods to estimate production, biomass and turnover of ectomycorrhizal mycelium in forests soils-A review, *Soil Biol. Biochem.*, 57, 1034-1047, doi:10.1016/j.soilbio.2012.08.027, 2013.
- [58] Waring, B. G., Averill, C., Hawkes, C. V., Holyoak, M.: Differences in fungal and bacterial physiology alter soil carbon and ni-trogen cycling: Insights from meta-analysis and theoretical models, *Ecol. Lett.*, 16, 887-894, doi:10.1111/ele.12125, 2013.
- [59] Wu, J., Joergensen, R. G., Pommerening, B., Chaussod, R., Brookes, P. C.: Measurement of soil microbial biomass C by fumigation extraction-an automated procedure, *Soil Biol. Biochem.*, 22, 1167-1169, doi:10.1016/0038-0717(90)90046-3, 1990.
- [60] Wu, J., Brookes, P. C., Jenkinson, D. S.: Formation and destruction of microbial biomass during the decomposition of glucose and ryegrass in soil, *Soil Biol. Biochem.*, 25, 1435-1441, doi:10.1016/0038-0717(93)90058-j, 1993.
- [61] Zelles, L.: Phospholipid fatty acid profiles in selected members of soil microbial communities, *Chemosphere*, 35, 275-294, doi:10.1016/s0045-6535(97)00155-0, 1997.
- [62] Zelles, L., Palojarvi, A., Kandeler, E., VonLutzow, M., Winter, K., Bai, Q. Y.: Changes in soil microbial properties and phospholipid fatty acid fractions after chloroform fumigation, *Soil Biol. Biochem.*, 29, 1325-1336, doi:10.1016/s0038-0717(97)00062-x, 1997.
- [63] Zhang, H., Ding, W., Yu, H., He, X.: Carbon uptake by a microbial community during 30-day treatment with C-13-glucose of a sandy loam soil fertilized for 20 years with NPK or compost as determined by a GC-C-IRMS analysis of phospholipid fatty acids, *Soil Biol. Biochem.*, 57, 228-236, doi:10.1016/j.soilbio.2012.08.024, 2013.
- [64] Zhang, X. D. and Amelung, W.: Gas chromatographic determination of muramic acid, glucosamine, mannosamine, and galactosamine in soils, *Soil Biol. Biochem.*, 28, 1201-1206, doi:10.1016/0038-0717(96)00117-4, 1996.
- [65] Ziegler, S. E., White, P. M., Wolf, D. C., Thoma, G. J.: Tracking the fate and recycling of C-13-labeled glucose in soil, *Soil Sci.*, 170, 767-778, doi:10.1097/01.ss.0000190511.64169.9c, 2005.