

Effect of hydrostatic pressure, temperature, and solvent on the rate of the Diels-Alder reaction between 9,10-anthracenedimethanol and maleic anhydride

Kiselev V., Kornilov D., Anikin O., Latypova L., Konovalov A.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2017, Pleiades Publishing, Ltd. The rate of the reaction between 9,10-anthracenedimethanol and maleic anhydride in 1,4-dioxane, acetonitrile, trichloromethane, and toluene is studied at 25, 35, 45°C in the pressure range of 1–1772 bar. The rate constants, enthalpies, entropies and activation volumes are determined. It is shown that the rate of reaction with 9,10-anthracenedimethanol is approximately one order of magnitude higher than with 9-anthracenemethanol.

<http://dx.doi.org/10.1134/S0036024417030128>

Keywords

9,10-dimethanol anthracene, Diels-Alder reaction, effect of pressure, effect of temperature, solvent effect

References

- [1] D. C. Rideout and R. Breslow, *J. Am. Chem. Soc.* 102, 7816 (1980).
- [2] R. Breslow, U. Maitra, and D. Rideout, *Tetrahedron Lett.* 24, 1901 (1983).
- [3] R. Breslow and U. Maitra, *Tetrahedron Lett.* 25, 1239 (1984).
- [4] R. Breslow and T. Guo, *J. Am. Chem. Soc.* 110, 5613 (1988).
- [5] J. Light and R. Breslow, *Tetrahedron Lett.* 31, 2957 (1990).
- [6] R. Breslow, *Acc. Chem. Res.* 24, 159 (1991).
- [7] W. Blokzijl, M. J. Blandamer, and J. B. F. N. Engberts, *J. Am. Chem. Soc.* 113, 4241 (1991).
- [8] W. Blokzijl and J. B. F. N. Engberts, *J. Am. Chem. Soc.* 114, 5440 (1992).
- [9] S. Horiuchi, T. Murase, and M. Fujita, *Asian J. Chem.* 6, 1839 (2011).
- [10] C.-J. Li, *Chem. Rev.* 93, 2023 (1993).
- [11] X. Chen, J. F. Dobson, and C. L. Raston, *Chem. Commun.* 48, 3703 (2012).
- [12] L. Yasmin, K. A. Stubbs, and C. L. Raston, *Tetrahedron Lett.* 55, 2246 (2014).
- [13] S. Narayan, J. Muldoon, M. G. Finn, et al., *Ang. Chem. Int. Ed.* 44, 3275 (2005).
- [14] J. E. Klijn and J. B. F. N. Engberts, *Nature* 435, 746 (2005).
- [15] Y. Jung and R. A. Marcus, *J. Am. Chem. Soc.* 127, 5492 (2007).
- [16] G. Desimoni, G. Faita, P. P. Righetti, and L. Toma, *Tetrahedron* 46, 7951 (1990).
- [17] V. D. Kiselev, I. I. Shakirova, and A. I. Konovalov, *Russ. Chem. Bull.* 62, 285 (2013).
- [18] V. D. Kiselev and A. I. Konovalov, *J. Phys. Org. Chem.* 22, 466 (2009).

- [19] V. D. Kiselev and J. G. Miller, *J. Am. Chem. Soc.* 97, 4036 (1975).
- [20] V. D. Kiselev, *Int. J. Chem. Kinet.* 45, 613 (2013).
- [21] V. D. Kiselev, D. A. Kornilov, I. I. Lekomtseva, and A. I. Konovalov, *Int. J. Chem. Kinet.* 47, 289 (2015).
- [22] V. D. Kiselev, E. A. Kashaeva, L. N. Potapova, et al., *Izv. Akad. Nauk, Ser. Khim.*, No. 4360 (2016).
- [23] J. Sauer, D. Lang, and A. Mielert, *Angew. Chem.* 74, 352 (1962).
- [24] M. W. Miller, R. W. Amidon, and P. O. Tawney, *J. Am. Chem. Soc.* 77, 2845 (1955).
- [25] M. H. Schwartz, S. M. Rosenfeld, C. I. Lee, et al., *Tetrahedron Lett.* 33, 6275 (1992).
- [26] J. A. Riddick, W. B. Bunger, and T. K. Sakano, *Organic Solvents*, 4th ed. (Wiley, New York, Chichester, Brisbane, Toronto, Singapore, 1986).
- [27] V. D. Kiselev, A. V. Bolotov, A. P. Satonin, et al., *J. Phys. Chem. B* 112, 6674 (2008).
- [28] V. D. Kiselev, I. I. Shakirova, D. A. Kornilov, E. A. Kashaeva, L. N. Potapova, and A. I. Konovalov, *Russ. J. Phys. Chem. A* 87, 160 (2013).
- [29] V. D. Kiselev, D. A. Kornilov, E. A. Kashaeva, L. N. Potapova, D. B. Krivolapov, I. A. Litvinov, and A. I. Konovalov, *Russ. J. Phys. Chem. A* 88, 2073 (2014).
- [30] D. A. Kornilov and V. D. Kiselev, *J. Chem. Eng. Data* 60, 3571 (2015).