Biochemistry (Moscow) Supplement Series B: Biomedical Chemistry 2016 vol.10 N1, pages 87-90

## The combined action of binase and bleomycin on human lung adenocarcinoma cells

Zelenikhin P., Makeeva A., Nguen T., Siraj Y., Ilinskaya O. Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

## Abstract

© 2016, Pleiades Publishing, Ltd.Some microbial ribonucleases (RNases) demonstrate selective cytotoxic effect against a wide range of tumor cells. In this context combined use of cytotoxic RNases in complex therapy with other chemotherapeutic agents appears to be especially promising. In this study we have investigated the apoptosisinduced effect of Bacillus pumilus RNase (binase) in combination with known antitumor antibiotic bleomycin on human lung adenocarcinoma A549 cells. The combined effect of high concentrations of these agents did not have any mutual increase in their apoptosisinduced action, while a combination of nonapoptotic concentrations resulted in the increase of proportion of apoptotic cells up to 22% as compared with individual effect of bleomycin (6%) and binase (12%) used separately. These results indicate that binase and bleomycin are effective in combination of their low concentrations and ineffective in combination of their high concentrations.

http://dx.doi.org/10.1134/S1990750816010121

## **Keywords**

antitumor activity, Bacillus pumilus, binase, bleomycin, cytotoxic ribonucleases, lung adenocarcinoma