

The integrated exploration of Raifa lake sediments and dendrochronological analysis of Raifa forestry pines

Kosareva L., Nurgalieva N., Frolova L., Gafiatullina L., Krylov P., Terekhin A., Tishin D., Batalin G., Gareev B., Kuzina D., Antonenko V., Akhmerov R.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2006-2017 Asian Research Publishing Network (ARPN). All rights reserved. The article presents some results of comprehensive research on the properties of Raifa lake bottom sediments and dendrochronological study of Raifa forestry pines (Russia). A preliminary seismic acoustic investigations have been carried out, which allowed to select the sampling site (E 48 ° 43'40.6" N 55 ° 54'21.7"). The length of core sample was 32 cm. Laboratory studies of the core, including the study of the elemental composition, magneto-mineralogical and carcinologic analysis revealed the features of sedimentation mass formation.

Keywords

Carcinologic analysis, Dendrochronological analysis, Elemental composition, Lacustrine sediments, Magnetic properties, Paleoclimate, Paleolimnology, Seismic acoustic studies

References

- [1] Unkovskaya E.N., V.V. Zharikov, S.V. Bykov, M.Yu. Gorbunov, M. Uman, N.G. Tarasova, O.V. Muhortova, O.V. Palagushkina, O.J. Derevenskaya. 2010. Plankton communities of Lake Raifa (Volga-Kama Natural Biosphere Reserve) I. Biodiversity plankton communities of different habitats of Lake Raifa. *Izvestia Samarskogo nauchnogo sentra, Russian Academy of Sciences*. 12(1) (5): 1453-1460
- [2] Bykova S.V., V.V. Zharikov. 2009. Ciliates of Lake Raifa (Volga-Kama biosphere reserve). *Samarskaya Luka: regional and global environmental problems*. 18(3): 121-131
- [3] Muhortova O.V., N.G. Tarasova, S.V. Bykov, T.N. Burkova, E.N. Unkovsky. 2011. Feature zooplankton phytophilic complex of Lakes Raifa and Dolgoe of Volga-Kama Natural Biosphere Reserve. *Ecology and geography of plants and plant communities of the Middle Volga Region*. pp. 346-352
- [4] Rysaeva I.A., Dvinskikh A.P. 2013. Features of macrocomponent composition of lake waters in the Raifsky area of the Volga Kama national natural biospheric reserve. "Vestnik AGU". 4: 88-94
- [5] Iassonov P.G., Nourgaliev D.K., Bourov B.V., Heller F. 1998. A modernized coercivity spectrometer/*Geologica Carpathica*. 49(N 3): 224-226
- [6] Kosareva L.R., Nourgaliev D.K., Kuzina D.M., Spassov S., Fattakhov A.V. 2015. Ferromagnetic, dia-/paramagnetic and superparamagnetic components of Aral sea sediments: significance for paleoenvironmental reconstruction. *ARPN Journal of Earth Sciences*. 4(1): 1-6
- [7] Kosareva L.R., Utemov E.V., Nurgaliev D.K., Shcherbakov V.P., Kosarev V.E., Yasonov P.G. 2015. Separation of Ferromagnetic Components by Analyzing the Hysteresis Loops of Remanent Magnetization *Izvestiya, Physics of the Solid Earth*. 51(5): 689-708
- [8] Sarmaja-Korjonen K., Szeroczyńska K. 2007. *Atlas of Subfossil Cladocera from Central and Northern Europe*. Friends of the Lower Vistula Society. p. 84

- [9] Korosi J. B., Smol John P. 2012. An illustrated guide to the identification of cladoceran subfossils from lake sediments in northeastern North America. *The Chydoridae*. *J. Paleolimnol.* 48: 571-586
- [10] Frolova L.A. 2013. Cladocera. Biological indicators in paleobiological studies: atlas/ Scientific Ed. L.B. Nazarova. Kazan: Kazan University Press. pp. 64-87
- [11] Kotov A.A., A.Y. Sinev, S.M. Glagolev, Smirnov N.N. 2010. Cladocerans (Cladocera). Determinant of zooplankton and zoobenthos in fresh waters of European Russia. V.1. Zooplankton/ ed. V.R. Alekseev, S.Y. Tsalolihin.-M.: Association of scientific editions KMK. pp. 151-276
- [12] Flössner D. 2000. Die Haplopoda und Cladocera (ohne Bosminidae) Mitteleuropas. Leiden: Backhuys Publishers. p. 428
- [13] Grimm EC. 1987 CONISS: a Fortran 77 program for stratigraphically constrained cluster analysis by the method of incremental sum of squares.-*ComputGeosci* 113-pp. 13-35
- [14] 1993. Grimm EC TILIA 2.0.b.4 (Computer software). Illinois StateMuseum, Research and Collections Center, Springfield
- [15] Shiyatov S.G. 1986. Dendrochronology upper forest border in the Urals. M.: Nauka. p. 137
- [16] Shiyatov S.G. Dendrochronology methods. Part I. 2000. Fundamentals of dendrochronology. Gathering and getting tree-ring data/ S.G. Shiyatov, E.A. Vaganov, A. Kirdyanov, V.B. Kruglov, V.S. Mazepa, M.M. Naurzbaev, R.M. Hantemirov.-Krasnoyarsk: Krasnoyarsk State University. p. 80
- [17] Rinn, F. 2005. TSAPWin-Time Series Analysis and Presentation for Dendrochronology and Related Applications, Version 0.53, User Reference/ F. Rinn.-Heidelberg. p. 91
- [18] Bunn, A.G. dplR: Dendrochronology Program Library in R. 2012 R package version 1.5.6/ A.G. Bunn, M. Korpela, F. Biondi, F. Campelo, P. Merian, F. Qeadan, Ch. Zang. URL: <http://CRAN.Rproject.org/package=dplR>
- [19] Hammer O. 2003. Past-PAlaeological STatistics, ver. 1.04, 59 p. (<http://www.folk.uio.no/ohammer/past>)
- [20] Anisimov M.A., Ivanova V.V., Pushina Z.V., Pitulko V.V. 2009. Lagoon sediments of Zhokhov Island: age, conditions of formation and significance for paleogeographical reconstruction of the New Siberian Islands. *Proceedings of the Russian Academy of Sciences. Series: Geography.* (5): 107-119
- [21] Shotyk W., Cheburkin A.K., Appleby P.G. et al. 1966. Two thousand years of atmospheric arsenic, antimony and lead deposition in an ombrotrophic bog profile, Jura Mountains, Switzerland. *Earth and Planetary Scien. Letter.* 145: 1-7
- [22] Li Yuan-Hui. 1991. Distribution patterns of the elements in the ocean: A synthesis. *Geochim. et. Cosmochim. Acta.* 55: 3223-3240
- [23] Leonova G.A. 2009. Geochemical role of plankton of continental reservoirs of Siberia in the concentration and biosedimentation of trace elements-Doctoral dissertation author's abstract
- [24] Miko S., Mesic S., Prohic E., Pen Z. 2003. Trace element distribution in surface sediments of lake Vrana and topsoil of Cres island, Croatia.-*Nat. Croat.* 12(2): 93-111
- [25] Gorbatov E.S. 2013. Features of lake sedimentation during the formation of a ternary complex Shatarsky Meshcherov. p. 26 (in Russian)
- [26] Pashkova G.V., Ivanova E.V., Aysueva T.S., Shchetnikov A.A., Markova U.N., Finkelstein A.L. 2015. X-ray fluorescence determination of bromine in the bottom sediments of lakes for paleoclimatic studies. *Analysis and control.* T.19, No 4. pp. 340-346
- [27] Egli R. 2004a. Characterization of individual rock magnetic components by analysis of remanence curves. 1. Unmixing natural sediments// *StudiaGeophysicaetGeodaetica.* 48(2): 391-446
- [28] Egli R. 2004b. Characterization of individual rock magnetic components by analysis of remanence curves. 2. Fundamental properties of coercivity distributions// *Physics and Chemistry of the Earth.* 29(13-14): 851-867
- [29] Frolova L. 2016. Subfossil Cladocera (Branchiopoda, Crustacea) in climatic and palaeoenvironmental investigations in Eastern Siberia (Russia)-International Multidisciplinary Scientific GeoConference, 2016.-Book 4. Energy and Clean Technologies, Conference Proceedings. II: 601-607
- [30] Frolova L.A., Nazarova L.B., Pestryakova L.A., Herzschuh U. 2013. Analysis of the effects of climate-dependent factors on the formation of zooplankton communities that inhabit Arctic lakes in the Anabar River basin.-*Contemporary Problems of Ecology.* 6(1): 1-11
- [31] Deneke R. 2003. Review of rotifers and crustaceans in highly acidic environments of pH values < 3. *Hydrobiologia.* 433: 167-172
- [32] Quade H.W. 1969. Cladoceran faunas associated with aquatic macrophytes in some lakes in northwestern Minnesota. *Ecology.* 50: 170-179
- [33] Williams J.B. 1983. A study of summer mortality factors for natural populations of Chydoridae (Cladocera). *Hydrobiologia.* 107: 131-139
- [34] Frolova L., Nazarova L., Pestryakova L., Herzschuh U. 2014. Subfossil cladoceran remains from sediment in thermokarst lakes in northeastern Siberia, Russia.-*Journal of Paleolimnology.* 52(1): 107-119

- [35] Frolova L., Ibragimova A., Fedorova I. 2016. Stratigraphy of Cladocera in a core from A Yamal Peninsula lake (Arctic Russia). International Multidisciplinary Scientific GeoConference.-Book 4.-Energy and Clean Technologies, Conference Proceedings. II: 579-587
- [36] www.nina.no
- [37] Nevalainen L., Luoto T. 2010. Temperature sensitivity of gamogenesis in littoral cladocerans and its ecological implications// J. Limnol. 69(1): 120-125
- [38] Shannon C.E., Weaver W. 1949. The mathematical theory of communication.-Urbana: Univ. Illinois Press. p. 117
- [39] Structures of zooplankton communities of lakes in Raifa part of Volzhsko-Kamaskiy Reserve and its buffer zone/ Derevenskaya, O.Yu., Unkovskaya, E.N., Mingazova, N.M., Pavlova, L.R. 2002. Transactions of Volzhsko-Kamskiy National Nature Reserve. Kazan. 5: 52-70
- [40] Bitvinskas T.T. 1974. Dendroclimatic research. L. Gidrometeoizdat. p. 172
- [41] Pugachev P.G. 1975. The dynamics of annual growth *Pinus sylvestris*. L. in Turgai basin due to climatic factors// Bot.zhurnal. 3: 401-412
- [42] Tishin D.V., Chizhikova N.A., Chugunov R.G. 2012. Assessing the impact of climatic factors on the radial growth of pine (*Pinus sylvestris* L) sphagnum bogs of the Volga-Kama Reserve. 2012,-Book of abstracts "Dendro 2012: perspectives of using tree-ring information for protection, restoration and sustainable use of tree vegetation. Materials of Conference. The Publishing House of the Moscow State University of Forest". (in Russian);
- [43] Rigling A., Waldner P., Forster T. et al. 2001. Ecological interpretation of tree-ring width and IADF in *Pinussylvestris* on dry sites in the central Alps and Siberia. Can. J. For. Res. 31: 18-31