

Frontiers in Immunology, 2017, vol.8, NMA5

---

# Serum cytokine profiles differentiating hemorrhagic fever with renal syndrome and hantavirus pulmonary syndrome

Khaiboullina S., Levis S., Morzunov S., Martynova E., Anokhin V., Gusev O., St Jeor S., Lombardi V., Rizvanov A.

*Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia*

---

## Abstract

© 2017 Khaiboullina, Levis, Morzunov, Martynova, Anokhin, Gusev, St Jeor, Lombardi and Rizvanov. Hantavirus infection is an acute zoonosis that clinically manifests in two primary forms, hemorrhagic fever with renal syndrome (HFRS) and hantavirus pulmonary syndrome (HPS). HFRS is endemic in Europe and Russia, where the mild form of the disease is prevalent in the Tatarstan region. HPS is endemic in Argentina, as well as other countries of North and South American. HFRS and HPS are usually acquired via the upper respiratory tract by inhalation of virus-contaminated aerosol. Although the pathogenesis of HFRS and HPS remains largely unknown, postmortem tissue studies have identified endothelial cells as the primary target of infection. Importantly, cell damage due to virus replication, or subsequent tissue repair, has not been documented. Since no single factor has been identified that explains the complexity of HFRS or HPS pathogenesis, it has been suggested that a cytokine storm may play a crucial role in the manifestation of both diseases. In order to identify potential serological markers that distinguish HFRS and HPS, serum samples collected during early and late phases of the disease were analyzed for 48 analytes using multiplex magnetic bead-based assays. Overall, serum cytokine profiles associated with HPS revealed a more pro-inflammatory milieu as compared to HFRS. Furthermore, HPS was strictly characterized by the upregulation of cytokine levels, in contrast to HFRS where cases were distinguished by a dichotomy in serum cytokine levels. The severe form of hantavirus zoonosis, HPS, was characterized by the upregulation of a higher number of cytokines than HFRS (40 vs 21). In general, our analysis indicates that, although HPS and HFRS share many characteristic features, there are distinct cytokine profiles for these diseases. These profiles suggest a strong activation of an innate immune and inflammatory responses are associated with HPS, relative to HFRS, as well as a robust activation of Th1-type immune responses. Finally, the results of our analysis suggest that serum cytokines profiles of HPS and HFRS cases are consistent with the presence of extracellular matrix degradation, increased mononuclear leukocyte proliferation, and transendothelial migration.

<http://dx.doi.org/10.3389/fimmu.2017.00567>

---

## Keywords

Blood serum, Cytokine profile, Hantavirus pulmonary syndrome, Hemorrhagic fever with renal syndrome, Nephropathia epidemica

## References

- [1] Jonsson CB, Figueiredo LT, Vapalahti O. A global perspective on hantavirus ecology, epidemiology, and disease. *Clin Microbiol Rev* (2010) 23(2):412-41. doi: 10.1128/CMR.00062-09
- [2] Tappero JW, Khan AS, Pinner RW, Wenger JD, Graber JM, Armstrong LR, et al. Utility of emergency, telephone-based national surveillance for hantavirus pulmonary syndrome. Hantavirus task force. *JAMA* (1996) 275(5):398-400. doi:10.1001/jama.275.5.398
- [3] Calderón G, Pini N, Bolpe J, Levis S, Mills J, Segura E, et al. Hantavirus reservoir hosts associated with peridomestic habitats in Argentina. *Emerg Infect Dis* (1999) 5(6):792-7. doi:10.3201/eid0506.990608
- [4] Heyman P, Vaheri A, ENIVD Members. Situation of hantavirus infections and haemorrhagic fever with renal syndrome in European countries as of December 2006. *Euro Surveill* (2008) 13(28):18925
- [5] Tkachenko EA, Bernshtein AD, Dzagurova TK, Morozov VG, Slonova RA, Ivanov LI, et al. [Actual problems of hemorrhagic fever with renal syndrome]. *Zh Mikrobiol Epidemiol Immunobiol* (2013) 1:51-8
- [6] Hjertqvist M, Klein SL, Ahlm C, Klingstrom J. Mortality rate patterns for hemorrhagic fever with renal syndrome caused by Puumala virus. *Emerg Infect Dis* (2010) 16(10):1584-6. doi:10.3201/eid1610.100242
- [7] Makary P, Kanerva M, Ollgren J, Virtanen MJ, Vapalahti O, Lyytikäinen O. Disease burden of Puumala virus infections, 1995-2008. *Epidemiol Infect* (2010) 138(10):1484-92. doi:10.1017/S0950268810000087
- [8] Garanina SB, Platonov AE, Zhuravlev VI, Murashkina AN, Yakimenko VV, Korneev AG, et al. Genetic diversity and geographic distribution of hantaviruses in Russia. *Zoonoses Public Health* (2009) 56(6-7):297-309. doi:10.1111/j.1863-2378.2008.01210.x
- [9] KabweN E, DavidyukS Y, Morzunov P, ShakirovaV VG, AnokhinG A, IsaevaR S, et al. Genetic characterization of small (s)-segment genome Puumala virus strain Kazan. *BioNanoScience* (2016). doi:10.1007/s12668-016-03-8-2
- [10] Niklasson BS. Haemorrhagic fever with renal syndrome, virological and epidemiological aspects. *Pediatr Nephrol* (1992) 6(2):201-4. doi:10.1007/BF00866318
- [11] Martinez-Valdebenito C, Calvo M, Vial C, Mansilla R, Marco C, Palma RE, et al. Person-to-person household and nosocomial transmission of Andes hantavirus, Southern Chile, 2011. *Emerg Infect Dis* (2014) 20(10):1629-36. doi:10.3201/eid2010.140353
- [12] Padula PJ, Edelstein A, Miguel SD, López NM, Rossi CM, Rabinovich RD. [Epidemic outbreak of Hantavirus pulmonary syndrome in Argentina. Molecular evidence of person to person transmission of Andes virus]. *Medicina (B Aires)* (1998) 58(Suppl 1):27-36
- [13] Enria DA, Briggiler AM, Pini N, Levis S. Clinical manifestations of New World hantaviruses. *Curr Top Microbiol Immunol* (2001) 256:117-34. doi:10.1007/978-3-642-56753-7\_7
- [14] Pini N. Hantavirus pulmonary syndrome in Latin America. *Curr Opin Infect Dis* (2004) 17(5):427-31. doi:10.1097/00001432-200410000-00007
- [15] Figueiredo LTM, Moreli ML, de-Sousa RLM, Borges AA, de-Figueiredo GG, Machado AM, et al. Hantavirus Pulmonary Syndrome, Central Plateau, Southeastern, and Southern Brazil. *Emerg Infect Dis* (2009) 15(4):561-7. doi:10.3201/eid1504.080289
- [16] Settergren B, Leschinskaya E, Zagidullin I, Fazlyeva R, Khunafina D, Niklasson B. Hemorrhagic fever with renal syndrome: comparison of clinical course in Sweden and in the Western Soviet Union. *Scand J Infect Dis* (1991) 23(5):549-52. doi:10.3109/00365549109105176
- [17] Peco-Antic A, Popovic-Rolovic M, Gligic A, Popovic D, Jovanovic O, Kostic M. Clinical characteristics of haemorrhagic fever with renal syndrome in children. *Pediatr Nephrol* (1992) 6(4):335-8. doi:10.1007/BF00869727
- [18] Bren AF, Pavlovcic SK, Koselj M, Kovac J, Kandus A, Kveder R. Acute renal failure due to hemorrhagic fever with renal syndrome. *Ren Fail* (1996) 18(4):635-8. doi:10.3109/08860229609047688
- [19] Settergren B. Nephropathia epidemica (hemorrhagic fever with renal syndrome) in Scandinavia. *Rev Infect Dis* (1991) 13(4):736-44. doi:10.1093/clinids/13.4.736
- [20] Mäkelä S, Ala-Houhala I, Mustonen J, Koivisto AM, Kouri T, Turjanmaa V, et al. Renal function and blood pressure five years after Puumala virus-induced nephropathy. *Kidney Int* (2000) 58(4):1711-8. doi:10.1046/j.1523-1755.2000.00332.x
- [21] Khaiboullina SF, Netski DM, Krumpel P, St Jeor SC. Effects of tumor necrosis factor alpha on sin nombre virus infection in vitro. *J Virol* (2000) 74(24):11966-71. doi:10.1128/JVI.74.24.11966-11971.2000
- [22] Temonen M, Vapalahti O, Holthöfer H, Brummer-Korvenkontio M, Vaheri A, Lankinen H. Susceptibility of human cells to Puumala virus infection. *J Gen Virol* (1993) 74(Pt 3):515-8. doi:10.1099/0022-1317-74-3-515
- [23] Zaki SR, Greer PW, Coffield LM, Goldsmith CS, Nolte KB, Foucar K, et al. Hantavirus pulmonary syndrome. Pathogenesis of an emerging infectious disease. *Am J Pathol* (1995) 146(3):552-79

- [24] Temonen M, Mustonen J, Helin H, Pasternack A, Vaheri A, Holthöfer H. Cytokines, adhesion molecules, and cellular infiltration in nephropathia epidemica kidneys: an immunohistochemical study. *Clin Immunol Immunopathol* (1996) 78(1):47-55. doi:10.1006/clin.1996.0007
- [25] Colby TV, Zaki SR, Feddersen RM, Nolte KB. Hantavirus pulmonary syndrome is distinguishable from acute interstitial pneumonia. *Arch Pathol Lab Med* (2000) 124(10):1463-6. doi:10.1043/0003-9985(2000)124<1463:HPSIDF>2.0.CO;2
- [26] Mori M, Rothman AL, Kurane I, Montoya JM, Nolte KB, Norman JE, et al. High levels of cytokine-producing cells in the lung tissues of patients with fatal hantavirus pulmonary syndrome. *J Infect Dis* (1999) 179(2):295-302. doi:10.1086/314597
- [27] Sadeghi M, Eckerle I, Daniel V, Burkhardt U, Opelz G, Schnitzler P. Cytokine expression during early and late phase of acute Puumala hantavirus infection. *BMC Immunol* (2011) 12:65. doi:10.1186/1471-2172-12-65
- [28] Khaiboullina SF, Martynova EV, Khamidullina ZL, Lapteva EV, Nikolaeva IV, Anokhin VV, et al. Upregulation of IFN-gamma and IL-12 is associated with a milder form of hantavirus hemorrhagic fever with renal syndrome. *Eur J Clin Microbiol Infect Dis* (2014) 33(12):2149-56. doi:10.1007/s10096-014-2176-x
- [29] Borges AA, Campos GM, Moreli ML, Moro Souza RL, Saggiaro FP, Figueiredo GG, et al. Role of mixed Th1 and Th2 serum cytokines on pathogenesis and prognosis of hantavirus pulmonary syndrome. *Microbes Infect* (2008) 10(10-11):1150-7. doi:10.1016/j.micinf.2008.06.006
- [30] Saksida A, Wraber B, Avsic-Zupanc T. Serum levels of inflammatory and regulatory cytokines in patients with hemorrhagic fever with renal syndrome. *BMC Infect Dis* (2011) 11:142. doi:10.1186/1471-2334-11-142
- [31] Zupanc TA, Korva M, Markotic A. HFRS and hantaviruses in the Balkans/South-East Europe. *Virus Res* (2014) 187:27-33. doi:10.1016/j.virusres.2013.12.042
- [32] Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* (2009) 4(1):44-57. doi:10.1038/nprot.2008.211
- [33] Santiago B, Baleux F, Palao G, Gutiérrez-Cañas I, Ramírez JC, Arenzana-Seisdedos F, et al. CXCL12 is displayed by rheumatoid endothelial cells through its basic amino-terminal motif on heparan sulfate proteoglycans. *Arthritis Res Ther* (2006) 8(2):R43. doi:10.1186/ar1900
- [34] Ding L, Morrison SJ. Haematopoietic stem cells and early lymphoid progenitors occupy distinct bone marrow niches. *Nature* (2013) 495(7440):231-5. doi:10.1038/nature11885
- [35] Gear ARL, Camerini D. Platelet chemokines and chemokine receptors: linking hemostasis, inflammation, and host defense. *Microcirculation* (2003) 10(3-4):335-50. doi:10.1038/sj.mn.7800198
- [36] Schober A, Karshovska E, Zerneck A, Weber C. SDF-1alpha-mediated tissue repair by stem cells: a promising tool in cardiovascular medicine? *Trends Cardiovasc Med* (2006) 16(4):103-8. doi:10.1016/j.tcm.2006.01.006
- [37] Abi-Younes S, Si-Tahar M, Luster AD. The CC chemokines MDC and TARC induce platelet activation via CCR4. *Thromb Res* (2001) 101(4):279-89. doi:10.1016/S0049-3848(00)00402-3
- [38] Chatterjee M, Gawaz M. Platelet-derived CXCL12 (SDF-1alpha): basic mechanisms and clinical implications. *J Thromb Haemost* (2013) 11(11):1954-67. doi:10.1111/jth.12404
- [39] Khaiboullina SF, Morzunov SP, St Jeor SC. Hantaviruses: molecular biology, evolution and pathogenesis. *Curr Mol Med* (2005) 5(8):773-90. doi:10.2174/156652405774962317
- [40] Clement J, Maes P, Lagrou K, Van Ranst M, Lameire N. A unifying hypothesis and a single name for a complex globally emerging infection: hantavirus disease. *Eur J Clin Microbiol Infect Dis* (2012) 31(1):1-5. doi:10.1007/s10096-011-1456-y
- [41] Gizzi M, Delaere B, Weynand B, Clement J, Maes P, Vergote V, et al. Another case of 'European hantavirus pulmonary syndrome' with severe lung, prior to kidney, involvement, and diagnosed by viral inclusions in lung macrophages. *Eur J Clin Microbiol Infect Dis* (2013) 32(10):1341-5. doi:10.1007/s10096-013-1885-x
- [42] Clement J, Maes P, Van Ranst M. Hemorrhagic fever with renal syndrome in the new, and hantavirus pulmonary syndrome in the Old World: paradisiacal lost or regained? *Virus Res* (2014) 187:55-8. doi:10.1016/j.virusres.2013.12.036
- [43] da Rosa Elkhoury M, da Silva Mendes W, Waldman EA, Dias JP, Carmo EH, Fernando da Costa Vasconcelos P. Hantavirus pulmonary syndrome: prognostic factors for death in reported cases in Brazil. *Trans R Soc Trop Med Hyg* (2012) 106(5):298-302. doi:10.1016/j.trstmh.2012.01.002
- [44] Cruikshank W, Little F. Interleukin-16: the ins and outs of regulating T-cell activation. *Crit Rev Immunol* (2008) 28(6):467-83. doi:10.1615/CritRevImmunol.v28.i6.10
- [45] Sánchez-Martín L, Estechea A, Samaniego R, Sánchez-Ramón S, Vega Má, Sánchez-Mateos P. The chemokine CXCL12 regulates monocyte-macrophage differentiation and RUNX3 expression. *Blood* (2011) 117(1):88-97. doi:10.1182/blood-2009-12-258186
- [46] D'Apuzzo M, Rolink A, Loetscher M, Hoxie JA, Clark-Lewis I, Melchers F, et al. The chemokine SDF-1, stromal cell-derived factor 1, attracts early stage B cell precursors via the chemokine receptor CXCR4. *Eur J Immunol* (1997) 27(7):1788-93. doi:10.1002/eji.1830270729

- [47] Roth SJ, Carr MW, Springer TA. C-C chemokines, but not the C-X-C chemokines interleukin-8 and interferon-gamma inducible protein-10, stimulate transendothelial chemotaxis of T lymphocytes. *Eur J Immunol* (1995) 25(12):3482-8. doi:10.1002/eji.1830251241
- [48] Proost P, Wuyts A, Van Damme J. Human monocyte chemotactic proteins-2 and-3: structural and functional comparison with MCP-1. *J Leukoc Biol* (1996) 59(1):67-74
- [49] Chintakuntlawar AV, Chodosh J. Chemokine CXCL1/KC and its receptor CXCR2 are responsible for neutrophil chemotaxis in adenoviral keratitis. *J Interferon Cytokine Res* (2009) 29(10):657-66. doi:10.1089/jir.2009.0006
- [50] De Filippo K, Dudeck A, Hasenberg M, Nye E, van Rooijen N, Hartmann K, et al. Mast cell and macrophage chemokines CXCL1/CXCL2 control the early stage of neutrophil recruitment during tissue inflammation. *Blood* (2013) 121(24):4930-7. doi:10.1182/blood-2013-02-486217
- [51] Tominaga K, Yoshimoto T, Torigoe K, Kurimoto M, Matsui K, Hada T, et al. IL-12 synergizes with IL-18 or IL-1beta for IFN-gamma production from human T cells. *Int Immunol* (2000) 12(2):151-60. doi:10.1093/intimm/12.2.151
- [52] Li F, Zhu H, Sun R, Wei H, Tian Z. Natural killer cells are involved in acute lung immune injury caused by respiratory syncytial virus infection. *J Virol* (2012) 86(4):2251-8. doi:10.1128/JVI.06209-11
- [53] Sitkovsky MV, Lukashev D, Apasov S, Kojima H, Koshiba M, Caldwell C, et al. Physiological control of immune response and inflammatory tissue damage by hypoxia-inducible factors and adenosine A2A receptors. *Annu Rev Immunol* (2004) 22:657-82. doi:10.1146/annurev.immunol.22.012703.104731
- [54] Prescott J, Safronetz D, Haddock E, Robertson S, Scott D, Feldmann H. The adaptive immune response does not influence hantavirus disease or persistence in the Syrian hamster. *Immunology* (2013) 140(2):168-78. doi:10.1111/imm.12116
- [55] Koivula TT, Tuulasvaara A, Hetemäki I, Mäkelä SM, Mustonen J, Sironen T, et al. Regulatory T cell response correlates with the severity of human hantavirus infection. *J Infect* (2014) 68(4):387-94. doi:10.1016/j.jinf.2013.11.007
- [56] Wang M, Wang J, Zhu Y, Xu Z, Yang K, Yang A, et al. Cellular immune response to Hantaan virus nucleocapsid protein in the acute phase of hemorrhagic fever with renal syndrome: correlation with disease severity. *J Infect Dis* (2009) 199(2):188-95. doi:10.1086/595834
- [57] Rasmuson J, Pourazar J, Linderholm M, Sandström T, Blomberg A, Ahlm C. Presence of activated airway T lymphocytes in human Puumala hantavirus disease. *Chest* (2011) 140(3):715-22. doi:10.1378/chest.10-2791
- [58] Björkström NK, Lindgren T, Stoltz M, Fauriat C, Braun M, Evander M, et al. Rapid expansion and long-term persistence of elevated NK cell numbers in humans infected with hantavirus. *J Exp Med* (2011) 208(1):13-21. doi:10.1084/jem.20100762
- [59] Onodera S, Kaneda K, Mizue Y, Koyama Y, Fujinaga M, Nishihira J. Macrophage migration inhibitory factor up-regulates expression of matrix metalloproteinases in synovial fibroblasts of rheumatoid arthritis. *J Biol Chem* (2000) 275(1):444-50. doi:10.1074/jbc.275.1.444
- [60] Kong YZ, Yu X, Tang JJ, Ouyang X, Huang XR, Fingerle-Rowson G, et al. Macrophage migration inhibitory factor induces MMP-9 expression: implications for destabilization of human atherosclerotic plaques. *Atherosclerosis* (2005) 178(1):207-15. doi:10.1016/j.atherosclerosis.2004.08.030
- [61] Leech M, Metz C, Hall P, Hutchinson P, Gianis K, Smith M, et al. Macrophage migration inhibitory factor in rheumatoid arthritis: evidence of proinflammatory function and regulation by glucocorticoids. *Arthritis Rheum* (1999) 42(8):1601-8. doi:10.1002/1529-0131(199908)42:8<1601::AID-ANR6>3.0.CO;2-B
- [62] Lin SG, Yu XY, Chen YX, Huang XR, Metz C, Bucala R, et al. De novo expression of macrophage migration inhibitory factor in atherogenesis in rabbits. *Circ Res* (2000) 87(12):1202-8. doi:10.1161/01.RES.87.12.1202
- [63] Amin MA, Haas CS, Zhu K, Mansfield PJ, Kim MJ, Lackowski NP, et al. Migration inhibitory factor up-regulates vascular cell adhesion molecule-1 and intercellular adhesion molecule-1 via Src, PI3 kinase, and NFkappaB. *Blood* (2006) 107(6):2252-61. doi:10.1182/blood-2005-05-2011
- [64] Sugimoto T, Ishikawa Y, Yoshimoto T, Hayashi N, Fujimoto J, Nakanishi K. Interleukin 18 acts on memory T helper cells type 1 to induce airway inflammation and hyperresponsiveness in a naive host mouse. *J Exp Med* (2004) 199(4):535-45. doi:10.1084/jem.20031368
- [65] Wereszczynska-Siemiatkowska U, Mroczko B, Siemiatkowski A. Serum profiles of interleukin-18 in different severity forms of human acute pancreatitis. *Scand J Gastroenterol* (2002) 37(9):1097-102. doi:10.1080/003655202320378310
- [66] Cerqueira BA, Boas WV, Zanette AD, Reis MG, Goncalves MS. Increased concentrations of IL-18 and uric acid in sickle cell anemia: contribution of hemolysis, endothelial activation and the inflammasome. *Cytokine* (2011) 56(2):471-6. doi:10.1016/j.cyto.2011.08.013
- [67] Johansson SE, Hall H, Björklund J, Höglund P. Broadly impaired NK cell function in non-obese diabetic mice is partially restored by NK cell activation in vivo and by IL-12/IL-18 in vitro. *Int Immunol* (2004) 16(1):1-11. doi:10.1093/intimm/dxh002

[68] Zaccone P, Phillips J, Conget I, Cooke A, Nicoletti F. IL-18 binding protein fusion construct delays the development of diabetes in adoptive transfer and cyclophosphamide-induced diabetes in NOD mouse. *Clin Immunol* (2005) 115(1):74-9. doi:10.1016/j.clim.2004.11.007