

BioNanoScience 2017 vol.7 N1, pages 254-257

Action of ATP on Ca²⁺-Transient in Different Parts of the Frog Motor Nerve Ending

Khaziev E., Golovyahina A., Bukharaeva E., Nikolsky E., Samigullin D. *Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia*

Abstract

© 2016, Springer Science+Business Media New York. Electrophysiological evidence indicates a difference in neurotransmitter secretion along the motor nerve terminals of the frog neuromuscular junction. This includes a decrease of the minimal synaptic delay value and a reduction in the quantal content of the evoked endplate currents from the proximal to distal portion of the motor nerve ending. Besides, various physiologically active compounds may have different effects on the acetylcholine secretion in the proximal and distal parts of the nerve terminal. Here, we explored the effects of ATP on Ca2+-transient using optical detection methods with high-speed camera in different parts of the frog nerve terminal. There was shown a significant inhibitory effect of ATP on Ca2+-transient in both the proximal and distal regions of nerve terminals. However, in different parts of nerve endings, any significant differences in ATP effects were not found. Thus, ATP decreases the Ca2+-transient along the entire presynaptic terminal.

http://dx.doi.org/10.1007/s12668-016-0350-6

Keywords

ATP, Ca -transient 2+, Frog, Neuromuscular junction, Proximal-distal gradient, Synapse

References

- [1] Bennett, M. (1983). Development of neuromuscular synapses. Physiological Reviews, 63, 915-1048.
- [2] Grinnell, A., & Trussell, L. (1983). Synaptic length as a function of motor unit size in the normal frog sartorius. Journal of Physiology, 338, 221–241.
- [3] Braun, M., & Schmidt, R. (1961). Potential changes recorded from motor nerve terminal during its activation. Pflügers Archiv, 287, 56–80.
- [4] Zefirov, A. L., & Khalilov, I. A. (1985). Characteristics of electrical activity in different sections of nerve endings in the frog. Bulletin of Experimental Biology and Medicine, 99(1), 7–10.
- [5] Mallart, A. (1984). Presynaptic currents in frog motor endings. Pflügers Archiv, 400, 8–20.
- [6] Nikolsky, E. E., Bukharaeva, E. A., Samigullin, D. V., Gainulov, R. (2002). Characteristics of the time course of evoked secretion of transmitter quanta in different parts of the motor nerve ending in the frog. Neuroscience and Behavioral Physiology, 32, 265–274.
- [7] Zefirov, A. L. (1983). Mediator secretion in the proximal and distal portions of a nerve ending of the sartorius muscle in the frog. Neirofiziologiia, 15(4), 362–369.
- [8] Bukharaeva, E., Samigullin, D., Nikolsky, E., Vyskočil, F. (2002). Protein kinase a cascade regulates quantal release dispersion at frog muscle endplate. Journal of Physiology, 538, 837–848.

- [9] Tsentsevitsky, A. N., Nikolsky, E. E., Giniatullin, R. A., Bukharaeva, E. A. (2011). Opposite modulation of time course of quantal release in two parts of the same synapse by reactive oxygen species. Neuroscience, 189, 93–99.
- [10] Nikolsky, E. E., Vyskocil, F., Bukharaeva, E. A., Samigullin, D. V., Magazanik, L. G. (2004). Cholinergic regulation of the evoked quantal release at frog neuromuscular junction. Journal of Physiology (L), 560, 77–88.
- [11] Ribeiro, J. A., & Walker, J. (1975). The effects of adenosine triphosphate and adenosine diphosphate on transmission at the rat and frog neuromuscular junctions. British Journal of Pharmacology, 54(2), 213–218.
- [12] Samigullin, D. V., Vasin, A. L., Bukharaeva, E. A., Nikolsky, E. E. (2010). Characteristics of calcium transient in different parts of frog nerve terminal in response to nerve impulse. Doklady Biological Sciences, 431, 83–85.
- [13] Sokolova, E., Grishin, S., Shakirzyanova, A., Talantova, M., Giniatullin, R. (2003). Distinct receptors and different transduction mechanisms for ATP and adenosine at the frog motor nerve endings. European Journal of Neuroscience, 63, 915–1048.
- [14] Tsien, R. Y. (1989). Fluorescent indicators of ion concentrations. Methods in Cell Biology, 30, 127–156.
- [15] Khaziev, E. F., Fatikhov, N. F., Samigullin, D. V., Barrett, G., Bukharaeva, E. A., Nikolsky, E. E. (2012). Decreased entry of calcium into motor nerve endings upon activation of presynaptic cholinergic receptors. Doklady Biological Sciences, 446, 283–285.
- [16] Kazakov, A., Aleksandrov, M., Zhilyakov, N., Khaziev, E., Samigullin, D. (2015). A simple suction electrode for electrical stimulation of biological objects. International Research Journal, 40, 13–16.