

BioNanoScience 2017 vol.7 N2, pages 376-377

---

# Peculiarity of Postoperative Period in Hemodialysis Patients after Parathyroidectomy

Damocev V., Belyaev A., Salieva A., Khasanova M., Minabutdinov R., Vakhitova R.  
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

---

## Abstract

© 2016, Springer Science+Business Media New York. Signs of hypocalcemia of varying degrees of intensity are often shown in hemodialysis patients after parathyroidectomy for severe hyperparathyroidism on the background of the “hungry bone” syndrome. Teaching of patients to distinguish early clinical signs of hypocalcemia and to inform the medical staff about them in time and regular monitoring of ionized calcium with a personalized approach to introduction of calcium preparations in parenteral and oral form promote the favorable course of the postoperative period and help to avoid severe complications. Significant reduction of bone pain syndrome and improvement of general health in dialysis patients are noted in the early postoperative period.

<http://dx.doi.org/10.1007/s12668-016-0373-z>

---

## Keywords

Hemodialysis patients, Hyperparathyroidism, Hypocalcemia, Parathyroidectomy

## References

- [1] Kidney Disease. Improving Global Outcomes (KDIGO) CKD- MBD Work Group. (2009). KDIGO clinical practice guideline for the diagnosis, evolution, prevention, and treatment of chronic kidney disease- mineral and bone disorder (CKD-MBD). *Kidney International*, 76(113), 1-130.
- [2] Ermolenko, V. M., Volgina, G. V., Dobronravov, V. A., et al. (2010). Nacional'nye rekomendacii po mineral'nym i kostnym narushenijam pri hronicheskoj bolezni pochek. Rossijskoe dializnoe obshhestvo [National recommendations of mineral and bone disorders in patients with chronic renal disease, Russian Dialysis Society]. *Nefrologija i dializ [Nephrology and Dialysis]*, 17(3), 33-51.
- [3] Shutov, E. V., Lashutin, S. V., Ljuosev, V. S., et al. (2011). Novye vozmozhnosti v lechenii vtorichnogo giperparatireoza u bol'nyh na programmnom gemodialize pri kombinirovannoj terapii cinakal'cetom i malymi dozami aktivnogo metabolita vitamina D [New opportunities in the treatment of secondary hyperparathyroidism in patients on haemodialysis therapy in combination with treatment with small doses of cinacalcet and the active metabolite of vitamin D. *Klinicheskaja Nefrologija [Clinical Nephrology]*, 5, 41-44.
- [4] Meola, M., Petrucci, I., Barsotti, G. (2009). Long-term treatment with cinacalcet and conventional therapy reduces parathyroid hyperplasia in severe secondary hyperparathyroidism. *Nephrology, Dialysis, Transplantation*, 24, 982-989.
- [5] Goldfarb, M., Gondek, S. S., Lim, S. M., et al. (2012). Postoperative hungry bone syndrome in patients with secondary hyperparathyroidism of renal origin. *World Journal of Surgery*, 36(6), 1314-1319.
- [6] Hamouda, M., Ben Dhia, N., Aloui, S., et al. (2013). Predictors of early post-operative hypocalcemia after parathyroidectomy for secondary hyperparathyroidism. *Saudi Journal of Kidney Diseases and Transplantation*, 24(6), 1165-1169.

- [7] Cozzolino, M., Gallieni, M., Corsi, C., Bastagli, A., Brancaccio, D. (2004). Management of calcium refilling post-parathyroidectomy in end-stage renal disease. *J. Nephrology*, 17(1), 3-8.
- [8] Dzhon T. Daugirdas, Piter Dzh.Blejk, Todd S. Ing. (2003). Rukovodstvo po dializu [Dialysis guideline] Izdatel'stvo «Triada» [publishing office «Triada»]. Twer.