BioNanoScience 2017 vol.7 N1, pages 73-74

Semaphorins Are Likely to Be Involved in the Control of Hibernation

Gazizova G., Kozlova O., Tyapkina O., Nurullin L., Shagimardanova E. *Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia*

Abstract

© 2016, Springer Science+Business Media New York.Hibernation is unique mammals' phenotype demonstrated surviving in seasonal adverse environment conditions. During hibernation, all of systems of organs undergo alterations in their metabolic activity and physiology similar to different physiological conditions associated with human diseases and injuries, which represent a proper model for studying new approaches in clinical treatment. In this study, we have analyzed expression of semaphorins in small hibernator edible dormouse (Glis glis). Semaphorins are factors which involved in a key process of axon guidance and cell-cell communication and can act as tumor suppressor. We found that gene coding Semaphorin-3D (SEMA3D), previously reported to be specific for brain and heart of mammals, represents one of the most upregulated transcripts in the muscles of the hibernating dormice. Furthermore, another member of the same family, Semaphorin-5B (SEMA5B), was strongly induced in the spinal cord of hibernating animals. These observations make semaphorin group, recently attracting more attention due to anti-tumor activity, one of the target for in-depth analysis in relation to the molecular mechanisms of hibernation.

http://dx.doi.org/10.1007/s12668-016-0292-z

Keywords

Edible dormice, Gene expression, Hibernation, Semaphorin

References

- Hoelzl, F., Bieber, C., Cornils, J. S., Gerritsmann, H., Stalder, G. L., Walzer, C., et al. (2015). How to spend the summer? Free-living dormice (Glis glis) can hibernate for 11 months in non-reproductive years. Journal of Comparative Physiology B, 185(8), 931–939. doi:10.1007/s00360-015-0929-1.
- [2] Bieber, C., & Ruf, T. (2009). Summer dormancy in edible dormice (Glis glis) without energetic constraints. Naturwissenschaften, 96, 165–171. doi:10.1007/s00114-008-0471-z.
- [3] Worzfeldand, T., & Offermanns, S. (2014). Semaphorins and plexins as therapeutic targets. Nature Reviews Drug Discovery, 13(8), 603–21. doi:10.1038/nrd4337.
- [4] Mecollari V, Nieuwenhuis B, Verhaagen J. (2014) A perspective on the role of class III semaphorin signaling in central nervous system trauma. Frontiers in Cellular Neuroscience, 8 328. doi:10.3389/fncel.2014.00328.
- [5] Schwartz, C., Hampton, M., Andrews, M. T. (2013). Seasonal and regional differences in gene expression in the brain of a hibernating mammal. PLoS One, 8(3), e58427. doi:10.1371/journal.pone.0058427.