

BioNanoScience 2017 vol.7 N1, pages 201-203

---

## Detection of Antibodies Recognizing Puumala Virus Nucleocapsid and Glycoprotein Peptides in NE Serum

Garanina E., Martynova E., Ivanova V., Okhezin E., Shakirova V., Anokhin V., Morzunov S., Rizvanov A., Khaiboullina S.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

---

### Abstract

© 2016, Springer Science+Business Media New York. Nephropatia epidemica (NE), a mild form of hemorrhagic fever with renal syndrome (HFRS), is an endemic zoonosis in the Republic of Tatarstan, Russia. Humans become infected by inhaling an aerosol contaminated with Puumala virus, a member of genus Hantavirus. NE diagnosis is based on detection of anti-hantavirus antibodies using ELISA. Antibodies to hantavirus nucleocapsid (N) protein are detected early in the course of infection, suggesting that this viral protein is the most immunogenic. Several epitopes were previously identified on N protein as well as glycoproteins of Puumala viruses endemic in Europe. However, there is limited knowledge about Puumala virus N protein antigenic epitopes in NE patients in the Republic of Tatarstan. The aim of the present study is to identify N protein and glycoprotein epitopes which induce a humoral immune response in NE cases. Analysis of NE serum using an array of overlapping N protein and glycoprotein peptides identified the most immunogenic epitopes, which can then be used for developing Puumala virus-specific vaccine.

<http://dx.doi.org/10.1007/s12668-016-0296-8>

---

### Keywords

ELISA, Hemorrhagic fever with renal syndrome, Nucleocapsid protein, Peptides

### References

- [1] Khismatullina, N. A., et al. (2016). Epidemiological dynamics of nephropatia epidemica in the Republic of Tatarstan, Russia, during the period of 1997-2013. *Epidemiology and Infection*, 144(3), 618-626.
- [2] Uchaykin V.F., N.N.I., Shamsheva O.V., *Infectious diseases in children*. 2006, Moscow: GEOTAR-Media. 326-331.
- [3] Groen, J., et al. (1992). Serum antibodies to structural proteins of Hantavirus arise at different times after infection. *Journal of Medical Virology*, 37(4), 283-287.
- [4] Marchand, E., et al. (2013). Autochthonous case of dengue in France, October 2013. *Euro Surveillance*, 18(50), 20661.
- [5] Latus, J., et al. (2015). Clinical course and long-term outcome of hantavirus-associated nephropatia epidemica, Germany. *Emerg Infect Dis*, 21(1), 76-83.
- [6] Lundkvist, A., et al. (1995). Mapping of B-cell determinants in the nucleocapsid protein of Puumala virus: definition of epitopes specific for acute immunoglobulin G recognition in humans. *Clinical and Diagnostic Laboratory Immunology*, 2(1), 82-86.
- [7] Tischler, N. D., et al. (2005). Human and rodent humoral immune responses to Andes virus structural proteins. *Virology*, 334(2), 319-326.

- [8] Vapalahti, O., et al. (1995). Human B-cell epitopes of Puumala virus nucleocapsid protein, the major antigen in early serological response. *Journal of Medical Virology*, 46(4), 293-303.
- [9] Hjelle, B., et al. (1994). Dominant glycoprotein epitope of four corners hantavirus is conserved across a wide geographical area. *Journal of General Virology*, 75(Pt 11), 2881-2888.
- [10] Wang, M., Rossi, C., Schmaljohn, C. S. (1993). Expression of non-conserved regions of the S genome segments of three hantaviruses: evaluation of the expressed polypeptides for diagnosis of haemorrhagic fever with renal syndrome. *Journal of General Virology*, 74(Pt 6), 1115-1124.