

Single-Molecule Interactions of a Monoclonal Anti-DNA Antibody with DNA

Nevzorova T., Zhao Q., Lomakin Y., Ponomareva A., Mukhitov A., Purohit P., Weisel J., Litvinov R.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2016, Springer Science+Business Media New York. Interactions of DNA with proteins are essential for key biological processes and have both a fundamental and practical significance. In particular, DNA binding to anti-DNA antibodies is a pathogenic mechanism in autoimmune pathology, such as systemic lupus erythematosus. Here we measured at the single-molecule level binding and forced unbinding of surface-attached DNA and a monoclonal anti-DNA antibody MRL4 from a lupus erythematosus mouse. In optical trap-based force spectroscopy, a microscopic antibody-coated latex bead is trapped by a focused laser beam and repeatedly brought into contact with a DNA-coated surface. After careful discrimination of non-specific interactions, we showed that the DNA-antibody rupture force spectra had two regimes, reflecting formation of weaker (20–40 pN) and stronger (>40 pN) immune complexes that implies the existence of at least two bound states with different mechanical stability. The two-dimensional force-free off-rate for the DNA-antibody complexes was $\sim 2.2 \times 10^{-3} \text{ s}^{-1}$, the transition state distance was $\sim 0.94 \text{ nm}$, the apparent on-rate was $\sim 5.26 \text{ s}^{-1}$, and the stiffness of the DNA-antibody complex was characterized by a spring constant of 0.0021 pN/nm, suggesting that the DNA-antibody complex is a relatively stable, but soft and deformable macromolecular structure. The stretching elasticity of the DNA molecules was characteristic of single-stranded DNA, suggesting preferential binding of the MRL4 antibody to one strand of DNA. Collectively, the results provide fundamental characteristics of formation and forced dissociation of DNA-antibody complexes that help to understand principles of DNA-protein interactions and shed light on the molecular basis of autoimmune diseases accompanied by formation of anti-DNA antibodies.

<http://dx.doi.org/10.1007/s12668-016-0303-0>

Keywords

Anti-DNA antibody, DNA, Nanomechanics, Optical trap, Single-molecule force spectroscopy, Two-dimensional kinetics

References

- [1] Pisetsky, D. S. (1998). Antibody responses to DNA in normal immunity and aberrant immunity. *Clinical and Diagnostic Laboratory Immunology*, 5, 1–6.
- [2] Williams, W. M., & Isenberg, D. A. (1996). A cross-sectional study of anti-DNA antibodies in the serum and IgG and IgM fraction of healthy individuals, patients with systemic lupus erythematosus and their relatives. *Lupus*, 5, 576–586.

- [3] Uchida, K. (2014). Natural antibodies as a sensor of electronegative damage-associated molecular patterns (DAMPs). *Free Radical Biology and Medicine*, 72, 156-161.
- [4] Lora, V., Bonaguri, C., Gisondi, P., Sandei, F., Battistelli, L., Russo, A., et al. (2013). Autoantibody induction and adipokine levels in patients with psoriasis treated with infliximab. *Immunologic Research*, 56, 382-389.
- [5] Quan, C. P., Watanabe, S., Pamonsinlapatham, P., Bouvet, J.-P. (2001). Different dysregulations of the natural antibody repertoire in treated and untreated HIV-1 patients. *Journal of Autoimmunity*, 17, 81-87.
- [6] Williamson, R. A., Burgoon, M. P., Owens, G. P., Ghausi, O., Leclerc, E., Firme, L., et al. (2001). Anti-DNA antibodies are a major component of the intrathecal B cell response in multiple sclerosis. *Proceedings of the National Academy of Sciences of the United States of America*, 98, 1793-1798.
- [7] Compagno, M., Rekvig, O. P., Bengtsson, A. A., Sturfelt, G., Heegaard, N. H., Jønsen, A., et al. (2014). Clinical phenotype associations with various types of anti-dsDNA antibodies in patients with recent onset of rheumatic symptoms. Results from a multicentre observational study. *Lupus Science and Medicine*, 1, e000007.
- [8] Gorny, M. K., Lawniczak, M., Jenek, R., Słowik-Gabryelska, A., Kaczmarek, E., Zeromski, J. (1988). Alloantibodies, autoantibodies, and immune complexes in patients with lung cancer. *Lung*, 166, 97-105.
- [9] Barbas, S. M., Ditzel, H. J., Saloneh, E. M., Yang, W.-P., Silverman, G. J., Burton, D. R. (1995). Human autoantibody recognition of DNA. *Proceedings of the National Academy of Sciences of the United States of America*, 92, 2529-2533.
- [10] Suenaga, R., & Abdou, N. I. (1993). Cationic and high affinity serum IgG anti-dsDNA antibodies in active lupus nephritis. *Clinical and Experimental Immunology*, 94, 418-422.
- [11] Deocharan, B., Qing, X., Beger, E., Puttermann, C. (2002). Antigenic triggers and molecular targets for anti-double-stranded DNA antibodies. *Lupus*, 11, 865-871.
- [12] Madaio, M. P., & Yanase, K. (1998). Cellular penetration and nuclear localization of anti-DNA antibodies: mechanisms, consequences, implications and applications. *Journal of Autoimmunity*, 11, 535-538.
- [13] Puttermann, C. (2004). New approaches to the renal pathogenicity of anti-DNA antibodies in systemic lupus erythematosus. *Autoimmunity Reviews*, 3, 7-11.
- [14] Ullala, A. J., Marion, T. N., Pisetsky, D. S. (2014). The role of antigen specificity in the binding of murine monoclonal anti-DNA antibodies to microparticles from apoptotic cells. *Clinical Immunology*, 154, 178-187.
- [15] Foster, M. H., Kieber-Emmons, T., Ohliger, M., Madaio, M. P. (1994). Molecular and structural analysis of nuclear localizing anti-DNA lupus antibodies. *Immunologic Research*, 13, 186-206.
- [16] Aranow C, Zhou D, Diamond B (2011) Anti-DNA antibodies: structure, regulation and pathogenicity. In: Lahita RG, Tsokos G, Buyon JP, Koike T (eds.) *Systemic lupus erythematosus*, Fifth Ed. Academic Press, pp 235-258.
- [17] Shlomchik, M., Mascelli, M., Shan, H., Radic, M. Z., Pisetsky, D., Marshak-Rothstein, A., et al. (1990). Anti-DNA antibodies from autoimmune mice arise by clonal expansion and somatic mutation. *Journal of Experimental Medicine*, 171, 265-292.
- [18] Uccellini, M. B., Bustó, P., Debatis, M., Marshak-Rothstein, A., Viglianti, G. A. (2012). Selective binding of anti-DNA antibodies to native dsDNA fragments of differing sequence. *Immunology Letters*, 143, 85-91.
- [19] Jin, H., Sepulveda, J., Burrone, O. R. (2004). Specific recognition of a dsDNA sequence motif by an immunoglobulin VH homodimer. *Protein Science*, 13, 3222-3229.
- [20] Eivazova, E. R., McDonnell, J. M., Sutton, B. J., Staines, N. A. (2000). Specificity and binding kinetics of murine lupus anti-DNA monoclonal antibodies implicate different stimuli for their production. *Immunology*, 101, 371-377.
- [21] Chaurasiya, K. R., Paramanathan, T., McCauley, M. J., Williams, M. C. (2010). Biophysical characterization of DNA binding from single molecule force measurements. *Physics of Life Reviews*, 7, 299-341.
- [22] Noy, A. (2011). Force spectroscopy 101: how to design, perform, and analyze an AFM-based single molecule force spectroscopy experiment. *Current Opinion in Chemical Biology*, 15, 710-718.
- [23] Rocha, M. S. (2015). Extracting physical chemistry from mechanics: a new approach to investigate DNA interactions with drugs and proteins in single molecule experiments. *Integrative Biology (Camb)*, 7, 967-986.
- [24] Chung, J. W., Shin, D., Kwak, J. M., Seog, J. (2013). Direct force measurement of single DNA-peptide interactions using atomic force microscopy. *Journal of Molecular Recognition*, 26, 268-275.
- [25] Bartels, F., McIntosh, M., Fuhrmann, A., Metzendorf, C., Plattner, P., Sewald, N., et al. (2007). Effector-stimulated single molecule protein-DNA interactions of a quorum-sensing system in *Sinorhizobium meliloti*. *Biophysical Journal*, 92, 4391-4400.
- [26] Wollschlager, K., Gaus, K., Kornig, A., Eckel, R., Wilking, S., McIntosh, M., et al. (2009). Single-molecule experiments to elucidate the minimal requirement for DNA recognition by transcription factor epitopes. *Small*, 5, 484-495.
- [27] Koch, S., & Wang, M. (2003). Dynamic force spectroscopy of protein-DNA interactions by unzipping DNA. *Physical Review Letters*, 91, 028103.
- [28] Krasnoslobodtsev, A. V., Shlyakhtenko, L. S., Lyubchenko, Y. L. (2007). Probing interactions within the synaptic DNA-Sfil complex by AFM force spectroscopy. *Journal of Molecular Biology*, 365, 1407-1416.

- [29] Sanchez, H., Suzuki, Y., Yokokawa, M., Takeyasu, K., Wyman, C. (2011). Protein-DNA interactions in high speed AFM: single molecule diffusion analysis of human RAD54. *Integrative Biology*, 3, 1127-1134.
- [30] Lionnet, T., Allemand, J. F., Revyakin, A., Strick, T. R., Saleh, O. A., Bensimon, D., et al. (2012). Single-molecule studies using magnetic traps. *Cold Spring Harbor Protocols*, 2012, 34-49.
- [31] Limmer, K., Pippig, D. A., Aschenbrenner, D., Gaub, H. E. (2014). A force-based, parallel assay for the quantification of protein-DNA interactions. *PLoS One*, 9, e89626.
- [32] Kofler, R., Strohal, R., Balderas, R. S., Johnson, M. E., Noonan, D. J., Duchosal, M. A., et al. (1988). Immunoglobulin kappa light chain variable region gene complex organization and immunoglobulin genes encoding anti-DNA autoantibodies in lupus mice. *Journal of Clinical Investigation*, 82, 852-860.
- [33] Litvinov, R. I., Gorkun, O. V., Owen, S. F., Shuman, H., Weisel, J. W. (2005). Polymerization of fibrin: specificity, strength, and stability of knob-hole interactions studied at the single-molecule level. *Blood*, 106, 2944-2951.
- [34] Litvinov, R. I., Yarovoi, S. V., Rauova, L., Barsegov, V., Sachais, B. S., Rux, A. H., et al. (2013). Distinct specificity and single-molecule kinetics characterize the interaction of pathogenic and non-pathogenic antibodies against platelet factor 4-heparin complexes with platelet factor 4. *Journal of Biological Chemistry*, 288, 33060-33070.
- [35] Litvinov, R. I., Shuman, H., Bennett, J. S., Weisel, J. W. (2002). Binding strength and activation state of single fibrinogen-integrin pairs on living cells. *Proceedings of the National Academy of Sciences of the United States of America*, 99, 7426-7431.
- [36] Litvinov, R. I., Mekler, A., Shuman, H., Bennett, J. S., Barsegov, V., Weisel, J. W. (2012). Resolving two-dimensional kinetics of the integrin $\alpha IIb\beta 3$ -fibrinogen interactions using binding-unbinding correlation spectroscopy. *Journal of Biological Chemistry*, 287, 35275-35285.
- [37] Evans, E., & Ritchie, K. (1997). Dynamic strength of molecular adhesion bonds. *Biophysical Journal*, 72, 1541-1555.
- [38] Tees, D. F. J., Woodward, J. T., IV, Hammer, D. A. (2001). Reliability theory for receptor-ligand bond dissociation. *Journal of Chemical Physics*, 114, 7483-7496.
- [39] Bell, G. I. (1978). Models for the specific adhesion of cells to cells. *Science*, 200, 618-627.
- [40] Bura, E., Zhmurov, A., Barsegov, V. (2009). Nonparametric density estimation and optimal bandwidth selection for protein unfolding and unbinding data. *Journal of Chemical Physics*, 130, 015102.
- [41] Nelson P (2004) Biological physics. In: Freeman WH and company, New York.
- [42] Zhang, Y., Zhou, H., Ou-Yang, Z. C. (2001). Stretching single-stranded DNA: interplay of electrostatic, base-pairing, and base-pair stacking interactions. *Biophysical Journal*, 81, 1133-1143.
- [43] Jorgensen, M. H., Rekvig, O. P., Jacobsen, R. S., Jacobsen, S., Fenton, K. A. (2011). Circulating levels of chromatin fragments are inversely correlated with anti-dsDNA antibody levels in human and murine systemic lupus erythematosus. *Immunology Letters*, 138, 179-186.
- [44] Spatz, L., Iliev, A., Saenko, V., Jones, L., Irigoyen, M., Manheimer-Lory, A., et al. (1997). Studies on the structure, regulation, and pathogenic potential of anti-dsDNA antibodies. *Methods*, 11, 70-78.
- [45] Kozyr, A. V., Kolesnikov, A. V., Khlyntseva, A. E., Bogun, A. G., Savchenko, G. A., Shemyakin, I. G., et al. (2012). Role of structure-based changes due to somatic mutation in highly homologous DNA-binding and DNA-hydrolyzing autoantibodies exemplified by A23P substitution in the VH domain. *Autoimmune Diseases*, 2012, 683829.
- [46] Lomakin, Y. A., Zakharova, M. Y., Stepanov, A. V., Dronina, M. A., Smirnov, I. V., Bobik, T. V., et al. (2014). Heavy-light chain interrelations of MS-associated immunoglobulins probed by deep sequencing and rational variation. *Molecular Immunology*, 62, 305-314.
- [47] Pavlovic, M., Kats, A., Cavallo, M., Chen, R., Hartmann, J. X., Shoenfeld, Y. (2010). Pathogenic and epiphenomenal anti-DNA antibodies in SLE. *Autoimmune Diseases*, 2010, 462841.
- [48] Liu, Y. Y., Guthold, M., Snyder, M. J., Lu, H. F. (2015). AFM of self-assembled lambda DNA-histone networks. *Colloids and Surfaces, B: Biointerfaces*, 134, 17-25.
- [49] Nawas, S., Sanches, P., Bodensiek, K., Li, S., Simons, M., Schaap, I. A. (2012). Cell visco-elasticity measured with AFM amp optical trapping at sub-micrometer deformations. *PLoS One*, 7, e45297.
- [50] Weisel, J. W., Shuman, H., Litvinov, R. I. (2003). Protein-protein unbinding induced by force: single-molecule studies. *Current Opinion in Structural Biology*, 13, 227-235.
- [51] Goldsmith, H. L., McIntosh, F. A., Shahin, J., Frojmovic, M. M. (2000). Time and force dependence of the rupture of glycoprotein IIb-IIIa-fibrinogen bonds between latex spheres. *Biophysical Journal*, 78, 1195-1206.
- [52] Lee, I., & Marchant, R. E. (2001). Force measurements on the molecular interactions between ligand (RGD) and human platelet $\alpha IIb\beta 3$ receptor system. *Surface Science*, 491, 433-443.
- [53] Zhu, C., Long, M., Chesla, S. E., Bongrand, P. (2002). Measuring receptor/ligand interaction at the single-bond level: experimental and interpretative issues. *Annals of Biomedical Engineering*, 30, 305-314.
- [54] Litvinov, R. I., Bennett, J. S., Weisel, J. W., Shuman, H. (2005). Multi-step fibrinogen binding to the integrin $\alpha IIb\beta 3$ detected using force spectroscopy. *Biophysical Journal*, 89, 2824-2834.

- [55] Litvinov, R. I., Gorkun, O. V., Galanakis, D. K., Yakovlev, S., Medved, L., Shuman, H., et al. (2007). Polymerization of fibrin: direct observation and quantification of individual B:b knob-hole interactions. *Blood*, 109, 130-138.
- [56] Chilkoti, A., & Stayton, P. S. (1995). Molecular origins of the slow streptavidin-biotin dissociation kinetics. *Journal of the American Chemical Society*, 117, 10622-10628.
- [57] Poongavanam, M.-V., Kisley, L., Kourentzi, K., Landes, C. F., Willson, R. C. (2016). Ensemble and single-molecule biophysical characterization of D17.4 DNA aptamer-IgE interactions. *Biochimica et Biophysica Acta, Proteins Proteomics*, 1864, 154-164.
- [58] Schwesinger, F., Ros, R., Strunz, T., Anselmetti, D., Güntherodt, H. J., Honegger, A., et al. (2000). Unbinding forces of single antibody-antigen complexes correlate with their thermal dissociation rates. *Proceedings of the National Academy of Sciences of the United States of America*, 97, 9972-9977.
- [59] Giuntoli, R. D., Linzer, N. B., Banigan, E. J., Sing, C. E., de la Cruz, M. O., Graham, J. S., et al. (2015). DNA-segment-facilitated dissociation of Fis and NHP6A from DNA detected via single-molecule mechanical response. *Journal of Molecular Biology*, 427, 3123-3136.