

International Multidisciplinary Scientific GeoConference Surveying Geology and Mining Ecology Management, SGEM 2018 vol.18 N1.4, pages 541-546

---

## Integration of the seismic and geochemistry data to evaluate hydrocarbon potential of the carbonate reservoirs in Tatarstan, Russia

Bagmanov I., Safina R., Platov B., Usmanov S.  
*Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia*

---

### Abstract

© SGEM2018. The article describes the results of the integration of the geochemical and seismic data on one of the oilfields of Tatarstan Republic in Russian Federation. The complex geological structure of the oil bearing formations of Carboniferous age results in misinterpretation of the geophysical data and drilling of the dry wells. Some potential structural oil traps find from the seismic data interpretation are water bearing. To avoid nonproductive drilling authors studied seismic faults and their connection with the geochemical anomalies. On the first step the faults in the potential oil-bearing formation of the Tournaisian age were traced in the 2D seismic lines. Then the geochemical parameter (propane concentration in the soils) was studied in 90 observation points. The gas anomaly represented by propane is indicative, because the biochemical genesis of methane homologues is practically excluded, and their content in coal is insignificant. That means that the increased content of propane is connected with the presence of hydrocarbons. It appears that in the presents of oil in the formation geochemical anomaly and the fault in the Tournaisian formation are coincide. That can be used as additional source of information to avoid nonproductive drilling.

<http://dx.doi.org/10.5593/sgem2018/1.4/S06.070>

---

### Keywords

Geochemical anomalies, Seismic data, Seismic faults

### References

- [1] Platov B.V, Khamidullina G.S, Nouriev I.A, Novikov, I.P., Ognev, I.N., The example of reconstructing carbonate sedimentation process on the basis of seismic-stratigraphic and paleogeomorphological analysis, *Neftyanoe Khozyaystvo, Russia*, issue 1, pp 18-22, 2018;
- [2] Kolchugin, A.N., Morozov, V.P., Korolev, E.A., Eskin, A.A., Gazeeva, F.M., Typical sections of bashkirian carbonate rocks and structure of oil deposits in southeast part of the Republic of Tatarstan, *Neftyanoe Khozyaystvo-Oil Industry, Russia*, issue 11, pp 84-86, 2013;
- [3] Khamidullina, G.S., Ziganshin, E.R., Minnibajeva, E.I., Khaliullin, R.R., Study of reservoir properties of carbonate rocks based on analysis of reservoir quality index, *Neftyanoe Khozyaystvo-Oil Industry, Russia*, issue 10, pp 64-66, 2015;
- [4] Nurgalieva N. G., Khaziev R. R., Silantiev V. V., Gareev B. I., Batalin G. A., Geochemical examination of the middle-upper Permian rocks, Volga river reference section, *ARPN Journal of Engineering and Applied Sciences, Pakistan*, vol 11, pp 11433 - 11444, 2016.

- [5] Nezhdanov A.A., Kosarev I.V., Agafonov Yu.K., The role of surface gas chemistry studies in improving the efficiency of oil and gas exploration (by the example of the south of the Tyumen region), The Second Scientific and Practical Conference Ways of realizing the oil and gas potential of Khanty-Mansiysk Autonomous Okrug, Russia vol 1, pp 237-249, 1999.
- [6] Kuznetsov O.L., Petukhov A.V., Zorkin L.M., Physicochemical foundations of direct searches for oil and gas deposits, Russia, 336 p., 1986.