

Mycoplasmas and novel HO-1 inducers: Recent advances

Chernov V., Chernova O., Mouzykantov A., Lopukhov L., Trushin M.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 Bentham Science Publishers. Inflammation and the ways for its regulation: The development of an effective system for the treatment of inflammatory diseases requires comprehensive studies of the cellular signaling molecular networks comprising responses to various stressors, including pathogenic and non-pathogenic microorganisms. Significant attention on fundamental and applied research has recently focused on inducers of heme oxygenase-1 (HO-1) and inhibitors of the expression of this enzyme, which regulates expression of this and other cytoprotective molecules and modulation of inflammation. Recent studies indicate that mycoplasmas (a major group of human pathogens of the Mollicutes) are capable of modulating inflammatory responses through the activation of the Nrf2 and the expression of HO-1. In vitro experiments demonstrate that the membrane lipoproteins (LAMPs), along with lipoprotein derivatives (lipopeptide MALP-2) in mycoplasmas cause a "cross-talk" between the pro- and anti-inflammatory signaling pathways. Importantly, lipopeptide/lipoprotein-induced expression of HO-1 tends to suppress inflammation. Conclusion: The study of the molecular network that causes the corresponding outcome can facilitate the development of new approaches for the treatment of inflammatory processes. The derivatives of LAMPs and MALP-2 and of their analogues may prove promising for the treatment of diseases associated with chronic inflammation.

<http://dx.doi.org/10.2174/1381612824666180716170128>

Keywords

Ho-1 inducers, Human pathogens, Inflammatory response modulation, Lipoprotein derivatives, Membrane lipoproteins, Mollicutes, Mycoplasmas

References

- [1] Hunter P. The inflammation theory of disease. *EMBO Rep* 2012; 13: 968-70.
- [2] Tabas I, Glass CK. Anti-inflammatory therapy in chronic disease: challenges and opportunities. *Science* 2013; 339: 166-72.
- [3] Vijayan V, Wagener FADTG, Immenschuh S. The macrophage heme-heme oxygenase-1 system and its role in inflammation. *Biochem Pharmacol* 2018; 13: S0006-2952(18)30067-4.
- [4] Lamkanfi M, Dixit VM. Modulation of inflammasome pathways by bacterial and viral pathogens. *J Immunol* 2011; 187: 597-602.
- [5] Chen K, Shanmugam NKN, Pazos MA, et al. Commensal bacteria-induced inflammasome activation in mouse and human macrophages is dependent on potassium efflux but does not require phagocytosis or bacterial viability. *PLoS One* 2016; 11: e0160937.
- [6] Prawan A, Saw CL, Khor TO, et al. Anti-NF-kappaB and anti-inflammatory activities of synthetic isothiocyanates: Effect of chemical structures and cellular signaling. *Chem Biol Interact* 2009; 179: 202-11.

- [7] Sundaresan NR, Vasudevan P, Zhong L, et al. The sirtuin SIRT6 blocks IGF-Akt signaling and development of cardiac hypertrophy by targeting c-Jun. *Nat Med* 2012; 18: 1643-50.
- [8] Yan X, Lee S, Gugiu BG, et al. Fatty acid epoxyisoprostane E2 stimulates an oxidative stress response in endothelial cells. *Biochem Biophys Res Commun* 2014; 444: 69-74.
- [9] Jaiswal AK. Nrf2 signaling in coordinated activation of antioxidant gene expression. *Free Radic Biol Med* 2004; 36: 1199-207.
- [10] Saidu NE, Touma R, Asali IA, et al. Diallyl tetrasulfane activates both the eIF2 α and Nrf2/HO-1 pathways. *Biochim Biophys Acta* 2013; 1830: 2214-25.
- [11] Bellezza I, Mierla AL, Minelli A. Nrf2 and NF- κ B and their concerted modulation in cancer pathogenesis and progression. *Cancers (Basel)* 2010; 2: 483-97.
- [12] Thimmulappa RK, Lee H, Rangasamy T, et al. Nrf2 is a critical regulator of the innate immune response and survival during experimental sepsis. *J Clin Invest* 2006; 116: 984-95.
- [13] Rushworth SA, MacEwan DJ, O'Connell MA. Lipopolysaccharide-induced expression of NAD(P)H:quinone oxidoreductase 1 and heme oxygenase-1 protects against excessive inflammatory responses in human monocytes. *J Immunol* 2008; 181: 6730-37.
- [14] Ma X, You X, Zeng Y, et al. *Mycoplasma fermentans* MALP-2 induces heme oxygenase-1 expression via mitogen-activated protein kinases and Nrf2 pathways to modulate cyclooxygenase 2 expression in human monocytes. *Clin Vaccine Immunol* 2013; 20: 827-34.
- [15] Razin S, Hayflick L. Highlights of mycoplasma research-an historical perspective. *Biologicals* 2010; 38: 183-90.
- [16] Yavlovich A, Tarshis M, Rottem S. Internalization and intracellular survival of *Mycoplasma pneumoniae* by non-phagocytic cells. *FEMS Microbiol Lett* 2004; 233: 241-6.
- [17] Paessler M, Levinson A, Patel JB, et al. Disseminated *Mycoplasma orale* infection in a patient with common variable immunodeficiency syndrome. *Diagn Microbiol Infect Dis* 2002; 44: 201-4.
- [18] Uphoff CC, Drexler HG. Detecting mycoplasma contamination in cell cultures by polymerase chain reaction. *Methods Mol Biol* 2011; 731: 93-103.
- [19] Rottem S, Kosower NS, Kornspan JD. Contamination of tissue cultures by mycoplasmas. In: L Ceccherini-Nelli, B Matteoli (eds) *Biomedical Tissue Culture*. InTech 2012 pp 35-8. Available from: (<https://www.intechopen.com/books/biomedical-tissue-culture/contamination-of-tissue-cultures-by-mycoplasmas>)
- [20] McGowin CL, Annan RS, Quayle AJ, et al. Persistent *Mycoplasma genitalium* infection of human endocervical epithelial cells elicits chronic inflammatory cytokine secretion. *Infect Immun* 2012; 80: 3842-9.
- [21] Zhang S, Tsai S, Lo S-C. Alteration of gene expression profiles during mycoplasma-induced malignant cell transformation. *BMC Cancer* 2006; 6: 116.
- [22] Elkind E, Vaisid T, Kornspan JD, et al. Calpastatin upregulation in *Mycoplasma hyorhinis*-infected cells is promoted by the mycoplasma lipoproteins via the NF- κ B pathway. *Cell Microbiol* 2012; 14: 840-51.
- [23] He L, You X, Li G, et al. *Mycoplasma genitalium*-derived lipid-associated membrane proteins negatively regulate cytokine secretion by inducing HO-1 expression in placental trophoblast cells. *Xi Bao Yu Fen Zi Mian Yi Xue Za Zhi* 2015; 31: 194-8.
- [24] Hu J, Chen C, Ou G, et al. Nrf2 regulates the inflammatory response, including heme oxygenase-1 induction, by *Mycoplasma pneumoniae* lipid-associated membrane proteins in THP-1 cells. *Pathog Dis* 2017; 75. Available from: (<https://academic.oup.com/femspd/article/75/4/ftx044/3738187>).
- [25] Okusawa T, Fujita M, Nakamura J, et al. Relationship between structures and biological activities of mycoplasmal diacylated lipopeptides and their recognition by Toll-like receptors 2 and 6. *Infect Immun* 2004; 72: 1657-65.
- [26] Shimizu T, Kida Y, Kuwano K. Triacylated lipoproteins derived from *Mycoplasma pneumoniae* activate nuclear factor- κ B through Toll-like receptors 1 and 2. *Immunology* 2007; 121: 473-83.
- [27] Yang H, Chen C. Cyclooxygenase-2 in synaptic signaling. *Curr Pharm Des* 2008; 14: 1443-51.
- [28] Shih RH, Yang CM. Induction of heme oxygenase-1 attenuates lipopolysaccharide-induced cyclooxygenase-2 expression in mouse brain endothelial cells. *J Neuroinflammation* 2010; 7: 86.
- [29] Hecke F, Steinwede K, Muhlradt P, et al. The synthetic TLR-2/6 agonist MALP-2 reduces the bacterial load in the murine *M. bovis* BCG infection. *Am J Respir Crit Care Med* 2009; 179: A5913.
- [30] Grote K, Sonnenschein K, Hillmer A, et al. The Toll-like receptor 2/6 agonist MALP-2 promotes reendothelialization following vascular injury. *Circulation* 2012; 124: A11704.
- [31] Reppe K, Tschernig T, Luhrmann A, et al. Immunostimulation with macrophage-activating lipopeptide-2 increased survival in murine pneumonia. *Am J Respir Cell Mol Biol* 2009; 40: 474-81.
- [32] Deiters U, Barsig J, Tawil B, et al. The macrophage-activating lipopeptide-2 accelerates wound healing in diabetic mice. *Exp Dermatol* 2004; 13: 731-9.
- [33] Chernov VM, Govorun VM, Demina IA, et al. Adaptation of mycoplasmas to adverse growth conditions: morphology, ultrastructure, and genome expression of *Mycoplasma gallisepticum* S6 cells. *Dokl Biochem Biophys* 2008; 421: 231-4.

- [34] Henderson B, Poole S, Wilson M. Bacterial modulins: A novel class of virulence factors which cause host tissue pathology by inducing cytokine synthesis. *Microbiol Rev* 1996; 60: 316-41.
- [35] Into T, Nodasaka Y, Hasebe A et al. Mycoplasmal lipoproteins induce toll-like receptor 2-and caspases-mediated cell death in lymphocytes and monocytes. *Microbiol Immunol* 2002; 46: 265-76.
- [36] Rawadi G, Ramez V, Lemercier B, et al. Activation of mitogen-activated protein kinase pathways by *Mycoplasma fermentans* membrane lipoproteins in murine macrophages: involvement in cytokine synthesis. *J Immunol* 1998; 160: 1330-9.
- [37] Chernov VM, Chernova OA, Medvedeva ES, et al. Unadapted and adapted to starvation *Acholeplasma laidlawii* cells induce different responses of *Oryza sativa*, as determined by proteome analysis. *J. Proteomics* 2011; 74: 2920-36.
- [38] Chernov VM, Mouzykantov AA, Baranova NB, et al. Extracellular membrane vesicles secreted by mycoplasma *Acholeplasma laidlawii* PG8 are enriched in virulence proteins. *J Proteomics* 2014; 110: 117-28.
- [39] Medvedeva ES, Siniagina MN, Malanin SY, et al. Genome sequences of *Acholeplasma laidlawii* strains differing in sensitivity to ciprofloxacin. *Genome Announc* 2017; 5: pii: e01189-17.
- [40] Medvedeva ES, Davydova MN, Mouzykantov AA, et al. Genomic and proteomic profiles of *Acholeplasma laidlawii* strains differing in sensitivity to ciprofloxacin. *Dokl Biochem Biophys* 2016; 466: 23-7.
- [41] Virgin HW. The virome in mammalian physiology and disease. *Cell* 2014; 157: 142-50.