

# The role of polyamine-dependent facilitation of calcium permeable ampars in short-term synaptic enhancement

Rozov A., Zakharova Y., Vazetdinova A., Valiullina-Rakhmatullina F.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

## Abstract

© 2018 Rozov, Zakharova, Vazetdinova and Valiullina-Rakhmatullina. Depending on subunit composition AMPA receptor channels can be subdivided into two groups: GluA2-containing calcium impermeable AMPARs, and GluA2-lacking calcium permeable, AMPARs. These two groups differ in a number of biophysical properties and, most likely, in their functional role at glutamatergic synapses. GluA2-lacking channels have received a lot of attention over the last two decades mainly due to high calcium permeability, which was suggested to play a significant role in the induction of long-term synaptic plasticity in healthy tissue and neuronal death under neuropathological conditions. However, calcium permeable AMPARs possess another property that can contribute substantially to frequency dependent dynamics of synaptic efficacy. In the closed state calcium permeable AMPARs are blocked by endogenous polyamines, however, repetitive activation leads to progressive relief from the block and to the facilitation of ion flux through these channels. Polyamine-dependent facilitation of AMPARs can contribute to short-term plasticity at synapses that have high initial release probability and express calcium permeable AMPARs. During synaptic transmission activity-dependent relief from polyamine block of postsynaptic calcium-permeable AMPARs either counteracts presynaptic short-term depression in a frequency-dependent manner or, under specific stimulation conditions, induces facilitation of a synaptic response. Taking into account the fact that expression of calcium permeable AMPARs is developmentally regulated, depends on network activity and increases in diseased brain states, polyamine-dependent facilitation of calcium permeable AMPARs is an important, entirely postsynaptic mechanism of synaptic gain regulation.

<http://dx.doi.org/10.3389/fncel.2018.00345>

## Keywords

Calcium permeable AMPA, GluA2-lacking AMPA, Homeostatic plasticity, Interneurons, Neurons, Polyamine, Short-term plasticity, Synapse

## References

- [1] Adesnik H., Nicoll R. A., (2007). Conservation of glutamate receptor 2-containing AMPA receptors during long-term potentiation. *J. Neurosci.* 27 4598–4602. 10.1523/JNEUROSCI.0325-07.2007 17460072.
- [2] Angulo M. C., Rossier J., Audinat E., (1999). Postsynaptic glutamate receptors and integrative properties of fast-spiking interneurons in the rat neocortex. *J. Neurophysiol.* 82 1295–1302. 10.1152/jn.1999.82.3.1295 10482748.

- [3] Bai X., Wong-Riley M. T., (2003). Neuronal activity regulates protein and gene expressions of GluR2 in postnatal rat visual cortical neurons in culture. *J. Neurocytol.* 32 71-78. 10.1023/A:1027380315902 14618102.
- [4] Bellone C., Mameli M., Luscher C., (2011). In utero exposure to cocaine delays postnatal synaptic maturation of glutamatergic transmission in the VTA. *Nat. Neurosci.* 14 1439-1446. 10.1038/nn.2930 21964489.
- [5] Bernstein H. G., Muller M., (1995). Increased immunostaining for L-ornithine decarboxylase occurs in neocortical neurons of Alzheimer's disease patients. *Neurosci. Lett.* 186 123-126. 10.1016/0304-3940(95)11301-C 7777179.
- [6] Bollmann J. H., Helmchen F., Borst J. G., Sakmann B., (1998). Postsynaptic Ca influx mediated by three different pathways during synaptic transmission at a calyx-type synapse. *J. Neurosci.* 18 10409-10419. 10.1523/JNEUROSCI.18-24-10409.1998.
- [7] Bowie D., Mayer M. L., (1995). Inward rectification of both AMPA and kainate subtype glutamate receptors generated by polyamine-mediated ion channel block. *Neuron* 15 453-462. 10.1016/0896-6273(95)90049-7 7646897.
- [8] Brill J., Huguenard J. R., (2008). Sequential changes in AMPA receptor targeting in the developing neocortical excitatory circuit. *J. Neurosci.* 28 13918-13928. 10.1523/JNEUROSCI.3229-08.2008 19091980.
- [9] Brown P. M. G. E., McGuire H., Bowie D., (2018). Stargazin and cornichon-3 relieve polyamine block of AMPA receptors by enhancing blocker permeation. *J. Gen. Physiol.* 150 67-82. 10.1085/jgp.201711895 29222130.
- [10] Burnashev N., Zhou Z., Neher E., Sakmann B., (1995). Fractional calcium currents through recombinant GluR channels of the NMDA, AMPA and kainate receptor subtypes. *J. Physiol.* 485(Pt 2), 403-418. 7666365.
- [11] Cho C. H., St-Gelais F., Zhang W., Tomita S., Howe J. R., (2007). Two families of TARP isoforms that have distinct effects on the kinetic properties of AMPA receptors and synaptic currents. *Neuron* 55 890-904. 10.1016/j.neuron.2007.08.024 17880893.
- [12] Clem R. L., Huganir R. L., (2010). Calcium-permeable AMPA receptor dynamics mediate fear memory erasure. *Science* 330 1108-1112. 10.1126/science.1195298 21030604.
- [13] Donevan S. D., Rogawski M. A., (1995). Intracellular polyamines mediate inward rectification of Ca<sup>2+</sup>-permeable alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptors. *Proc. Natl. Acad. Sci. U.S.A.* 92 9298-9302. 10.1073/pnas.92.20.9298 7568121.
- [14] Friedman L. K., Pellegrini-Giampietro D. E., Sperber E. F., Bennett M. V., Moshe S. L., Zukin R. S., (1994). Kainate-induced status epilepticus alters glutamate and GABAA receptor gene expression in adult rat hippocampus: an in situ hybridization study. *J. Neurosci.* 14 2697-2707. 10.1523/JNEUROSCI.14-05-02697.1994 8182436.
- [15] Geiger J. R., Melcher T., Koh D. S., Sakmann B., Seeburg P. H., Jonas P., (1995). Relative abundance of subunit mRNAs determines gating and Ca permeability of AMPA receptors in principal neurons and interneurons in rat CNS. *Neuron* 15 193-204. 10.1016/0896-6273(95)90076-4.
- [16] Goel A., Xu L. W., Snyder K. P., Song L., Goenaga-Vazquez Y., Megill A., (2011). Phosphorylation of AMPA receptors is required for sensory deprivation-induced homeostatic synaptic plasticity. *PLoS One* 6:e18264. 10.1371/journal.pone.0018264 21483826.
- [17] Gorter J. A., Petrozzino J. J., Aronica E. M., Rosenbaum D. M., Opitz T., Bennett M. V., (1997). Global ischemia induces downregulation of Glur2 mRNA and increases AMPA receptor-mediated Ca influx in hippocampal CA1 neurons of gerbil. *J. Neurosci.* 17 6179-6188. 10.1523/JNEUROSCI.17-16-06179.1997.
- [18] Gray E. E., Fink A. E., Sarinana J., Vissel B., O'Dell T. J., (2007). Long-term potentiation in the hippocampal CA1 region does not require insertion and activation of GluR2-lacking AMPA receptors. *J. Neurophysiol.* 98 2488-2492. 10.1152/jn.00473.2007 17652419.
- [19] Greger I. H., Watson J. F., Cull-Candy S. G., (2017). Structural and functional architecture of AMPA-Type glutamate receptors and their auxiliary proteins. *Neuron* 94 713-730. 10.1016/j.neuron.2017.04.009 28521126.
- [20] Grooms S. Y., Opitz T., Bennett M. V., Zukin R. S., (2000). Status epilepticus decreases glutamate receptor 2 mRNA and protein expression in hippocampal pyramidal cells before neuronal death. *Proc. Natl. Acad. Sci. U.S.A.* 97 3631-3636. 10.1073/pnas.97.7.3631 10725374.
- [21] Guire E. S., Oh M. C., Soderling T. R., Derkach V. A., (2008). Recruitment of calcium-permeable AMPA receptors during synaptic potentiation is regulated by CaM-kinase I. *J. Neurosci.* 28 6000-6009. 10.1523/JNEUROSCI.0384-08.2008 18524905.
- [22] Hestrin S., (1993). Different glutamate receptor channels mediate fast excitatory synaptic currents in inhibitory and excitatory cortical neurons. *Neuron* 11 1083-1091. 10.1016/0896-6273(93)90221-C 7506044.
- [23] Ho M. T., Pelkey K. A., Topolnik L., Petralia R. S., Takamiya K., Xia J., (2007). Developmental expression of Ca<sup>2+</sup>-permeable AMPA receptors underlies depolarization-induced long-term depression at mossy fiber CA3 pyramid synapses. *J. Neurosci.* 27 11651-11662. 10.1523/JNEUROSCI.2671-07.2007 17959808.
- [24] Huang Y., Doherty J. J., Dingledine R., (2002). Altered histone acetylation at glutamate receptor 2 and brain-derived neurotrophic factor genes is an early event triggered by status epilepticus. *J. Neurosci.* 22 8422-8428. 10.1523/JNEUROSCI.22-19-08422.2002 12351716.

- [25] Hull C., Isaacson J. S., Scanziani M., (2009). Postsynaptic mechanisms govern the differential excitation of cortical neurons by thalamic inputs. *J. Neurosci.* 29 9127–9136. 10.1523/JNEUROSCI.5971-08.2009 19605650.
- [26] Ichikawa M., Maeda T., Handa J., (1997). Ornithine decarboxylase in developing brain of normal and hydrocephalic rats. *Biochemical and immunohistochemical studies. Childs Nerv. Syst.* 13 578–583. 10.1007/s003810050143 9454972.
- [27] Kamboj S. K., Swanson G. T., Cull-Candy S. G., (1995). Intracellular spermine confers rectification on rat calcium-permeable AMPA and kainate receptors. *J. Physiol.* 486(Pt 2), 297–303. 7473197.
- [28] Kato A. S., Siuda E. R., Nisenbaum E. S., Bredt D. S., (2008). AMPA receptor subunit-specific regulation by a distinct family of type II TARPs. *Neuron* 59 986–996. 10.1016/j.neuron.2008.07.034 18817736.
- [29] Keinanen R., Miettinen S., Yrjanheikki J., Koistinaho J., (1997). Induction of ornithine decarboxylase mRNA in transient focal cerebral ischemia in the rat. *Neurosci. Lett.* 239 69–72. 10.1016/S0304-3940(97)00890-2 9469658.
- [30] Kim S., Ziff E. B., (2014). Calcineurin mediates synaptic scaling via synaptic trafficking of Ca<sup>2+</sup>-permeable AMPA receptors. *PLoS Biol.* 12:e1001900. 10.1371/journal.pbio.1001900 24983627.
- [31] Koh D. S., Burnashev N., Jonas P., (1995). Block of native Ca<sup>2+</sup>-permeable AMPA receptors in rat brain by intracellular polyamines generates double rectification. *J. Physiol.* 486(Pt 2), 305–312. 7473198.
- [32] Krauss M., Weiss T., Langnaese K., Richter K., Kowski A., Veh R. W., (2007). Cellular and subcellular rat brain spermidine synthase expression patterns suggest region-specific roles for polyamines, including cerebellar pre-synaptic function. *J. Neurochem.* 103 679–693. 10.1111/j.1471-4159.2007.04770.x 17635671.
- [33] Laube G., Bernstein H. G., Wolf G., Veh R. W., (2002). Differential distribution of spermidine/spermine-like immunoreactivity in neurons of the adult rat brain. *J. Comp. Neurol.* 444 369–386. 10.1002/cne.10157 11891649.
- [34] Liu S. J., Zukin R. S., (2007). Ca<sup>2+</sup>-permeable AMPA receptors in synaptic plasticity and neuronal death. *Trends Neurosci.* 30 126–134. 10.1016/j.tins.2007.01.006 17275103.
- [35] Lu J., Tucciarone J., Lin Y., Huang Z. J., (2014). Input-specific maturation of synaptic dynamics of parvalbumin interneurons in primary visual cortex. *Proc. Natl. Acad. Sci. U.S.A.* 111 16895–16900. 10.1073/pnas.1400694111 25385583.
- [36] Lu Y., Allen M., Halt A. R., Weisenhaus M., Dallapiazza R. F., Hall D. D., (2007). Age-dependent requirement of AKAP150-anchored PKA and GluR2-lacking AMPA receptors in LTP. *EMBO J.* 26 4879–4890. 10.1038/sj.emboj.7601884 17972919.
- [37] Mattison H. A., Bagal A. A., Mohammadi M., Pulimood N. S., Reich C. G., Alger B. E., (2014). Evidence of calcium-permeable AMPA receptors in dendritic spines of CA1 pyramidal neurons. *J. Neurophysiol.* 112 263–275. 10.1152/jn.00578.2013 24760782.
- [38] McCutcheon J. E., Wang X., Tseng K. Y., Wolf M. E., Marinelli M., (2011). Calcium-permeable AMPA receptors are present in nucleus accumbens synapses after prolonged withdrawal from cocaine self-administration but not experimenter-administered cocaine. *J. Neurosci.* 31 5737–5743. 10.1523/JNEUROSCI.0350-11.2011 21490215.
- [39] McGee T. P., Bats C., Farrant M., Cull-Candy S. G., (2015). Auxiliary subunit GSG1L Acts to suppress calcium-permeable AMPA receptor function. *J. Neurosci.* 35 16171–16179. 10.1523/JNEUROSCI.2152-15.2015 26658868.
- [40] Morrison L. D., Kish S. J., (1995). Brain polyamine levels are altered in Alzheimer's disease. *Neurosci. Lett.* 197 5–8. 10.1016/0304-3940(95)11881-V.
- [41] Nilsson T., Bogdanovic N., Volkman I., Winblad B., Folkesson R., Benedikz E., (2006). Altered subcellular localization of ornithine decarboxylase in Alzheimer's disease brain. *Biochem. Biophys. Res. Commun.* 344 640–646. 10.1016/j.bbrc.2006.03.191 16630547.
- [42] Opitz T., Grooms S. Y., Bennett M. V., Zukin R. S., (2000). Remodeling of alpha-amino-3-hydroxy-5-methyl-4-isoxazole-propionic acid receptor subunit composition in hippocampal neurons after global ischemia. *Proc. Natl. Acad. Sci. U.S.A.* 97 13360–13365. 10.1073/pnas.97.24.13360 11087875.
- [43] Paschen W., Csiba L., Rohn G., Bereczki D., (1991). Polyamine metabolism in transient focal ischemia of rat brain. *Brain Res.* 566 354–357. 10.1016/0006-8993(91)91726-H.
- [44] Pellegrini-Giampietro D. E., Zukin R. S., Bennett M. V., Cho S., Pulsinelli W. A., (1992). Switch in glutamate receptor subunit gene expression in CA1 subfield of hippocampus following global ischemia in rats. *Proc. Natl. Acad. Sci. U.S.A.* 89 10499–10503. 10.1073/pnas.89.21.10499 1438239.
- [45] Plant K., Pelkey K. A., Bortolotto Z. A., Morita D., Terashima A., McBain C. J., (2006). Transient incorporation of native GluR2-lacking AMPA receptors during hippocampal long-term potentiation. *Nat. Neurosci.* 9 602–604. 10.1038/nn1678 16582904.
- [46] Reyes A., Lujan R., Rozov A., Burnashev N., Somogyi P., Sakmann B., (1998). Target-cell-specific facilitation and depression in neocortical circuits. *Nat. Neurosci.* 1 279–285. 10.1038/1092 10195160.
- [47] Rozov A., Burnashev N., (1999). Polyamine-dependent facilitation of postsynaptic AMPA receptors counteracts paired-pulse depression. *Nature* 401 594–598. 10.1038/44151 10524627.

- [48] Rozov A., Burnashev N., Sakmann B., Neher E., (2001). Transmitter release modulation by intracellular Ca<sup>2+</sup> + buffers in facilitating and depressing nerve terminals of pyramidal cells in layer 2/3 of the rat neocortex indicates a target cell-specific difference in presynaptic calcium dynamics. *J. Physiol.* 531 807-826. 10.1111/j.1469-7793.2001.0807h.x 11251060.
- [49] Rozov A., Zilberter Y., Wollmuth L. P., Burnashev N., (1998). Facilitation of currents through rat Ca<sup>2+</sup>-permeable AMPA receptor channels by activity-dependent relief from polyamine block. *J. Physiol.* 511(Pt 2),361-377.
- [50] Rozov A., Zivkovic A. R., Schwarz M. K., (2012). Homer1 gene products orchestrate Ca<sup>2+</sup>-permeable AMPA receptor distribution and LTP expression. *Front. Synaptic Neurosci.* 4:4. 10.3389/fnsyn.2012.00004 23133416.
- [51] Sanderson J. L., Gorski J. A., Dell'Acqua M. L., (2016). NMDA receptor-dependent LTD requires transient synaptic incorporation of Ca<sup>2+</sup>-permeable AMPARs mediated by AKAP150-anchored PKA and Calcineurin. *Neuron* 89 1000-1015. 10.1016/j.neuron.2016.01.043 26938443.
- [52] Shanks N. F., Savas J. N., Maruo T., Cais O., Hirao A., Oe S., (2012). Differences in AMPA and kainate receptor interactomes facilitate identification of AMPA receptor auxiliary subunit GSG1L. *Cell Rep.* 1 590-598. 10.1016/j.celrep.2012.05.004 22813734.
- [53] Shin J., Shen F., Huguenard J., (2007). PKC and polyamine modulation of GluR2-deficient AMPA receptors in immature neocortical pyramidal neurons of the rat. *J. Physiol.* 581 679-691. 10.1113/jphysiol.2007.130963 17379631.
- [54] Shin J., Shen F., Huguenard J. R., (2005). Polyamines modulate AMPA receptor-dependent synaptic responses in immature layer v pyramidal neurons. *J. Neurophysiol.* 93 2634-2643. 10.1152/jn.01054.2004 15574796.
- [55] Soares C., Lee K. F., Nassrallah W., Beique J. C., (2013). Differential subcellular targeting of glutamate receptor subtypes during homeostatic synaptic plasticity. *J. Neurosci.* 33 13547-13559. 10.1523/JNEUROSCI.1873-13.2013 23946413.
- [56] Sommer B., Kohler M., Sprengel R., Seeburg P. H., (1991). RNA editing in brain controls a determinant of ion flow in glutamate-gated channels. *Cell* 67 11-19. 10.1016/0092-8674(91)90568-j.
- [57] Soto D., Coombs I. D., Gratacos-Batlle E., Farrant M., Cull-Candy S. G., (2014). Molecular mechanisms contributing to TARP regulation of channel conductance and polyamine block of calcium-permeable AMPA receptors. *J. Neurosci.* 34 11673-11683. 10.1523/JNEUROSCI.0383-14.2014 25164663.
- [58] Soto D., Coombs I. D., Kelly L., Farrant M., Cull-Candy S. G., (2007). Stargazin attenuates intracellular polyamine block of calcium-permeable AMPA receptors. *Nat. Neurosci.* 10 1260-1267. 10.1038/nn1966 17873873.
- [59] Temiz C., Dogan A., Baskaya M. K., Dempsey R. J., (2005). Effect of difluoromethylornithine on reperfusion injury after temporary middle cerebral artery occlusion. *J. Clin. Neurosci.* 12 449-452. 10.1016/j.jocn.2004.05.019 15925780.
- [60] Thiagarajan T. C., Lindskog M., Tsien R. W., (2005). Adaptation to synaptic inactivity in hippocampal neurons. *Neuron* 47 725-737. 10.1016/j.neuron.2005.06.037 16129401.
- [61] Toth K., McBain C. J., (1998). Afferent-specific innervation of two distinct AMPA receptor subtypes on single hippocampal interneurons. *Nat. Neurosci.* 1 572-578. 10.1038/2807 10196564.
- [62] Toth K., Soares G., Lawrence J. J., Philips-Tansey E., McBain C. J., (2000). Differential mechanisms of transmission at three types of mossy fiber synapse. *J. Neurosci.* 20 8279-8289. 10.1523/JNEUROSCI.20-2-08279.2000 11069934.
- [63] Verdoorn T. A., Burnashev N., Monyer H., Seeburg P. H., Sakmann B., (1991). Structural determinants of ion flow through recombinant glutamate receptor channels. *Science* 252 1715-1718. 10.1126/science.1710829 1710829.
- [64] Voinova O., Valiullina F., Zakharova Y., Mukhtarov M., Draguhn A., Rozov A., (2015). Layer specific development of neocortical pyramidal to fast spiking cell synapses. *Front. Cell. Neurosci.* 9:518. 10.3389/fncel.2015.00518.
- [65] Wang H. X., Gao W. J., (2010). Development of calcium-permeable AMPA receptors and their correlation with NMDA receptors in fast-spiking interneurons of rat prefrontal cortex. *J. Physiol.* 588 2823-2838. 10.1113/jphysiol.2010.187591 20547673.
- [66] Watanabe J., Rozov A., Wollmuth L. P., (2005). Target-specific regulation of synaptic amplitudes in the neocortex. *J. Neurosci.* 25 1024-1033. 10.1523/JNEUROSCI.3951-04.2005 15673684.
- [67] Wong-Riley M. T., Jacobs P., (2002). AMPA glutamate receptor subunit 2 in normal and visually deprived macaque visual cortex. *Vis. Neurosci.* 19 563-573. 10.1017/S0952523802195022 12507323.
- [68] Yang Y., Wang X. B., Zhou Q., (2010). Perisynaptic GluR2-lacking AMPA receptors control the reversibility of synaptic and spines modifications. *Proc. Natl. Acad. Sci. U.S.A.* 107 11999-12004. 10.1073/pnas.0913004107 20547835.