

Genome sequences of *Acholeplasma laidlawii* strains with increased resistance to tetracycline and melittin

Chernova O., Chernov V.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 Baranova et al. *Acholeplasma laidlawii* is a well-suited model for studying the molecular basis for adapting mollicutes to environmental conditions. Here, we present the whole-genome sequences of two strains of *A. laidlawii* with increased resistance to tetracycline and melittin.

<http://dx.doi.org/10.1128/genomeA.01446-17>

References

- [1] Blanchard A, Bébéar CM. 2002. Mycoplasmas of humans, p 45-71. In Razin Sh, Herrmann R (ed), Molecular biology and pathogenicity of mycoplasmas. Kluwer Academic Publishers, New York, NY
- [2] Razin S, Hayflick L. 2010. Highlights of mycoplasma research-an historical perspective. *Biologicals* 38:183-190. <https://doi.org/10.1016/j.biologicals.2009.11.008>
- [3] UphoffCC, Drexler HG. 2014. Eradication of mycoplasma contaminations from cell cultures. *Curr Protoc Mol Biol* 106:28.5.1-28.5.12. <https://doi.org/10.1002/0471142727.mb2805s106>
- [4] Waites KB, Lysnyansky I, Bebear CM. 2014. Emerging antimicrobial resistance in mycoplasmas of humans and animals, p 289-322. In Browning GF, Citti C (ed), Mollicutes: molecular biology and pathogenesis. Caister Academic Press, Norfolk, United Kingdom
- [5] Béven L, Castano S, Dufourcq J, Wieslander A, Wróblewski H. 2003. The antibiotic activity of cationic linear amphipathic peptides: lessons from the action of leucine/lysine copolymers on bacteria of the class Mollicutes. *Eur J Biochem* 270:2207-2217. <https://doi.org/10.1046/j.1432-1033.2003.03587.x>
- [6] Lazarev VN, Stipkovits L, Biro J, Miklodi D, Shkarupeta MM, Titova GA, Akopian TA, Govorun VM. 2004. Induced expression of the antimicrobial peptide melittin inhibits experimental infection by *Mycoplasma gallisepticum* in chickens. *Microbes Infect* 6:536-541. <https://doi.org/10.1016/j.micinf.2004.02.006>
- [7] Tully JG. 1979. Special features of the acholeplasmas, p 431-449. In Barile MF, Razin Sh (ed), The mycoplasmas, vol 1. Academic Press, New York, NY
- [8] Lazarev VN, Levitskii SA, Basovskii YI, Chukin MM, Akopian TA, Vereshchagin VV, Kostrjukova ES, Kovaleva GY, Kazanov MD, Malko DB, Vitreschak AG, Sernova NV, Gelfand MS, Demina IA, Serebryakova MV, Galyamina MA, Vtyurin NN, Rogov SI, Alexeev DG, Ladygina VG, Govorun VM. 2011. Complete genome and proteome of *Acholeplasma laidlawii*. *J Bacteriol* 193:4943-4953. <https://doi.org/10.1128/JB.05059-11>
- [9] Medvedeva ES, Baranova NB, Mouzykantov AA, Grigorieva TY, Davydova MN, Trushin MV, Chernova OA, Chernov VM. 2014. Adaptation of mycoplasmas to antimicrobial agents: *Acholeplasma laidlawii* extracellular vesicles mediate the export of ciprofloxacin and a mutant gene related to the antibiotic target. *ScientificWorldJournal* 2014:150615. <https://doi.org/10.1155/2014/150615>
- [10] Medvedeva ES, Davydova MN, Mouzykantov AA, Baranova NB, Grigoreva TY, Siniagina MN, Boulygina EA, Chernova OA, Chernov VM. 2016. Genomic and proteomic profiles of *Acholeplasma laidlawii* strains differing in sensitivity to ciprofloxacin. *Dokl Biochem Biophys* 466:23-27. <https://doi.org/10.1134/S1607672916010075>
- [11] Medvedeva ES, Siniagina MN, Malanin SY, Boulygina EA, Malygina TY, Baranova NB, Mouzykantov AA, Davydova MN, Chernova OA, Chernov VM. 2017. Genome sequences of *Acholeplasma laidlawii* strains differing in sensitivity to ciprofloxacin. *Genome Announc* 5(44):e01189-17. <https://doi.org/10.1128/genomeA.01189-17>
- [12] Sambrook J, Fritsch EF, Maniatis T. 1989. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY

- [13] Nurk S, Bankevich A, Antipov D, Gurevich AA, Korobeynikov A, Lapidus A, Prjibelski AD, Pyshkin A, Sirotkin A, Sirotkin Y, Stepanauskas R, Clingenpeel SR, Woyke T, McLean JS, Lasken R, Tesler G, Alekseyev MA, Pevzner PA. 2013. Assembling single-cell genomes and mini-metagenomes from chimeric MDA products. *J Comput Biol* 20:714-737. <https://doi.org/10.1089/cmb.2013.0084>
- [14] Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie *Nat Methods* 9:357-359. <https://doi.org/10.1038/nmeth.1923>
- [15] Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25:2078-2079. <https://doi.org/10.1093/bioinformatics/btp352>
- [16] Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM. 2012. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff. *Fly* 6:80-92. <https://doi.org/10.4161/fly.19695>
- [17] Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Ciuffo S, Li W. 2013. Prokaryotic genome annotation pipeline. In *The NCBI handbook*, 2nd ed. National Center for Biotechnology Information, Bethesda, MD
- [18] Joo HS, Fu CI, Otto M. 2016. Bacterial strategies of resistance to antimicrobial peptides. *Philos Trans R Soc Lond B Biol Sci* 371:20150292. <https://doi.org/10.1098/rstb.2015.0292>